

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68824 A2

(51) International Patent Classification⁷: C12N 9/02, C07K 14/435, C12N 15/12, 15/10, 15/11, 15/66, A61K 49/00, A01H 5/00, A01K 67/027, A61K 38/17, F41C 3/00, F21S 10/00, A23L 1/00, C12G 1/00, C07K 16/18, C12N 5/10

Pittsburgh, PA 15237 (US). **SZCZEPANIAK, William** [US/US]; 372 South Highland Avenue #203, Pittsburgh, PA 15206 (US).

(21) International Application Number: PCT/US01/08277

(74) Agents: **SEIDMAN, Stephanie, L.** et al.; Heller Ehrman White & McAuliffe LLP, Suite 700, 4250 Executive Square, La Jolla, CA 92037 (US).

(22) International Filing Date: 15 March 2001 (15.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/189,691 15 March 2000 (15.03.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/189,691 (CON)
Filed on 15 March 2000 (15.03.2000)

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **PROL-UME, LTD.** [US/US]; 1085 William Pitt Way, Pittsburgh, PA 15236 (US).

Published:

(71) Applicant and
(72) Inventor: **BRYAN, Bruce, J.** [US/US]; 716 N. Arden Drive, Beverly Hills, CA 90210 (US).

— without international search report and to be republished upon receipt of that report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **SZENT-GY-ORGYI, Christopher** [US/US]; 719 Duncan Avenue,

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/68824 A2

(54) Title: RENILLA RENIFORMIS FLUORESCENT PROTEINS, NUCLEIC ACIDS ENCODING THE FLUORESCENT PROTEINS AND THE USE THEREOF IN DIAGNOSTICS, HIGH THROUGHPUT SCREENING AND NOVELTY ITEMS

(57) Abstract: Isolated and purified nucleic acids encoding green fluorescent proteins from *Renilla reniformis* and the green fluorescent protein encoded thereby are also provided. Mutants of the nucleic acid molecules and the modified encoded proteins are also provided. Compositions and combinations comprising the green fluorescent proteins and/or the luciferase are further provided.

BEST AVAILABLE COPY

-1-

**RENILLA RENIFORMIS FLUORESCENT PROTEINS, NUCLEIC ACIDS ENCODING
THE FLUORESCENT PROTEINS AND THE USE THEREOF IN DIAGNOSTICS,
HIGH THROUGHPUT SCREENING AND NOVELTY ITEMS**

RELATED APPLICATIONS

- 5 Benefit of priority is claimed to U.S. provisional application Serial No. 60/189,691, filed March 15, 2000, to Bryan and Prolume, LTD, entitled "RENILLA RENIFORMIS FLUORESCENT PROTEINS, NUCLEIC ACIDS ENCODING THE FLUORESCENT PROTEINS AND THE USE THEREOF IN DIAGNOSTICS, HIGH THROUGHPUT SCREENING AND NOVELTY ITEMS" is claimed.
- 10 This application is related to allowed U.S. application Serial No. 09/277,716, filed March 26, 1999, to Bruce Bryan and Christopher Szent-Gyorgyi, entitled "LUCIFERASES, FLUORESCENT PROTEINS, NUCLEIC ACIDS ENCODING THE LUCIFERASES AND FLUORESCENT PROTEINS AND THE USE THEREOF IN DIAGNOSTICS, HIGH THROUGHPUT SCREENING AND NOVELTY
- 15 ITEMS." This application is related to International PCT application No. WO 99/49019 to Bruce Bryan and Prolume, LTD., entitled "LUCIFERASES, FLUORESCENT PROTEINS, NUCLEIC ACIDS ENCODING THE LUCIFERASES AND FLUORESCENT PROTEINS AND THE USE THEREOF IN DIAGNOSTICS, HIGH THROUGHPUT SCREENING AND NOVELTY ITEMS."
- 20 This application is also related to subject matter in U.S. application Serial No. 08/757,046, filed November 25, 1996, to Bruce Bryan entitled "BIOLUMINESCENT NOVELTY ITEMS", now U.S. Patent No. 5,876,995, issued March 2, 1999, and in U.S. application Serial No. 08/597,274, filed February 6, 1996, to Bruce Bryan, entitled "BIOLUMINESCENT NOVELTY ITEMS". This
- 25 application is also related to U.S. application Serial No. 08/908,909, filed August 8, 1997, to Bruce Bryan entitled "DETECTION AND VISUALIZATION OF NEOPLASTIC TISSUE AND OTHER TISSUES". The application is also related to U.S. application Serial No. 08/990,103, filed December 12, 1997, to Bruce Bryan entitled "APPARATUS AND METHODS FOR DETECTING AND
- 30 IDENTIFYING INFECTIOUS AGENTS".

Where permitted, the subject matter of each of the above noted U.S. applications and patents herein incorporated by reference in its entirety.


-2-

FIELD OF INVENTION

Provided herein are isolated and purified nucleic acids and encoded fluorescent proteins from *Renilla reniformis* and uses thereof.

BACKGROUND OF THE INVENTION

- 5 Luminescence is a phenomenon in which energy is specifically channeled to a molecule to produce an excited state. Return to a lower energy state is accompanied by release of a photon ($h\nu$). Luminescence includes fluorescence, phosphorescence, chemiluminescence and bioluminescence. Bioluminescence is the process by which living organisms emit light that is visible to other
- 10 organisms. Luminescence may be represented as follows:



- where X^* is an electronically excited molecule and $h\nu$ represents light emission upon return of X^* to a lower energy state. Where the luminescence is
- 15 bioluminescence, creation of the excited state is derived from an enzyme catalyzed reaction. The color of the emitted light in a bioluminescent (or chemiluminescent or other luminescent) reaction is characteristic of the excited molecule, and is independent from its source of excitation and temperature.

- An essential condition for bioluminescence is the use of molecular oxygen, either bound or free in the presence of a luciferase. Luciferases, are
- 20 oxygenases, that act on a substrate, luciferin, in the presence of molecular oxygen and transform the substrate to an excited state. Upon return to a lower energy level, energy is released in the form of light (for reviews see, *e.g.*, McElroy *et al.* (1966) in *Molecular Architecture in Cell Physiology*, Hayashi *et al.*, eds., Prentice-Hall, Inc., Englewood Cliffs, NJ, pp. 63-80; Ward *et al.*, Chapter 7 in *Chemi-and Bioluminescence*, Burr, ed., Marcel Dekker, Inc. NY, pp.321-358; Hastings, J. W. in (1995) *Cell Physiology:Source Book*, N. Sperelakis (ed.), Academic Press, pp. 665-681; *Luminescence, Narcosis and Life in the Deep Sea*, Johnson, Vantage Press, NY, see, esp. pp. 50-56).
- 25

-3-

Though rare overall, bioluminescence is more common in marine organisms than in terrestrial organisms. Bioluminescence has developed from as many as thirty evolutionarily distinct origins and, thus, is manifested in a variety of ways so that the biochemical and physiological mechanisms responsible for

5 bioluminescence in different organisms are distinct. Bioluminescent species span many genera and include microscopic organisms, such as bacteria (primarily marine bacteria including *Vibrio* species), fungi, algae and dinoflagellates, to marine organisms, including arthropods, mollusks, echinoderms, and chordates, and terrestrial organisms including annelid worms

10 and insects.

Assays employing bioluminescence

During the past twenty years, high-sensitivity biochemical assays used in research and in medicine have increasingly employed luminescence and fluorescence rather than radioisotopes. This change has been driven partly by

15 the increasing expense of radioisotope disposal and partly by the need to find more rapid and convenient assay methods. More recently, the need to perform biochemical assays *in situ* in living cells and whole animals has driven researchers toward protein-based luminescence and fluorescence. The uses of firefly luciferase for ATP assays, aequorin and obelin as calcium reporters,

20 *Vargula* luciferase as a neurophysiological indicator, and the *Aequorea* green fluorescent protein as a protein tracer and pH indicator show the potential of bioluminescence-based methods in research laboratories.

Bioluminescence is also beginning to directly impact medicine and biotechnology; for example, *Aequorea* green fluorescent protein (GFP) is

25 employed to mark cells in murine model systems and as a reporter in high throughput drug screening. *Renilla* luciferase is under development for use in diagnostic platforms.

Bioluminescence generating systems

Bioluminescence, as well as other types of chemiluminescence, is used

30 for quantitative determinations of specific substances in biology and medicine. For example, luciferase genes have been cloned and exploited as reporter genes in numerous assays, for many purposes. Since the different luciferase systems

-4-

have different specific requirements, they may be used to detect and quantify a variety of substances. The majority of commercial bioluminescence applications are based on firefly (*Photinus pyralis*) luciferase. One of the first and still widely used assays involves the use of firefly luciferase to detect the presence of ATP.

- 5 It is also used to detect and quantify other substrates or co-factors in the reaction. Any reaction that produces or utilizes NAD(H), NADP(H) or long chain aldehyde, either directly or indirectly, can be coupled to the light-emitting reaction of bacterial luciferase.

- Another luciferase system that has been used commercially for analytical
10 purposes is the *Aequorin* system. The purified jellyfish photoprotein, aequorin, is used to detect and quantify intracellular Ca^{2+} and its changes under various experimental conditions. The *Aequorin* photoprotein is relatively small (~20kDa), nontoxic, and can be injected into cells in quantities adequate to detect calcium over a large concentration range (3×10^{-7} to 10^{-4} M).

- 15 Because of their analytical utility, luciferases and substrates have been studied and well-characterized and are commercially available (e.g., firefly luciferase is available from Sigma, St. Louis, MO, and Boehringer Mannheim Biochemicals, Indianapolis, IN; recombinantly produced firefly luciferase and other reagents based on this gene or for use with this protein are available from
20 Promega Corporation, Madison, WI; the aequorin photoprotein luciferase from jellyfish and luciferase from *Renilla* are commercially available from Sealite Sciences, Bogart, GA; coelenterazine, the naturally-occurring substrate for these luciferases, is available from Molecular Probes, Eugene, OR). These luciferases and related reagents are used as reagents for diagnostics, quality control,
25 environmental testing and other such analyses.

- Because of the utility of luciferases as reagents in analytical systems and the potential for use in high throughput screening systems, there is a need to identify and isolate a variety of luciferases that have improved or different spectral properties compared to those presently available. For all these reasons,
30 it would be advantageous to have luciferases from a variety of species, such as *Gaussia* and various *Renilla* species available.

Fluorescent Proteins

Reporter genes, when co-transfected into recipient cells with a gene of interest, provide a means to detect transfection and other events. Among reporter genes are those that encode fluorescent proteins. The
5 bioluminescence generating systems described herein are among those used as reporter genes. To increase the sensitivity bioluminescence generating systems have been combined with fluorescent compounds and proteins, such as naturally fluorescent phycobiliproteins. Also of interest are the fluorescent proteins that are present in a variety of marine invertebrates, such as the green
10 and blue fluorescent proteins, particularly the green fluorescent protein (GFP) of *Aequorea victoria*.

The green fluorescent proteins (GFP) constitute a class of chromoproteins found only among certain bioluminescent coelenterates. These accessory proteins are fluorescent and function as the ultimate bioluminescence
15 emitter in these organisms by accepting energy from enzyme-bound, excited-state oxyluciferin (*e.g.*, see Ward *et al.* (1979) *J. Biol. Chem.* 254:781-788; Ward *et al.* (1978) *Photochem. Photobiol.* 27:389-396; Ward *et al.* (1982) *Biochemistry* 21:4535-4540).

The best characterized GFPs are those isolated from the jellyfish species
20 *Aequorea*, particularly *Aequorea victoria* (*A. victoria*) and *Aequorea forskålea* (Ward *et al.* (1982) *Biochemistry* 21:4535-4540; Prendergast *et al.* (1978) *Biochemistry* 17:3448-3453). Purified *A. victoria* GFP is a monomeric protein of about 27 kDa that absorbs blue light with excitation wavelength maximum of 395 nm, with a minor peak at 470 nm, and emits green fluorescence with an
25 emission wavelength of about 510 nm and a minor peak near 540 nm (Ward *et al.* (1979) *Photochem. Photobiol. Rev* 4:1-57). This GFP has certain limitations. The excitation maximum of the wildtype GFP is not within the range of wavelengths of standard fluorescein detection optics.

The detection of green fluorescence does not require any exogenous
30 substrates or co-factors. Instead, the high level of fluorescence results from the intrinsic chromophore of the protein. The chromophore includes modified amino acid residues within the polypeptide chain. For example, the fluorescent

-6-

chromophore of *A. victoria* GFP is encoded by the hexapeptide sequence, FSYGVQ, encompassing amino acid residues 64-69. The chromophore is formed by the intramolecular cyclization of the polypeptide backbone at residues Ser65 and Gly67 and the oxidation of the α - β bond of residue Tyr66 (*e.g.*, see

5 Cody *et al.* (1993) *Biochemistry* 32:1212-1218; Shimomura (1978) *FEBS Letters* 104:220-222; Ward *et al.* (1989) *Photochem. Photobiol.* 49:62S). The emission spectrum of the isolated chromophore and the denatured protein at neutral pH do not match the spectrum of the native protein, suggesting that chromophore formation occurs post-translationally (*e.g.*, see Cody *et al.* (1993)

10 *Biochemistry* 32:1212-1218).

In addition, the crystal structure of purified *A. victoria* GFP has been determined (*e.g.*, see Ormö (1996) *Science* 273:1392-1395). The predominant structural features of the protein are an 11-stranded β barrel that forms a nearly perfect cylinder wrapping around a single central α -helix, which contains the

15 modified *p*-hydroxybenzylideneimidazolidinone chromophore. The chromophore is centrally located within the barrel structure and is completely shielded from exposure to bulk solvent.

DNA encoding an isotype of *A. victoria* GFP has been isolated and its nucleotide sequence has been determined (*e.g.*, see Prasher (1992) *Gene*

20 111:229-233). The *A. victoria* cDNA contains a 714 nucleotide open reading frame that encodes a 238 amino acid polypeptide of a calculated M_r of 26,888 Da. Recombinantly expressed *A. victoria* GFPs retain their ability to fluoresce *in vivo* in a wide variety organisms, including bacteria (*e.g.*, see Chalfie *et al.* (1994) *Science* 263:802-805; Miller *et al.* (1997) *Gene* 191:149-153), yeast

25 and fungi (Fey *et al.* (1995) *Gene* 165:127-130; Straight *et al.* (1996) *Curr. Biol.* 6:1599-1608; Cormack *et al.* (1997) *Microbiology* 143:303-311), *Drosophila* (*e.g.*, see Wang *et al.* (1994) *Nature* 369:400-403; Plautz (1996) *Gene* 173:83-87), plants (Heinlein *et al.* (1995); Casper *et al.* (1996) *Gene* 173:69-73), fish (Amsterdam *et al.* (1995)), and mammals (Ikawa *et al.*

30 (1995). *Aequorea* GFP vectors and isolated *Aequorea* GFP proteins have been

-7-

used as markers for measuring gene expression, cell migration and localization, microtubule formation and assembly of functional ion channels (*e.g.*, see Terry *et al.* (1995) *Biochem. Biophys. Res. Commun.* 217:21-27; Kain *et al.* (1995) *Biotechniques* 19:650-655). The *A. victoria* GFP, however, is not ideal for use
5 in analytical and diagnostic processes. Consequently GFP mutants have been selected with the hope of identifying mutants that have single excitation spectral peaks shifted to the red.

In fact a stated purpose in constructing such mutants has been to attempt to make the *A. victoria* GFP more like the GFP from *Renilla*, but which
10 has properties that make it far more ideal for use as an analytical tool. For many practical applications, the spectrum of *Renilla* GFP is preferable to that of the *Aequorea* GFP, because wavelength discrimination between different fluorophores and detection of resonance energy transfer are easier if the component spectra are tall and narrow rather than low and broad (see, U.S.
15 Patent No. 5,625,048). Furthermore, the longer wavelength excitation peak (475 nm) of *Renilla* GFP is almost ideal for fluorescein filter sets and is resistant to photobleaching, but has lower amplitude than the shorter wavelength peak at 395 nm, which is more susceptible to photobleaching (Chalfie *et al.* (1994) *Science* 263:802-805).

20 There exists a phylogenetically diverse and largely unexplored repertoire of bioluminescent proteins that are a reservoir for future development. For these reasons, it would be desirable to have a variety of new luciferases and fluorescent proteins, particularly, *Renilla reniformis* GFP available rather than use mutants of *A. victoria* GFP. Published International PCT application No. WO
25 99/49019 (see, also, allowed U.S. application Serial No. 09/277,716) provides a variety of GFPs including those from *Renilla* species. It remains desirable to have a variety of GFPs and luciferases available in order to optimize systems for particular applications and to improve upon existing methods. Therefore, it is an object herein to provide isolated nucleic acid molecules encoding *Renilla*
30 *reniformis* GFP and the protein encoded thereby. It is also an object herein to provide bioluminescence generating systems that include the luciferases, luciferins, and also include *Renilla reniformis* GFP.

-8-

SUMMARY OF THE INVENTION

Isolated nucleic acid molecules that encode *Renilla reniformis* fluorescent proteins are provided. Nucleic acid probes and primers derived therefrom are also provided. Functionally equivalent nucleic acids, such as those that

5 hybridize under conditions of high stringency to the disclosed molecules and those that have high sequence identity, are also contemplated. Nucleic acid molecules and the encoded proteins are set forth in SEQ ID Nos. 23-27, an exemplary mutin is set forth in SEQ ID No. 33. Also contemplated are nucleic acid molecules that encode the protein set forth in SEQ ID No. 27.

10 Host cells, including bacterial, yeast and mammalian host cells, and plasmids for expression of the nucleic acids encoding the *Renilla reniformis* green fluorescent protein (GFP), are also provided. Combinations of luciferases and the *Renilla reniformis* GFP are also provided.

The genes can be modified by substitution of codons optimized for

15 expression in selected host cells or hosts, such as humans and other mammals, or can be mutagenized to alter the emission properties. Mutations that alter spectral properties are also contemplated.

Such mutations may be identified by substituting each codon with one encoding another amino acid, such as alanine, and determining the effect on the

20 spectral properties of the resulting protein. Particular regions of interest are those in which corresponding the sites mutated in other GFPs, such as Aequora to produce proteins with altered spectral properties are altered.

The *Renilla reniformis* GFP may be used in combination with nucleic acids encoding luciferases, such as those known to those of skill in the art and

25 those that are described in copending allowed U.S. application Serial No. 09/277,716 (see, also, Published International PCT application No. WO 99/49019).

Compositions containing the *Renilla reniformis* GFP or the *Renilla reniformis* GFP and luciferase combination are provided. The compositions can

30 take any of a number of forms, depending on the intended method of use therefor. In certain embodiments, for example, the compositions contain a *Gaussia* luciferase, *Gaussia* luciferase peptide or *Gaussia* luciferase fusion

-9-

protein, formulated for use in luminescent novelty items, immunoassays, donors in FET (fluorescent energy transfer) assays, FRET (fluorescent resonance energy transfer) assays, HTRF (homogeneous time-resolved fluorescence) assays or used in conjunction with multi-well assay devices containing integrated

5 photodetectors, such as those described herein.

The bioluminescence-generating system includes, in addition to the luciferase, a *Renilla reniformis* GFP or mutated form thereof. These compositions can be used in a variety of methods and systems, such as those included in conjunction with diagnostic systems for the *in vivo* detection of

10 neoplastic tissues and other tissues, such as those methods described herein.

Combinations of the *Renilla reniformis* GFP with articles of manufacture to produce novelty items are provided. These novelty items are designed for entertainment, recreation and amusement, and include, but are not limited to: toys, particularly squirt guns, toy cigarettes, toy "Halloween" eggs, footbags

15 and board/card games; finger paints and other paints, slimy play material; textiles, particularly clothing, such as shirts, hats and sports gear suits, threads and yarns; bubbles in bubble making toys and other toys that produce bubbles; balloons; figurines; personal items, such as cosmetics, bath powders, body lotions, gels, powders and creams, nail polishes, make-up, toothpastes and

20 other dentifrices, soaps, body paints, and bubble bath; items such as inks, paper; foods, such as gelatins, icings and frostings; fish food containing luciferins and transgenic fish, particularly transgenic fish that express a luciferase; plant food containing a luciferin or luciferase, preferably a luciferin for use with transgenic plants that express luciferase; and beverages, such as beer,

25 wine, champagne, soft drinks, and ice cubes and ice in other configurations; fountains, including liquid "fireworks" and other such jets or sprays or aerosols of compositions that are solutions, mixtures, suspensions, powders, pastes, particles or other suitable form. The combinations optionally include a bioluminescence generating system. The bioluminescence generating systems

30 can be provided as two compositions: a first composition containing a luciferase and a second composition containing one or more additional components of a bioluminescence generating system.

-10-

Any article of manufacture that can be combined with a bioluminescence-generating system as provided herein and thereby provide entertainment, recreation and/or amusement, including use of the items for recreation or to attract attention, such as for advertising goods and/or services that are associated with a logo or trademark is contemplated herein. Such uses may be in addition to or in conjunction with or in place of the ordinary or normal use of such items. As a result of the combination, the items glow or produce, such as in the case of squirt guns and fountains, a glowing fluid or spray of liquid or particles. The novelty in the novelty item derives from its bioluminescence.

GFPS

Isolated nucleic acids that encode GFP from *Renilla reniformis* are provided herein. Also provided are isolated and purified nucleic acids that encode a component of the bioluminescence generating system and the green fluorescent protein (GFP) (see SEQ ID Nos. 23-27). In particular, nucleic acid molecules that encode *Renilla reniformis* green fluorescent protein (GFPs) and nucleic acid probes and primers derived therefrom are provided. Nucleic acid molecules encoding *Renilla reniformis* GFP are provided (see SEQ ID Nos. 23-26).

Nucleic acid probes and primers containing 14, 16, 30, 100 or more contiguous nucleotides from any of SEQ ID Nos. 23-26 are provided. Nucleic acid probes can be labeled, if needed, for detection, containing at least about 14, preferably at least about 16, or, if desired, 20 or 30 or more, contiguous nucleotides of the sequence of nucleotides encoding the *Renilla reniformis* GFP.

Methods using the probes for the isolation and cloning of GFP-encoding DNA in *Renilla reniformis* are also provided. Vectors containing DNA encoding the *Renilla reniformis* GFP are provided. In particular, expression vectors that contain DNA encoding a *Renilla reniformis* or in operational association with a promoter element that allows for the constitutive or inducible expression of *Renilla reniformis* are provided.

The vectors are capable of expressing the *Renilla reniformis* GFP in a wide variety of host cells. Vectors for producing chimeric *Renilla reniformis*

-11-

GFP/luciferase fusion proteins and/or polycistronic mRNA containing a promoter element and a multiple cloning site located upstream or downstream of DNA encoding *Renilla reniformis* GFP are also provided.

- Recombinant cells containing heterologous nucleic acid encoding a
- 5 *Renilla reniformis* GFP are also provided. Purified *Renilla reniformis* GFP peptides and compositions containing the *Renilla* GFPs and GFP peptides alone or in combination with at least one component of a bioluminescence-generating system, such as a *Renilla* luciferase, are provided. The *Renilla* GFP and GFP peptide compositions can be used, for example, to provide fluorescent
- 10 illumination of novelty items or used in methods of detecting and visualizing neoplastic tissue and other tissues, detecting infectious agents using immunoassays, such as homogeneous immunoassays and *in vitro* fluorescent-based screening assays using multi-well assay devices, or provided in kits for carrying out any of the above-described methods. In particular, these proteins may be
- 15 used in FP (fluorescence polarization) assays, FET (fluorescent energy transfer) assays, FRET (fluorescent resonance energy transfer) assays and HTRF (homogeneous time-resolved fluorescence) assays and also in the BRET assays and sensors provided herein.

- Non-radioactive energy transfer reactions, such as FET or FRET, FP and
- 20 HTRF assays, are homogeneous luminescence assays based on energy transfer and are carried out between a donor luminescent label and an acceptor label (see, e.g., Cardullo *et al.* (1988) *Proc. Natl. Acad. Sci. U.S.A.* 85:8790-8794; Pearce *et al.* (1986) *Proc. Natl. Acad. Sci. U.S.A.* 83:8092-8096; U.S. Patent No. 4,777,128; U.S. Patent No. 5,162,508; U.S. Patent No. 4,927,923; U.S.
- 25 Patent No. 5,279,943; and International PCT Application No. WO 92/01225). Non-radioactive energy transfer reactions using GFPs have been developed (see, International PCT application Nos. WO 98/02571 and WO 97/28261). Non-radioactive energy transfer reactions using GFPs and luciferases, such as a luciferase and its cognate GFP (or multimers thereof), such as in a fusion
- 30 protein, are contemplated herein.

Nucleic acids that exhibit substantial sequence identity with the nucleic acids provided herein are also contemplated. These are nucleic acids that can

-12-

be produced by substituting codons that encode conservative amino acids and also nucleic acids that exhibit at least about 80%, preferably 90 or 95% sequence identity. Sequence identity refers to identity as determined using standard programs with default gap penalties and other defaults as provided by the manufacturer thereof.

The nucleic acids provide an opportunity to produce luciferases and GFPs, which have advantageous application in all areas in which luciferase/luciferins and GFPs have application. The nucleic acids can be used to obtain and produce GFPs and GFPs from other, particularly *Renilla*, species using the probes described herein that correspond to conserved regions. These GFPs have advantageous application in all areas in which GFPs and/or luciferase/luciferins have application. For example, the GFPs provide a means to amplify the output signal of bioluminescence generating systems. *Renilla* GFP has a single excitation absorbance peak in blue light (and around 498 nm) and a predominantly single emission peak around 510 nm (with a small shoulder near 540). This spectrum provides a means for it to absorb blue light and efficiently convert it to green light. This results in an amplification of the output. When used in conjunction with a bioluminescence generating system that yields blue light, such as *Aequorea* or *Renilla* or *Vargula* (*Cypridina*), the output signal for any application, including diagnostic applications, is amplified. In addition, this green light can serve as an energy donor in fluorescence-based assays, such as fluorescence polarization assays, FET (fluorescent energy transfer) assays, FRET (fluorescent resonance energy transfer) assays and HTRF (homogeneous time-resolved fluorescence) assays. Particular assays, herein referred to as BRET (bioluminescence resonance energy transfer assays in which energy is transferred from a bioluminescence reaction of a luciferase to a fluorescent protein), are provided.

Non-radioactive energy transfer reactions, such as FET or FRET, FP and HTRF assays, are homogeneous luminescence assays based on energy transfer that are carried out between a donor luminescent label and an acceptor label (see, e.g., Cardullo *et al.* (1988) *Proc. Natl. Acad. Sci. U.S.A.* 85:8790-8794; Pearce *et al.* (1986) *Proc. Natl. Acad. Sci. U.S.A.* 83:8092-8096; U.S. Patent

-13-

No. 4,777,128; U.S. Patent No. 5,162,508; U.S. Patent No. 4,927,923; U.S. Patent No. 5,279,943; and International PCT Application No. WO 92/01225). Non-radioactive energy transfer reactions using GFPs have been developed (see, International PCT application Nos. WO 98/02571 and WO 97/28261).

- 5 Mutagenesis of the GFPs is contemplated herein, particularly mutagenesis that results in modified GFPs that have red-shifted excitation and emission spectra. The resulting systems have higher output compared to the unmutagenized forms. These GFPs may be selected by random mutagenesis and selection for GFPs with altered spectra or by selected mutagenesis of the
- 10 chromophore region of the GFP.
- The DNA may be introduced as a linear DNA molecule (fragment) or may be included in an expression vector for stable or transient expression of the encoding DNA. In certain embodiments, the cells that contain DNA or RNA encoding a *Renilla* GFP also express the recombinant *Renilla* GFP or polypeptide.
- 15 It is preferred that the cells are selected to express functional GFPs that retain the ability to fluorescence and that are not toxic to the host cell. In some embodiments, cells may also include heterologous nucleic acid encoding a component of a bioluminescence-generating system, preferably a photoprotein or luciferase. In preferred embodiments, the nucleic acid encoding the
- 20 bioluminescence-generating system component is isolated from the species *Aequorea*, *Vargula*, *Pleuromamma*, *Ptilosarcus* or *Renilla*. In more preferred embodiments, the bioluminescence-generating system component is a *Renilla reniformis* luciferase or *mulleri* including the amino acid sequence set forth in SEQ ID No. 18 or the *Pleuromamma* luciferase set forth in SEQ ID No. 28, or
- 25 the *Gaussia* luciferase set forth in SEQ ID No. 19.

The GFPs provided herein may be used in combination with any suitable bioluminescence generating system, but is preferably used in combination with a *Renilla* or *Aequorea*, *Pleuromamma* or *Gaussia* luciferase.

-14-

Purified *Renilla* GFPs, particularly purified *Renilla reniformis* GFP peptides are provided. Presently preferred *Renilla* GFP for use in the compositions herein is *Renilla reniformis* GFP including the sequence of amino acids set forth above and in the Sequence Listing.

- 5 Fusions of the nucleic acid, particularly DNA, encoding *Renilla* GFP with DNA encoding a luciferase are also provided herein.

The cells that express functional luciferase and/or GFP, which may be used alone or in conjunction with a bioluminescence-generating system, in cell-based assays and screening methods, such as those described herein.

- 10 Presently preferred host cells for expressing GFP and luciferase are bacteria, yeasts, fungi, plant cells, insect cells and animal cells.

- The luciferases and GFPs or cells that express them also may be used in methods of screening for bacterial contamination and methods of screening for metal contaminants. To screen for bacterial contamination, bacterial cells that
- 15 express the luciferase and/or GFP are put in autoclaves or in other areas in which testing is contemplated. After treatment or use of the area, the area is tested for the presence of glowing bacteria. Presence of such bacteria is indicative of a failure to eradicate other bacteria. Screening for heavy metals and other environmental contaminants can also be performed with cells that
- 20 contain the nucleic acids provided herein, if expression is linked to a system that is dependent upon the particular heavy metal or contaminant.

- The systems and cells provided herein can be used for high throughout screening protocols, intracellular assays, medical diagnostic assays, environmental testing, such as tracing bacteria in water supplies, in conjunction
- 25 with enzymes for detecting heavy metals, in spores for testing autoclaves in hospital, foods and industrial autoclaves. Non-pathogenic bacteria containing the systems can be included in feed to animals to detect bacterial contamination in animal products and in meats.

- Compositions containing a *Renilla* GFP are provided. The compositions
- 30 can take any of a number of forms, depending on the intended method of use therefor. In certain embodiments, for example, the compositions contain a *Renilla* GFP or GFP peptide, preferably *Renilla mulleri* GFP or *Renilla reniformis*

-15-

GFP peptide, formulated for use in luminescent novelty items, immunoassays, FET (fluorescent energy transfer) assays, FRET (fluorescent resonance energy transfer) assays, HTRF (homogeneous time-resolved fluorescence) assays or used in conjunction with multi-well assay devices containing integrated
5 photodetectors, such as those described herein. In other instances, the GFPs are used in beverages, foods or cosmetics.

Compositions that contain a *Renilla reniformis* GFP or GFP peptide and at least one component of a bioluminescence-generating system, preferably a luciferase, luciferin or a luciferase and a luciferin, are provided. In preferred
10 embodiments, the luciferase/luciferin bioluminescence- generating system is selected from those isolated from: an insect system, a coelenterate system, a ctenophore system, a bacterial system, a mollusk system, a crustacea system, a fish system, an annelid system, and an earthworm system. Bioluminescence-generating systems include those isolated from *Renilla*, *Aequorea*, and *Vargula*,
15 *Gaussia* and *Pleuromamma*.

Combinations containing a first composition containing a *Renilla reniformis* GFP or *Ptilosarcus* GFP or mixtures thereof and a second composition containing a bioluminescence- generating system for use with inanimate articles of manufacture to produce novelty items are provided. These novelty items,
20 which are articles of manufacture, are designed for entertainment, recreation and amusement, and include, but are not limited to: toys, particularly squirt guns, toy cigarettes, toy "Halloween" eggs, footbags and board/card games; finger paints and other paints, slimy play material; textiles, particularly clothing, such as shirts, hats and sports gear suits, threads and yarns; bubbles in bubble
25 making toys and other toys that produce bubbles; balloons; figurines; personal items, such as bath powders, body lotions, gels, powders and creams, nail polishes, cosmetics including make-up, toothpastes and other dentifrices, soaps, cosmetics, body paints, and bubble bath, bubbles made from non-detergent sources, particularly proteins such as albumin and other non-toxic proteins; in
30 fishing lures and glowing transgenic worms, particularly crosslinked polyacrylamide containing a fluorescent protein and/or components of a bioluminescence generating system, which glow upon contact with water; items

-16-

such as inks, paper; foods, such as gelatins, icings and frostings; fish food containing luciferins and transgenic animals, such as transgenic fish, worms, monkeys, rodents, ungulates, ovine, ruminants and others, that express a luciferase and/or *Renilla reniformis* GFP; transgenic worms that express *Renilla*
5 *reniformis* GFP and are used as lures; plant food containing a luciferin or luciferase, preferably a luciferin for use with transgenic plants that express luciferase and *Renilla reniformis* GFP, transgenic plants that express *Renilla reniformis* GFP, particularly ornamental plants, such as orchids, roses, and other plants with decorative flowers; transgenic plants and animals in which the
10 *Renilla reniformis* GFP is a marker for tracking introduction of other genes; and beverages, such as beer, wine, champagne, soft drinks, milk and ice cubes and ice in other configurations containing *Renilla reniformis* GFP; fountains, including liquid "fireworks" and other such jets or sprays or aerosols of compositions that are solutions, mixtures, suspensions, powders, pastes, particles or other
15 suitable forms.

Any article of manufacture that can be combined with a bioluminescence-generating system and *Renilla reniformis* GFP or with just a *Renilla reniformis* GFP, as provided herein, that thereby provides entertainment, recreation and/or amusement, including use of the items for recreation or to
20 attract attention, such as for advertising goods and/or services that are associated with a logo or trademark is contemplated herein. Such uses may be in addition to or in conjunction with or in place of the ordinary or normal use of such items. As a result of the combination, the items glow or produce, such as in the case of squirt guns and fountains, a glowing fluid or spray of liquid or
25 particles.

Methods for diagnosis and visualization of tissues *in vivo* or *in situ* using compositions containing a *Renilla reniformis* GFP and/or a *Renilla reniformis* or *mulleri* luciferase or others of the luciferases and/or GFPs provided herein are provided. For example, the *Renilla reniformis* GFP protein can be used in
30 conjunction with diagnostic systems that rely on bioluminescence for visualizing tissues *in situ*. The systems are particularly useful for visualizing and detecting neoplastic tissue and specialty tissue, such as during non-invasive and invasive

-17-

procedures. The systems include compositions containing conjugates that include a tissue specific, particularly a tumor-specific, targeting agent linked to a targeted agent, a *Renilla reniformis* GFP, a luciferase or luciferin. The systems also include a second composition that contains the remaining components of a bioluminescence generating reaction and/or the *Renilla reniformis* GFP. In some embodiments, all components, except for activators, which are provided *in situ* or are present in the body or tissue, are included in a single composition.

Methods for diagnosis and visualization of tissues *in vivo* or *in situ* using compositions containing a *Gaussia* luciferase are provided. For example, the *Gaussia* luciferase or *Gaussia* luciferase peptide can be used in conjunction with diagnostic systems that rely on bioluminescence for visualizing tissues *in situ*. The systems are particularly useful for visualizing and detecting neoplastic tissue and specialty tissue, such as during non-invasive and invasive procedures. The systems include compositions containing conjugates that include a tissue specific, particularly a tumor-specific, targeting agent linked to a targeted agent, a *Gaussia* luciferase, a GFP or luciferin. The systems also include a second composition that contains the remaining components of a bioluminescence generating reaction and/or the *Gaussia* luciferase. In some embodiments, all components, except for activators, which are provided *in situ* or are present in the body or tissue, are included in a single composition.

In particular, the diagnostic systems include two compositions. A first composition that contains conjugates that, in preferred embodiments, include antibodies directed against tumor antigens conjugated to a component of the bioluminescence generating reaction, a luciferase or luciferin, preferably a luciferase are provided. In certain embodiments, conjugates containing tumor-specific targeting agents are linked to luciferases or luciferins. In other embodiments, tumor-specific targeting agents are linked to microcarriers that are coupled with, preferably more than one of the bioluminescence generating components, preferably more than one luciferase molecule.

The second composition contains the remaining components of a bioluminescence generating system, typically the luciferin or luciferase substrate. In some embodiments, these components, particularly the luciferin

-18-

are linked to a protein, such as a serum albumin, or other protein carrier. The carrier and time release formulations, permit systemically administered components to travel to the targeted tissue without interaction with blood cell components, such as hemoglobin that deactivates the luciferin or luciferase.

- 5 Methods for diagnosing diseases, particularly infectious diseases, using chip methodology (see, *e.g.*, copending U.S. application Serial No. 08/990,103) a luciferase/luciferin bioluminescence-generating system and a *Renilla reniformis* GFP are provided. In particular, the chip includes an integrated photodetector that detects the photons emitted by the bioluminescence-generating system,
10 particularly using luciferase encoded by the nucleic acids provided herein and/or *Renilla reniformis* GFP.

- In one embodiment, the chip is made using an integrated circuit with an array, such as an X-Y array, of photodetectors. The surface of circuit is treated to render it inert to conditions of the diagnostic assays for which the chip is
15 intended, and is adapted, such as by derivatization for linking molecules, such as antibodies. A selected antibody or panel of antibodies, such as an antibody specific for a bacterial antigen, is affixed to the surface of the chip above each photodetector. After contacting the chip with a test sample, the chip is contacted with a second antibody linked to a *Renilla* GFP, a chimeric antibody-
20 *Renilla* GFP fusion protein or an antibody linked to a component of a bioluminescence generating system, such as a luciferase or luciferin, that are specific for the antigen. The remaining components of the bioluminescence generating reaction are added, and, if any of the antibodies linked to a component of a bioluminescence generating system are present on the chip,
25 light will be generated and detected by the adjacent photodetector. The photodetector is operatively linked to a computer, which is programmed with information identifying the linked antibodies, records the event, and thereby identifies antigens present in the test sample.

-19-

Methods for generating chimeric GFP fusion proteins are provided. The methods include linking DNA encoding a gene of interest, or portion thereof, to DNA encoding a GFP coding region in the same translational reading frame. The encoded-protein of interest may be linked in-frame to the amino- or carboxyl-terminus of the GFP. The DNA encoding the chimeric protein is then linked in operable association with a promoter element of a suitable expression vector. Alternatively, the promoter element can be obtained directly from the targeted gene of interest and the promoter-containing fragment linked upstream of the GFP coding sequence to produce chimeric GFP proteins or to produce polycistronic mRNAs that encode the *Renilla reniformis* GFP and a luciferase, preferably a *Renilla* luciferase, more preferably *Renilla reniformis* luciferase.

Methods for identifying compounds using recombinant cells that express heterologous DNA encoding a *Renilla reniformis* GFP under the control of a promoter element of a gene of interest are provided. The recombinant cells can be used to identify compounds or ligands that modulate the level of transcription from the promoter of interest by measuring *Renilla reniformis* GFP-mediated fluorescence. Recombinant cells expressing the chimeric *Renilla reniformis* GFP or polycistronic mRNA encoding *Renilla reniformis* and a luciferase, may also be used for monitoring gene expression or protein trafficking, or determining the cellular localization of the target protein by identifying localized regions of GFP-mediated fluorescence within the recombinant cell.

Other assays using the GFPs and/or luciferases are contemplated herein. Any assay or diagnostic method known used by those of skill in the art that employ *Aequora* GFPs and/or other luciferases are contemplated herein.

Kits containing the GFPs for use in the methods, including those described herein, are provided. In one embodiment, the kits containing an article of manufacture and appropriate reagents for generating bioluminescence are provided. The kits containing such soap compositions, with preferably a

-20-

moderate pH (between 5 and 8) and bioluminescence generating reagents, including luciferase and luciferin and the GFP are provided herein. These kits, for example, can be used with a bubble-blowing or producing toy. These kits can also include a reloading or charging cartridge or can be used in connection
5 with a food.

In another embodiment, the kits are used for detecting and visualizing neoplastic tissue and other tissues and include a first composition that contains the GFP and at least one component of a bioluminescence generating system, and a second that contains the activating composition, which contains the
10 remaining components of the bioluminescence generating system and any necessary activating agents.

Thus, these kits will typically include two compositions, a first composition containing the GFP formulated for systemic administration (or in some embodiments local or topical application), and a second composition
15 containing the components or remaining components of a bioluminescence generating system, formulated for systemic, topical or local administration depending upon the application. Instructions for administration will be included.

In other embodiments, the kits are used for detecting and identifying diseases, particularly infectious diseases, using multi-well assay devices and
20 include a multi-well assay device containing a plurality of wells, each having an integrated photodetector, to which an antibody or panel of antibodies specific for one or more infectious agents are attached, and composition containing a secondary antibody, such as an antibody specific for the infectious agent that is linked to a *Renilla reniformis* GFP protein, a chimeric antibody-*Renilla reniformis*
25 GFP fusion protein or F(Ab)₂ antibody fragment-*Renilla reniformis* GFP fusion protein. A second composition contains a bioluminescence generating system that emits a wavelength of light within the excitation range of the *Renilla mulleri* GFP, such as species of *Renilla* or *Aequorea*, for exciting the *Renilla reniformis*, which produces light that is detected by the photodetector of the device to
30 indicate the presence of the agent.

-21-

As noted above, fusions of nucleic acid encoding the luciferases and or GFPs provided herein with other luciferases and GFPs are provided. Of particular interest are fusions that encode pairs of luciferases and GFPs, such as a *Renilla* luciferase and a *Renilla* GFP (or a homodimer or other multiple of a *Renilla* GFP). The luciferase and GFP bind and in the presence of a luciferin will produced fluorescence that is red shifted compared to the luciferase in the absence of the GFP. This fusion or fusions in which the GFP and luciferase are linked via a target, such as a peptide, can be used as a tool to assess anything that interacts with the linker.

Mutants of the GFPs and luciferases are provided. Of particular interest are mutants, such as temperature sensitive mutants, of the GFP and luciferases that alter their interaction, such as mutations in the *Renilla* luciferase and *Renilla* GFP that alters their interaction at a critical temperature.

Antibodies, polyclonal and monoclonal antibodies that specifically bind to any of the proteins encoded by the nucleic acids provided herein are also provided. These antibodies, monoclonal or polyclonal, can be prepared employing standard techniques, known to those of skill in the art. In particular, immunoglobulins or antibodies obtained from the serum of an animal immunized with a substantially pure preparation of a luciferase or GFP provided herein or an or epitope-containing fragment thereof are provided. Monoclonal antibodies are also provided. The immunoglobulins that are produced have, among other properties, the ability to specifically and preferentially bind to and/or cause the immunoprecipitation of a GFP or luciferase, particularly a *Renilla* or *Ptilosarcus* GFP or a *Pleuromamma*, *Gaussia* or *Renilla mulleri* luciferase, that may be present in a biological sample or a solution derived from such a biological sample.

DESCRIPTION OF THE FIGURES

FIGURE 1 depicts phylogenetic relationships among the anthozoan GFPs.

FIGURES 2A-D illustrate the underlying principle of Bioluminescent Resonance Energy Transfer (BRET) and its use as sensor: A) in isolation, a luciferase, preferably an anthozoan luciferase, emits blue light from the coelenterazine-derived chromophore; B) in isolation, a GFP, preferably an

RECTIFIED SHEET (RULE91)
ISA/EP

-22-

anthozoan GFP that binds to the luciferase, that is excited with blue-green light emits green light from its integral peptide based fluorophore; C) when the luciferase and GFP associate as a complex *in vivo* or *in vitro*, the luciferase non-radiatively transfers its reaction energy to the GFP fluorophore, which then emits the green light; D) any molecular interaction that disrupts the luciferase-GFP complex can be quantitatively monitored by observing the spectral shift from green to blue light.

FIGURE 3 illustrates exemplary BRET sensor architecture.

FIGURE 4 depicts the substitution of altered fluorophores into the background of *Ptilosarcus*, *Renilla mulleri* and *Renilla reniformis* GFPs (the underlined regions correspond to amino acids 56-75 of SEQ ID No. 27 *Renilla reniformis* GFP; amino acids 59-78 of SEQ ID No. 16 *Renilla mulleri* GFP; and amino acids 59-78 of SEQ ID No. 32 for *Ptilosarcus* GFP).

FIGURE 5 depicts the three anthozoan fluorescent proteins for which a crystal structure exists; another is available commercially from Clontech as dsRed (from *Discosoma striata*; also known as drFP583, as in this alignment); a dark gray background depicts amino acid conservation, and a light gray background depicts shared physicochemical properties.

FIGURE 6 compares the sequences of a variety of GFPs, identifying sites for mutation to reduce multimerization; abbreviations are as follows: *Amemonia majona* is amFP486; *Zoanthus* sp. zFP506 and zFP538; *Discosoma* sp. "red" is drFP583; *Clavularia* sp. is cFP484; and the GFP from the anthozoal *A. sulcata* is designated FP595.

DETAILED DESCRIPTION OF THE INVENTION

- 25 A. DEFINITIONS
- B. Fluorescent Proteins
 - 1. Green and blue fluorescent proteins
 - 2. *Renilla reniformis* GFP
- C. BIOLUMINESCENCE GENERATING SYSTEMS AND COMPONENTS

-23-

1. General description
 - a. Luciferases
 - b. Luciferins
 - c. Activators
 - d. Reactions
2. The *Renilla* system
3. Ctenophore systems
4. The aequorin system
 - a. Aequorin and related photoproteins
 - b. Luciferin
5. Crustacean, particularly *Cypridina* systems
 - a. *Vargula* luciferase
 - (1) Purification from *Cypridina*
 - (2) Preparation by Recombinant Methods
 - b. *Vargula* luciferin
 - c. Reaction
6. Insect bioluminescent systems including fireflies, click beetles, and other insect system
 - a. Luciferase
 - b. Luciferin
 - c. Reaction
7. Other systems
 - a. Bacterial systems
 - (1) Luciferases
 - (2) Luciferins
 - (3) Reactions
 - b. Dinoflagellate bioluminescence generating systems
- D. ISOLATION AND IDENTIFICATION OF NUCLEIC ACIDS ENCODING LUCIFERASES AND GFPs
 1. Isolation of specimens of the genus *Renilla*
 2. Preparation of *Renilla* cDNA expression libraries
 - a. RNA isolation and cDNA synthesis
 - b. Construction of cDNA expression libraries
 3. Cloning of *Renilla reniformis* Green Fluorescent Protein
 4. Isolation and identification of DNA encoding *Renilla mulleri* GFP
 5. Isolation and identification of DNA encoding *Renilla mulleri* luciferase
- E. RECOMBINANT EXPRESSION OF PROTEINS
 1. DNA encoding *Renilla* proteins

-24-

2. DNA constructs for recombinant production of *Renilla reniformis* and other proteins
 3. Host organisms for recombinant production of *Renilla* proteins
 4. Methods for recombinant production of *Renilla* proteins
 5. Recombinant cells expressing heterologous nucleic acid encoding luciferases and GFPs
- 5
- F. COMPOSITIONS AND CONJUGATES
1. *Renilla* GFP compositions
 2. *Renilla* luciferase compositions
 3. Conjugates
 - a. Linkers
 - b. Targeting Agents
 - c. Anti-tumor Antigen Antibodies
 - d. Preparation of the conjugates
 4. Formulation of the compositions for use in the diagnostic systems
 - a. The first composition: formulation of the conjugates
 - b. The second composition
 - c. Practice of the reactions in combination with targeting agents
- 10
- 15
- 20
- G. COMBINATIONS
- H. Exemplary uses of *Renilla reniformis* GFPs and encoding nucleic acid molecules
1. Methods for diagnosis of neoplasms and other tissues
 2. Methods of diagnosing diseases
 3. Methods for generating *Renilla mulleri* luciferase, *Pleuromamma* luciferase and *Gaussia* luciferase fusion proteins with *Renilla reniformis* GFP
 4. Cell-based assays for identifying compounds
- 25
- 30
- I. KITS
- J. Muteins
1. Mutation of GFP surfaces to disrupt multimerization
 2. Use of advantageous GFP surfaces with substituted fluorophores
- 35
- K. Transgenic plants and animals
- L. Bioluminescence Resonance Energy Transfer (BRET) System
1. Design of sensors based on BRET
 2. BRET Sensor Architectures
 3. Advantages of BRET sensors

-25-

A. DEFINITIONS

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which this invention belongs. All patents, applications and publications of
5 referred to throughout the disclosure are incorporated by reference in their entirety.

As used herein, chemiluminescence refers to a chemical reaction in which energy is specifically channeled to a molecule causing it to become electronically excited and subsequently to release a photon thereby emitting visible light.
10 Temperature does not contribute to this channeled energy. Thus, chemiluminescence involves the direct conversion of chemical energy to light energy.

As used herein, luminescence refers to the detectable EM radiation, generally, UV, IR or visible EM radiation that is produced when the excited
15 product of an exergic chemical process reverts to its ground state with the emission of light. Chemiluminescence is luminescence that results from a chemical reaction. Bioluminescence is chemiluminescence that results from a chemical reaction using biological molecules (or synthetic versions or analogs thereof) as substrates and/or enzymes.

As used herein, bioluminescence, which is a type of chemiluminescence, refers to the emission of light by biological molecules, particularly proteins. The essential condition for bioluminescence is molecular oxygen, either bound or free
20 in the presence of an oxygenase, a luciferase, which acts on a substrate, a luciferin. Bioluminescence is generated by an enzyme or other protein (luciferase) that is an oxygenase that acts on a substrate luciferin (a
25 bioluminescence substrate) in the presence of molecular oxygen and transforms the substrate to an excited state, which upon return to a lower energy level releases the energy in the form of light.

As used herein, the substrates and enzymes for producing
30 bioluminescence are generically referred to as luciferin and luciferase, respectively. When reference is made to a particular species thereof, for clarity,

-26-

each generic term is used with the name of the organism from which it derives, for example, bacterial luciferin or firefly luciferase.

As used herein, luciferase refers to oxygenases that catalyze a light emitting reaction. For instance, bacterial luciferases catalyze the oxidation of
5 flavin mononucleotide (FMN) and aliphatic aldehydes, which reaction produces light. Another class of luciferases, found among marine arthropods, catalyzes the oxidation of *Cypridina* (*Vargula*) luciferin, and another class of luciferases catalyzes the oxidation of *Coleoptera* luciferin.

Thus, luciferase refers to an enzyme or photoprotein that catalyzes a
10 bioluminescent reaction (a reaction that produces bioluminescence). The luciferases, such as firefly and *Gaussia* and *Renilla* luciferases, that are enzymes which act catalytically and are unchanged during the bioluminescence generating reaction. The luciferase photoproteins, such as the aequorin photoprotein to which luciferin is non-covalently bound, are changed, such as by release of the
15 luciferin, during bioluminescence generating reaction. The luciferase is a protein that occurs naturally in an organism or a variant or mutant thereof, such as a variant produced by mutagenesis that has one or more properties, such as thermal stability, that differ from the naturally-occurring protein. Luciferases and modified mutant or variant forms thereof are well known. For purposes herein,
20 reference to luciferase refers to either the photoproteins or luciferases.

Thus, reference, for example, to "*Gaussia* luciferase" means an enzyme isolated from member of the genus *Gaussia* or an equivalent molecule obtained from any other source, such as from another related copepod, or that has been prepared synthetically. It is intended to encompass *Gaussia*
25 luciferases with conservative amino acid substitutions that do not substantially alter activity. Suitable conservative substitutions of amino acids are known to those of skill in this art and may be made generally without altering the biological activity of the resulting molecule. Those of skill in this art recognize that, in general, single amino acid substitutions in non-essential regions of a polypeptide
30 do not substantially alter biological activity (see, *e.g.*, Watson *et al. Molecular Biology of the Gene*, 4th Edition, 1987, The Bejacmin/Cummings Pub. co., p.224).

-27-

"*Renilla* GFP" refers to GFPs from the genus *Renilla* and to mutants or variants thereof. It is intended to encompass *Renilla* GFPs with conservative amino acid substitutions that do not substantially alter activity and physical properties, such as the emission spectra and ability to shift the spectral output of bioluminescence generating systems.

Such substitutions are preferably made in accordance with those set forth in TABLE 1 as follows:

TABLE 1		
	Original residue	Conservative substitution
10	Ala (A)	Gly; Ser
	Arg (R)	Lys
	Asn (N)	Gln; His
	Cys (C)	Ser
	Gln (Q)	Asn
15	Glu (E)	Asp
	Gly (G)	Ala; Pro
	His (H)	Asn; Gln
	Ile (I)	Leu; Val
	Leu (L)	Ile; Val
20	Lys (K)	Arg; Gln; Glu
	Met (M)	Leu; Tyr; Ile
	Phe (F)	Met; Leu; Tyr
	Ser (S)	Thr
	Thr (T)	Ser
25	Trp (W)	Tyr
	Tyr (Y)	Trp; Phe
	Val (V)	Ile; Leu

Other substitutions are also permissible and may be determined empirically or in accord with known conservative substitutions.

The luciferases and luciferin and activators thereof are referred to as bioluminescence generating reagents or components. Typically, a subset of these reagents will be provided or combined with an article of manufacture. Bioluminescence will be produced upon contacting the combination with the remaining reagents. Thus, as used herein, the component luciferases, luciferins, and other factors, such as O₂, Mg²⁺, Ca²⁺ are also referred to as bioluminescence generating reagents (or agents or components).

As used herein, a *Renilla reniformis* green fluorescent protein (GFP) refers to a fluorescent protein that is encoded by a sequence of nucleotides that

-28-

encodes the protein of SEQ ID No. 27 or to a green fluorescent protein from *Renilla reniformis* having at least 80%, 90% or 95% or greater sequence identity thereto; or that is encoded by a sequence of nucleotides that hybridizes under high stringency along its full length to the coding portion of the sequence of nucleotides set forth in any of SEQ ID Nos. 23-25. A *Renilla reniformis* GFP is protein that is fluorescent and is produced in a *Renilla reniformis*.

As used herein, bioluminescence substrate refers to the compound that is oxidized in the presence of a luciferase, and any necessary activators, and generates light. These substrates are referred to as luciferins herein, are substrates that undergo oxidation in a bioluminescence reaction. These bioluminescence substrates include any luciferin or analog thereof or any synthetic compound with which a luciferase interacts to generate light. Preferred substrates are those that are oxidized in the presence of a luciferase or protein in a light-generating reaction. Bioluminescence substrates, thus, include those compounds that those of skill in the art recognize as luciferins. Luciferins, for example, include firefly luciferin, *Cypridina* (also known as *Vargula*) luciferin (coelenterazine), bacterial luciferin, as well as synthetic analogs of these substrates or other compounds that are oxidized in the presence of a luciferase in a reaction the produces bioluminescence.

As used herein, capable of conversion into a bioluminescence substrate means susceptible to chemical reaction, such as oxidation or reduction, that yields a bioluminescence substrate. For example, the luminescence producing reaction of bioluminescent bacteria involves the reduction of a flavin mononucleotide group (FMN) to reduced flavin mononucleotide (FMNH₂) by a flavin reductase enzyme. The reduced flavin mononucleotide (substrate) then reacts with oxygen (an activator) and bacterial luciferase to form an intermediate peroxy flavin that undergoes further reaction, in the presence of a long-chain aldehyde, to generate light. With respect to this reaction, the reduced flavin and the long chain aldehyde are substrates.

As used herein, a bioluminescence generating system refers to the set of reagents required to conduct a bioluminescent reaction. Thus, the specific luciferase, luciferin and other substrates, solvents and other reagents that may

-29-

be required to complete a bioluminescent reaction form a bioluminescence system. Thus a bioluminescence generating system refers to any set of reagents that, under appropriate reaction conditions, yield bioluminescence. Appropriate reaction conditions refers to the conditions necessary for a bioluminescence
5 reaction to occur, such as pH, salt concentrations and temperature. In general, bioluminescence systems include a bioluminescence substrate, luciferin, a luciferase, which includes enzymes luciferases and photoproteins, and one or more activators. A specific bioluminescence system may be identified by reference to the specific organism from which the luciferase derives; for
10 example, the *Vargula* (also called *Cypridina*) bioluminescence system (or *Vargula* system) includes a *Vargula* luciferase, such as a luciferase isolated from the ostracod, *Vargula* or produced using recombinant means or modifications of these luciferases. This system would also include the particular activators necessary to complete the bioluminescence reaction, such as oxygen and a
15 substrate with which the luciferase reacts in the presence of the oxygen to produce light.

The luciferases provided herein may be incorporated into bioluminescence generating systems and used, as appropriate, with the GFPs provided herein or with other GFPs. Similarly, the GFPs provided herein may be used with known
20 bioluminescence generating systems.

As used herein, the amino acids, which occur in the various amino acid sequences appearing herein, are identified according to their well-known, three-letter or one-letter abbreviations. The nucleotides, which occur in the various DNA molecules, are designated with the standard single-letter
25 designations used routinely in the art.

As used herein, a fluorescent protein refers to a protein that possesses the ability to fluoresce (*i.e.*, to absorb energy at one wavelength and emit it at another wavelength). These proteins can be used as a fluorescent label or marker and in any applications in which such labels would be used, such as
30 immunoassays, CRET, FRET, and FET assays, and in the assays designated herein as BRET assays. For example, a green fluorescent protein refers to a polypeptide that has a peak in the emission spectrum at about 510 nm.

-30-

As used herein, the term BRET (Bioluminescence Resonance Energy Transfer) refers to non-radiative luciferase-to-FP energy transfer. It differs from (Fluorescence Resonance Energy Transfer), which refers to energy transfer between chemical fluors.

- 5 As used herein, a BRET system refers the combination of a FP, in this case *Renilla reniformis* GFP and a luciferase for resonance energy transfer. BRET refers to any method in which the luciferase is used to generate the light upon reaction with a luciferin which is then non-radiatively transferred to a FP. The energy is transferred to a FP, particularly a GFP, which focuses and shifts the
- 10 energy and emits it at a different wavelength. In preferred embodiments, the BRET system includes a bioluminescence generating system and a *Renilla reniformis* GFP. The bioluminescence generating system is preferably a *Renilla* system. Hence, the preferred pair is a *Renilla* luciferase and a *Renilla* GFP, which specifically interact. Alterations in the binding will be reflected in changes in the
- 15 emission spectra of light produced by the luciferase. As a result the pair can function as a sensor of external events.

As used herein, a biosensor (or sensor) refers to a BRET system for use to detect alterations in the environment *in vitro* or *in vivo* in which the BRET system is used.

- 20 As used herein, modulator with reference to a BRET system refers to a molecule or molecules that undergo a conformation change in response to interaction with another molecule thereby affecting the proximity and/or orientation of the GFP and luciferase in the BRET system. Modulators include, but are not limited to, a protease site, a second messenger binding site, an ion
- 25 binding molecule, a receptor, an oligomer, an enzyme substrate, a ligand, or other such binding molecule. If the GFP and luciferase are each linked to the modulator, changes in conformation alter the spacial relationship between the GFP and luciferase. The modulator can be a single entity covalently attached to one or both of the luciferase and GFP; it can be two separate entities each linked
- 30 to either the luciferase or GFP. The modulator(s), GFP and luciferase can be a single fusion protein, or a fusion protein of at least two of the entities. The components can be chemically linked, such as through thiol or disulfide linkages,

-31-

using linkers as provided herein. The GFP and luciferase can be linked directly or via linker, which can be a chemical linkage.

As used herein, "not strictly catalytically" means that the photoprotein acts as a catalyst to promote the oxidation of the substrate, but it is changed in the reaction, since the bound substrate is oxidized and bound molecular oxygen is used in the reaction. Such photoproteins are regenerated by addition of the substrate and molecular oxygen under appropriate conditions known to those of skill in this art.

As used herein, "nucleic acid" refers to a polynucleotide containing at least two covalently linked nucleotide or nucleotide analog subunits. A nucleic acid can be a deoxyribonucleic acid (DNA), a ribonucleic acid (RNA), or an analog of DNA or RNA. Nucleotide analogs are commercially available and methods of preparing polynucleotides containing such nucleotide analogs are known (Lin *et al.* (1994) *Nucl. Acids Res.* 22:5220-5234; Jellinek *et al.* (1995) *Biochemistry* 34:11363-11372; Pagratis *et al.* (1997) *Nature Biotechnol.* 15:68-73). The nucleic acid can be single-stranded, double-stranded, or a mixture thereof. For purposes herein, unless specified otherwise, the nucleic acid is double-stranded, or it is apparent from the context.

As used herein, a second messenger includes, but are not limited to, cAMP, cGMP, inositol phosphates, such as IP2 and IP3, NO (nitric oxide), Ca^{2+} , ceramide; DAG and arachidonic acid.

Hence, the term "nucleic acid" refers to single-stranded and/or double-stranded polynucleotides, such as deoxyribonucleic acid (DNA) and ribonucleic acid (RNA), as well as analogs or derivatives of either RNA or DNA. Also included in the term "nucleic acid" are analogs of nucleic acids such as peptide nucleic acid (PNA), phosphorothioate DNA, and other such analogs and derivatives.

As used herein, the term "nucleic acid molecule" and "nucleic acid fragment" are used interchangeably.

As used herein, DNA is meant to include all types and sizes of DNA molecules including cDNA, plasmids and DNA including modified nucleotides and nucleotide analogs.

-32-

As used herein, nucleotides include nucleoside mono-, di-, and triphosphates. Nucleotides also include modified nucleotides, such as, but are not limited to, phosphorothioate nucleotides and deazapurine nucleotides and other nucleotide analogs.

5 As used herein, a nucleic acid probe is single-stranded DNA or RNA that has a sequence of nucleotides that includes at least 14 contiguous bases, preferably at least 16 contiguous bases, typically about 30, that are the same as (or the complement of) any 14 or more contiguous bases set forth in any of SEQ ID Nos. 23-25 and herein. Among the preferred regions from which to construct
10 probes include 5' and/or 3' coding sequences, sequences predicted to encode regions that are conserved among *Renilla* species. Probes from regions conserved among *Renilla* species GFPs are for isolating GFP-encoding nucleic acid from *Renilla* libraries.

In preferred embodiments, the nucleic acid probes are degenerate probes
15 of at least 14 nucleotides, preferably 16 to 30 nucleotides.

In preferred embodiments, the nucleic acid probes are degenerate probes of at least 14 nucleotides, preferably 16 to 30 nucleotides, that are based on amino acids of *Renilla reniformis* set forth in above.

As used herein, vector (or plasmid) refers to discrete elements that are
20 used to introduce heterologous DNA into cells for either expression or replication thereof. Selection and use of such vehicles are well within the skill of the artisan. An expression vector includes vectors capable of expressing DNA operatively linked with regulatory sequences, such as promoter regions, that are capable of effecting expression of such DNA molecules. Thus, an expression
25 vector refers to a recombinant DNA or RNA construct, such as a plasmid, a phage, recombinant virus or other vector that, upon introduction into an appropriate host cell, results in expression of the cloned DNA. Appropriate expression vectors are well known to those of skill in the art and include those that are replicable in eukaryotic cells and/or prokaryotic cells and those that
30 remain episomal or those which integrate into the host cell genome. Presently preferred plasmids for expression of *Gaussia* luciferase, *Renilla* GFP and

-33-

luciferase are those that are expressed in bacteria and yeast, such as those described herein.

As used herein, a promoter region or promoter element refers to a segment of DNA or RNA that controls transcription of the DNA or RNA to which it is operatively linked. The promoter region includes specific sequences that are sufficient for RNA polymerase recognition, binding and transcription initiation. This portion of the promoter region is referred to as the promoter. In addition, the promoter region includes sequences that modulate this recognition, binding and transcription initiation activity of RNA polymerase. These sequences may be *cis* acting or may be responsive to *trans* acting factors. Promoters, depending upon the nature of the regulation, may be constitutive or regulated. Exemplary promoters contemplated for use in prokaryotes include the bacteriophage T7 and T3 promoters, and the like.

As used herein, operatively linked or operationally associated refers to the functional relationship of DNA with regulatory and effector sequences of nucleotides, such as promoters, enhancers, transcriptional and translational stop sites, and other signal sequences. For example, operative linkage of DNA to a promoter refers to the physical and functional relationship between the DNA and the promoter such that the transcription of such DNA is initiated from the promoter by an RNA polymerase that specifically recognizes, binds to and transcribes the DNA. In order to optimize expression and/or *in vitro* transcription, it may be necessary to remove, add or alter 5' untranslated portions of the clones to eliminate extra, potentially inappropriate alternative translation initiation (i.e., start) codons or other sequences that may interfere with or reduce expression, either at the level of transcription or translation. Alternatively, consensus ribosome binding sites (see, *e.g.*, Kozak (1991) *J. Biol. Chem.* 266:19867-19870) can be inserted immediately 5' of the start codon and may enhance expression. The desirability of (or need for) such modification may be empirically determined.

As used herein, to target a targeted agent, such as a luciferase, means to direct it to a cell that expresses a selected receptor or other cell surface protein by linking the agent to a such agent. Upon binding to or interaction with the

-34-

receptor or cell surface protein the targeted agent, can be reacted with an appropriate substrate and activating agents, whereby bioluminescent light is produced and the tumorous tissue or cells distinguished from non-tumorous tissue.

- 5 As used herein, an effective amount of a compound for treating a particular disease is an amount that is sufficient to ameliorate, or in some manner reduce the symptoms associated with the disease. Such amount may be administered as a single dosage or may be administered according to a regimen, whereby it is effective. The amount may cure the disease but,
- 10 typically, is administered in order to ameliorate the symptoms of the disease. Repeated administration may be required to achieve the desired amelioration of symptoms.

- As used herein, an effective amount of a conjugate for diagnosing a disease is an amount that will result in a detectable tissue. The tissues are
- 15 detected by visualization either without aid from a detector more sensitive than the human eye, or with the use of a light source to excite any fluorescent products.

- As used herein, visualizable means detectable by eye, particularly during surgery under normal surgical conditions, or, if necessary, slightly dimmed light.
- 20 As used herein, pharmaceutically acceptable salts, esters or other derivatives of the conjugates include any salts, esters or derivatives that may be readily prepared by those of skill in this art using known methods for such derivatization and that produce compounds that may be administered to animals or humans without substantial toxic effects and that either are pharmaceutically
- 25 active or are prodrugs.

 As used herein, treatment means any manner in which the symptoms of a conditions, disorder or disease are ameliorated or otherwise beneficially altered. Treatment also encompasses any pharmaceutical use of the compositions herein.

- As used herein, amelioration of the symptoms of a particular disorder by
- 30 administration of a particular pharmaceutical composition refers to any lessening, whether permanent or temporary, lasting or transient that can be attributed to or associated with administration of the composition.

-35-

As used herein, substantially pure means sufficiently homogeneous to appear free of readily detectable impurities as determined by standard methods of analysis, such as thin layer chromatography (TLC), gel electrophoresis and high performance liquid chromatography (HPLC), used by those of skill in the art to assess such purity, or sufficiently pure such that further purification would not detectably alter the physical and chemical properties, such as enzymatic and biological activities, of the substance. Methods for purification of the compounds to produce substantially chemically pure compounds are known to those of skill in the art. A substantially chemically pure compound may, however, be a mixture of stereoisomers or isomers. In such instances, further purification might increase the specific activity of the compound.

As used herein, a prodrug is a compound that, upon *in vivo* administration, is metabolized or otherwise converted to the biologically, pharmaceutically or therapeutically active form of the compound. To produce a prodrug, the pharmaceutically active compound is modified such that the active compound will be regenerated by metabolic processes. The prodrug may be designed to alter the metabolic stability or the transport characteristics of a drug, to mask side effects or toxicity, to improve the flavor of a drug or to alter other characteristics or properties of a drug. By virtue of knowledge of pharmacodynamic processes and drug metabolism *in vivo*, those of skill in this art, once a pharmaceutically active compound is known, can design prodrugs of the compound (see, *e.g.*, Nogrady (1985) *Medicinal Chemistry A Biochemical Approach*, Oxford University Press, New York, pages 388-392).

As used herein, biological activity refers to the *in vivo* activities of a compound or physiological responses that result upon *in vivo* administration of a compound, composition or other mixture. Biological activity, thus, encompasses therapeutic effects and pharmaceutical activity of such compounds, compositions and mixtures. Biological activities may be observed in *in vitro* systems designed to test or use such activities. Thus, for purposes herein the biological activity of a luciferase is its oxygenase activity whereby, upon oxidation of a substrate, light is produced.

-36-

As used herein, targeting agent (TA) refers to an agent that specifically or preferentially targets a linked targeted agent, a luciferin or luciferase, to a neoplastic cell or tissue.

As used herein, tumor antigen refers to a cell surface protein expressed or
5 located on the surface of tumor cells.

As used herein, neoplastic cells include any type of transformed or altered cell that exhibits characteristics typical of transformed cells, such as a lack of contact inhibition and the acquisition of tumor-specific antigens. Such cells include, but are not limited to leukemic cells and cells derived from a tumor.

10 As used herein, neoplastic disease is any disease in which neoplastic cells are present in the individual afflicted with the disease. Such diseases include, any disease characterized as cancer.

As used herein, metastatic tumors refers to tumors that are not localized in one site.

15 As used herein, specialty tissue refers to non-tumorous tissue for which information regarding location is desired. Such tissues include, for example, endometriotic tissue, ectopic pregnancies, tissues associated with certain disorders and myopathies or pathologies.

As used herein, a receptor refers to a molecule that has an affinity for a
20 given ligand. Receptors may be naturally-occurring or synthetic molecules. Receptors may also be referred to in the art as anti-ligands. As used herein, the receptor and anti-ligand are interchangeable. Receptors can be used in their unaltered state or as aggregates with other species. Receptors may be attached, covalently or noncovalently, or in physical contact with, to a binding member,
25 either directly or indirectly via a specific binding substance or linker. Examples of receptors, include, but are not limited to: antibodies, cell membrane receptors surface receptors and internalizing receptors, monoclonal antibodies and antisera reactive with specific antigenic determinants (such as on viruses, cells, or other materials), drugs, polynucleotides, nucleic acids, peptides, cofactors, lectins,
30 sugars, polysaccharides, cells, cellular membranes, and organelles.

-37-

Examples of receptors and applications using such receptors, include but are not restricted to:

- a) enzymes: specific transport proteins or enzymes essential to survival of microorganisms, which could serve as targets for antibiotic (ligand) selection;
- 5 b) antibodies: identification of a ligand-binding site on the antibody molecule that combines with the epitope of an antigen of interest may be investigated; determination of a sequence that mimics an antigenic epitope may lead to the development of vaccines of which the immunogen is based on one or more of such sequences or lead to the development of related diagnostic agents
- 10 c) nucleic acids: identification of ligand, such as protein or RNA, binding sites;
- d) catalytic polypeptides: polymers, preferably polypeptides, that are capable of promoting a chemical reaction involving the conversion of one or
- 15 more reactants to one or more products; such polypeptides generally include a binding site specific for at least one reactant or reaction intermediate and an active functionality proximate to the binding site, in which the functionality is capable of chemically modifying the bound reactant (see, *e.g.*, U.S. Patent No. 5,215,899);
- 20 e) hormone receptors: determination of the ligands that bind with high affinity to a receptor is useful in the development of hormone replacement therapies; for example, identification of ligands that bind to such receptors may lead to the development of drugs to control blood pressure; and
- f) opiate receptors: determination of ligands that bind to the opiate
- 25 receptors in the brain is useful in the development of less-addictive replacements for morphine and related drugs.

As used herein, antibody includes antibody fragments, such as Fab fragments, which are composed of a light chain and the variable region of a heavy chain.

- 30 As used herein, an antibody conjugate refers to a conjugate in which the targeting agent is an antibody.

-38-

As used herein, antibody activation refers to the process whereby activated antibodies are produced. Antibodies are activated upon reaction with a linker, such as heterobifunctional reagent.

As used herein, a surgical viewing refers to any procedure in which an opening is made in the body of an animal. Such procedures include traditional surgeries and diagnostic procedures, such as laparoscopies and arthroscopic procedures.

As used herein, humanized antibodies refer to antibodies that are modified to include "human" sequences of amino acids so that administration to a human will not provoke an immune response. Methods for preparation of such antibodies are known. For example, the hybridoma that expresses the monoclonal antibody is altered by recombinant DNA techniques to express an antibody in which the amino acid composition of the non-variable regions is based on human antibodies. Computer programs have been designed to identify such regions.

As used herein, ATP, AMP, NAD⁺ and NADH refer to adenosine triphosphate, adenosine monophosphate, nicotinamide adenine dinucleotide (oxidized form) and nicotinamide adenine dinucleotide (reduced form), respectively.

As used herein, production by recombinant means by using recombinant DNA methods means the use of the well known methods of molecular biology for expressing proteins encoded by cloned DNA.

As used herein, substantially identical to a product means sufficiently similar so that the property of interest is sufficiently unchanged so that the substantially identical product can be used in place of the product.

As used herein equivalent, when referring to two sequences of nucleic acids means that the two sequences in question encode the same sequence of amino acids or equivalent proteins. When "equivalent" is used in referring to two proteins or peptides, it means that the two proteins or peptides have substantially the same amino acid sequence with only conservative amino acid substitutions (see, *e.g.*, Table 1, above) that do not substantially alter the activity or function of the protein or peptide. When "equivalent" refers to a

-39-

property, the property does not need to be present to the same extent (*e.g.*, two peptides can exhibit different rates of the same type of enzymatic activity), but the activities are preferably substantially the same. "Complementary," when referring to two nucleotide sequences, means that the two sequences of
5 nucleotides are capable of hybridizing, preferably with less than 25%, more preferably with less than 15%, even more preferably with less than 5%, most preferably with no mismatches between opposed nucleotides. Preferably the two molecules will hybridize under conditions of high stringency.

As used herein: stringency of hybridization in determining percentage
10 mismatch is as follows:

- 1) high stringency: 0.1 x SSPE, 0.1% SDS, 65°C
- 2) medium stringency: 0.2 x SSPE, 0.1% SDS, 50°C
- 3) low stringency: 1.0 x SSPE, 0.1% SDS, 50°C

It is understood that equivalent stringencies may be achieved using alternative
15 buffers, salts and temperatures.

The term "substantially" identical or homologous or similar varies with the context as understood by those skilled in the relevant art and generally means at least 70%, preferably means at least 80%, more preferably at least 90%, and most preferably at least 95% identity. The terms "homology" and "identity" are
20 often used interchangeably. In general, sequences are aligned so that the highest order match is obtained (see, *e.g.*: *Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part I*, Griffin, A.M., and Griffin,
25 H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991; Carillo *et al.* (1988) *SIAM J Applied Math* 48:1073). By sequence identity, the number of conserved amino acids are determined by standard alignment
30 algorithms programs, and are used with default gap penalties established by each supplier. Substantially homologous nucleic acid molecules would hybridize typically at moderate stringency or at high stringency all along the length of the

-40-

nucleic acid of interest. Also contemplated are nucleic acid molecules that contain degenerate codons in place of codons in the hybridizing nucleic acid molecule.

Whether any two nucleic acid molecules have nucleotide sequences that are at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% "identical" can be determined using known computer algorithms such as the "FAST A" program, using for example, the default parameters as in Pearson *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:2444 (other programs include the GCG program package (Devereux, J., *et al.*, *Nucleic Acids Research* 12(II):387 (1984)), BLASTP, BLASTN, FASTA (Atschul, S.F., *et al.*, *J Molec Biol* 215:403 (1990); Guide to Huge Computers, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo *et al.* (1988) *SIAM J Applied Math* 48:1073). For example, the BLAST function of the National Center for Biotechnology Information database may be used to determine identity. Other commercially or publicly available programs include, DNASTar "MegAlign" program (Madison, WI) and the University of Wisconsin Genetics Computer Group (UWG) "Gap" program (Madison WI)). Percent homology or identity of proteins and/or nucleic acid molecules may be determined, for example, by comparing sequence information using a GAP computer program (*e.g.*, Needleman *et al.* (1970) *J. Mol. Biol.* 48:443, as revised by Smith and Waterman ((1981) *Adv. Appl. Math.* 2:482). Briefly, the GAP program defines similarity as the number of aligned symbols (*i.e.*, nucleotides or amino acids) which are similar, divided by the total number of symbols in the shorter of the two sequences. Default parameters for the GAP program may include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) and the weighted comparison matrix of Gribskov *et al.* (1986) *Nucl. Acids Res.* 14:6745, as described by Schwartz and Dayhoff, eds., *ATLAS OF PROTEIN SEQUENCE AND STRUCTURE*, National Biomedical Research Foundation, pp. 353-358 (1979); (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps.

Therefore, as used herein, the term "identity" represents a comparison between a test and a reference polypeptide or polynucleotide. For example, a

-41-

test polypeptide may be defined as any polypeptide that is 90% or more identical to a reference polypeptide. As used herein, the term at least "90% identical to" refers to percent identities from 90 to 99.99 relative to the reference polypeptides. Identity at a level of 90% or more is indicative of the fact that, assuming for exemplification purposes a test and reference polynucleotide length of 100 amino acids are compared. No more than 10% (i.e., 10 out of 100) amino acids in the test polypeptide differs from that of the reference polypeptides. Similar comparisons may be made between a test and reference polynucleotides. Such differences may be represented as point mutations randomly distributed over the entire length of an amino acid sequence or they may be clustered in one or more locations of varying length up to the maximum allowable, e.g. 10/100 amino acid difference (approximately 90% identity). Differences are defined as nucleic acid or amino acid substitutions, or deletions. At level of homologies or identities above about 85-90%, the result should be independent of the program and gap parameters set; such high levels of identity readily can be assess, often without relying on software.

As used herein, primer refers to an oligonucleotide containing two or more deoxyribonucleotides or ribonucleotides, preferably more than three, from which synthesis of a primer extension product can be initiated. Experimental conditions conducive to synthesis include the presence of nucleoside triphosphates and an agent for polymerization and extension, such as DNA polymerase, and a suitable buffer, temperature and pH.

As used herein, a composition refers to any mixture. It may be a solution, a suspension, liquid, powder, a paste, aqueous, non-aqueous or any combination thereof.

As used herein, a combination refers to any association between two or among more items.

As used herein, fluid refers to any composition that can flow. Fluids thus encompass compositions that are in the form of semi-solids, pastes, solutions, aqueous mixtures, gels, lotions, creams and other such compositions.

Examples of receptors and applications using such receptors, include but are not restricted to:

-42-

a) enzymes: specific transport proteins or enzymes essential to survival of microorganisms, which could serve as targets for antibiotic (ligand) selection;

b) antibodies: identification of a ligand-binding site on the antibody molecule that combines with the epitope of an antigen of interest may be

5 investigated; determination of a sequence that mimics an antigenic epitope may lead to the development of vaccines of which the immunogen is based on one or more of such sequences or lead to the development of related diagnostic agents or compounds useful in therapeutic treatments such as for auto-immune diseases

c) nucleic acids: identification of ligand, such as protein or RNA, binding
10 sites;

d) catalytic polypeptides: polymers, preferably polypeptides, that are capable of promoting a chemical reaction involving the conversion of one or more reactants to one or more products; such polypeptides generally include a binding site specific for at least one reactant or reaction intermediate and an
15 active functionality proximate to the binding site, in which the functionality is capable of chemically modifying the bound reactant (see, *e.g.*, U.S. Patent No. 5,215,899);

e) hormone receptors: determination of the ligands that bind with high affinity to a receptor is useful in the development of hormone replacement
20 therapies; for example, identification of ligands that bind to such receptors may lead to the development of drugs to control blood pressure; and

f) opiate receptors: determination of ligands that bind to the opiate receptors in the brain is useful in the development of less-addictive replacements for morphine and related drugs.

25 As used herein, complementary refers to the topological compatibility or matching together of interacting surfaces of a ligand molecule and its receptor. Thus, the receptor and its ligand can be described as complementary, and furthermore, the contact surface characteristics are complementary to each other.

30 As used herein, a ligand-receptor pair or complex formed when two macromolecules have combined through molecular recognition to form a complex.

-43-

As used herein, a substrate refers to any matrix that is used either directly or following suitable derivatization, as a solid support for chemical synthesis, assays and other such processes. Preferred substrates herein, are silicon substrates or siliconized substrates that are derivitized on the surface
5 intended for linkage of anti-ligands and ligands and other macromolecules, including the fluorescent proteins, phycobiliproteins and other emission shifters.

As used herein, a matrix refers to any solid or semisolid or insoluble support on which the molecule of interest, typically a biological molecule, macromolecule, organic molecule or biospecific ligand is linked or contacted.
10 Typically a matrix is a substrate material having a rigid or semi-rigid surface. In many embodiments, at least one surface of the substrate will be substantially flat, although in some embodiments it may be desirable to physically separate synthesis regions for different polymers with, for example, wells, raised regions, etched trenches, or other such topology. Matrix materials include any materials
15 that are used as affinity matrices or supports for chemical and biological molecule syntheses and analyses, such as, but are not limited to: polystyrene, polycarbonate, polypropylene, nylon, glass, dextran, chitin, sand, pumice, polytetrafluoroethylene, agarose, polysaccharides, dendrimers, buckyballs, polyacrylamide, Kieselguhr-polyacrylamide non-covalent composite, polystyrene-
20 polyacrylamide covalent composite, polystyrene-PEG (polyethyleneglycol) composite, silicon, rubber, and other materials used as supports for solid phase syntheses, affinity separations and purifications, hybridization reactions, immunoassays and other such applications.

As used herein, the attachment layer refers the surface of the chip device
25 to which molecules are linked. Typically, the chip is a semiconductor device, which is coated on at least a portion of the surface to render it suitable for linking molecules and inert to any reactions to which the device is exposed. Molecules are linked either directly or indirectly to the surface, linkage may be effected by absorption or adsorption, through covalent bonds, ionic interactions or any other
30 interaction. Where necessary the attachment layer is adapted, such as by derivatization for linking the molecules.

-44-

B. FLUORESCENT PROTEINS

The GFP from *Aequorea* and that of the sea pansy *Renilla reniformis* share the same chromophore, yet *Aequorea* GFP has two absorbance peaks at 395 and 475 nm, whereas *Renilla* GFP has only a single absorbance peak at 498 nm, with about 5.5 fold greater monomer extinction coefficient than the major 395 nm peak of the *Aequorea* protein (Ward, W. W. in Bioluminescence and Chemiluminescence (eds. DeLuca, M. A. & McElroy, W. D.) 235-242 (Academic Press, New York, 1981)). The spectra of the isolated chromophore and denatured protein at neutral pH do not match the spectra of either native protein (Cody, C. W. *et al.* (1993) *Biochemistry* 32:1212-1218).

1. Green and blue fluorescent proteins

As described herein, blue light is produced using the *Renilla* luciferase or the *Aequorea* photoprotein in the presence of Ca^{2+} and the coelenterazine luciferin or analog thereof. This light can be converted into a green light if a green fluorescent protein (GFP) is added to the reaction. Green fluorescent proteins, which have been purified (see, *e.g.*, Prasher *et al.* (1992) *Gene* 111:229-233) and also cloned (see, *e.g.*, International PCT Application No. WO 95/07463, which is based on U.S. application Serial No. 08/119,678 and U.S. application Serial No. 08/192,274, which are herein incorporated by reference), are used by cnidarians as energy-transfer acceptors. GFPs fluoresce *in vivo* upon receiving energy from a luciferase-oxyluciferin excited-state complex or a Ca^{2+} -activated photoprotein. The chromophore is modified amino acid residues within the polypeptide. The best characterized GFPs are those of *Aequorea* and *Renilla* (see, *e.g.*, Prasher *et al.* (1992) *Gene* 111:229-233; Hart, *et al.* (1979) *Biochemistry* 18:2204-2210). For example, a green fluorescent protein (GFP) from *Aequorea victoria* contains 238 amino acids, absorbs blue light and emits green light. Thus, inclusion of this protein in a composition containing the aequorin photoprotein charged with coelenterazine and oxygen, can, in the presence of calcium, result in the production of green light. Thus, it is

-45-

contemplated that GFPs may be included in the bioluminescence generating reactions that employ the aequorin or *Renilla* luciferases or other suitable luciferase in order to enhance or alter color of the resulting bioluminescence.

2. *Renilla reniformis* GFP

- 5 Purified *Renilla reniformis* GFP and muteins thereof are provided. Presently preferred *Renilla* GFP for use in the compositions herein is *Renilla reniformis* GFP having the sequence of amino acids set forth in SEQ ID No. 27. The *Renilla* GFP and GFP peptides can be isolated from natural sources or isolated from a prokaryotic or eukaryotic cell transfected with nucleic acid that
- 10 encodes the *Renilla* GFP and/or GFP peptides, such as those encoded by the sequences of nucleotides set forth in SEQ ID Nos. 23-25.

The encoding nucleic acid molecules are provided. Preferred are those that encode the protein having the sequence of amino acids (SEQ ID No. 27):

- 15 mdlaklgkevmptkinleglvgdhafsmevgvgegnilegtqevkisvtkgaplpfafdivsvafsygn
raytgypeeisdyflqsfpegtyniryqdggtaivkdsiledgkfivnvdfkakdlrrmgpvmqqd
ivgmqpsyesmytnvtsvigeciiafklqtgkfhfymrtvykskpvvetmplyhfiqhrlvktnvdta
sgyvvqhetaiaahstikkiegsip,

and is preferably the sequence set forth in SEQ ID No. 26.

- In particular, nucleic acid molecules encoding a *Renilla reniformis* GFP
- 20 having any of the following sequences are provided (see SEQ ID Nos. 23-25):

Renilla reniformis GFP Clone-1

- GGCACGAGGGTTTCCTGACACAATAAAAACCTTTCAAATTGTTTCTC
TGTAGCAGTAAGTATGGATCTCGCAAACCTTGGTTTGAAGGAAGTG
ATGCCTACTAAAATCAACTTAGAAGGACTGGTTGGCGACCACGCTT
- 25 TCTCAATGGAAGGAGTTGGCGAAGGCAACATATTGGAAGGAACTCA
AGAGGTGAAGATATCGGTAACAAAAGGCGCACCCTCCATTTCGC
ATTTGATATCGTATCTGTGGCTTTTTCATATGGGAACAGAGCTTA
TACCGGTTACCCAGAAGAAATTTCCGACTACTTCCTCCAGTCGTT
TCCAGAAGGCTTTACTTACGAGAGAAACATTTCGTTATCAAGATGG
- 30 AGGAACTGCAATTGTTAAATCTGATATAAGCTTGGAAGATGGTAA
ATTCATAGTGAATGTAGACTTCAAAGCGAAGGATCTACGTCGCAT
GGGACCAGTCATGCAGCAAGACATCGTGGGTATGCAGCCATCGTA
TGAGTCAATGTACACCAATGTCACTTCAGTTATAGGGGAATGTAT
AATAGCATTCAAACCTTCAAACCTGGCAAGCATTTCACCTTACCACAT
- 35 GAGGACAGTTTACAAATCAAAGAAGCCAGTGGAACCTATGCCATTG
TATCATTTTCATCCAGCATCGCCTCGTTAAGACCAATGTGGACACAG
CCAGTGGTTACGTTGTGCAACACGAGACAGCAATTGCAGCGCATT
TACAATCAAAAAAATTGAAGGCTCTTTACCATAGATACCTGTACACA

-46-

ATTATTCTATGCACGTAGCATTTTTTTTGGAAATATAAGTGGTATTGT
TCAATAAAATATTAAATATAAAAAAAAAAAAAAAAAAAAAAAAAA;

Renilla reniformis GFP Clone-2

5
GGCACGAGGCTGACACAATAAAAAACCTTTCAAATTGTTTCTCTGTAGCAGG
AAGTATGGATCTCGCAAACTTGGTTTGAAGGAAGTGATGCCTACTAAAATC
AACTTAGAAGGACTGGTTGGCGACCACGCTTTCTCAATGGAAGGAGTTGGCG
AAGGCAACATATTGGAAGGAACTCAAGAGGTGAAGATATCGGTAACAAAAGG
10 CGCACCACTCCCATTCGCATTTGATATCGTATCTGTTGCTTTCTCATATGGG
AACAGAGCTTATACTGGTTACCCAGAAGAAATTTCCGACTACTTCCTCCAGT
CGTTTCCAGAAGGCTTTACTTACGAGAGAAACATTTCGTTATCAAGATGGAGG
AACTGCAATTGTTAAATCTGATATAAGCTTGAAGATGGTAAATTCATAGTG
AATGTAGACTTCAAAGCGAAGGATCTACGTCGCATGGGACCAGTCATGCAGC
15 AAGACATCGTGGGTATGCAGCCATCGTATGAGTCAATGTACACCAATGTCAC
TTCAGTTATAGGGGAATGTATAATAGCATTCAAACCTCAAACCTGGCAAACAT
TTCATTACCACATGAGGACAGTTTACAAATCAAAGAAGCCAGTGGAAACTA
TGCCATTGTATCATTTTCATCCAGCATCGCTCGTTAAGACCAATGTGGACAC
AGCCAGTGGTTACGTTGTGCAACACGAGACAGCAATTGCAGCGCATTCTACA
20 ATCAAAAAAATTGAAGGCTCTTTACCATAGATATCTATACACAATTATTCTA
TGCACGTAGCATTTTTTTTGGAAATATAAGTGGTATTGTTCAATAAAATATTA
AATATAAAAAAAAAAAAAAAAAAAAAAAAAA; and

Renilla reniformis GFP Clone-3

25
GGCACGAGGGTTTCCTGACACAATAAAAAACCTTTCAAATTGTTTCTCTGTAG
CAGTAAGTATGGATCTCGCAAACTTGGTTTGAAGGAAGTGATGCCTACTA
AAATCAACTTAGAAGGACTGGTTGGCGACCACGCTTTCTCAATGGAAGGAGTT
GGCGAAGGCAACATATTGGAAGGAACTCAAGAGGTGAAGATATCGGTAACAA
30 AAGGCGCACCCTCCCATTCGCATTTGATATCGTATCTGTGGCTTTTTTCATAT
GGGAACAGAGCTTATACCGGTTACCCAGAAGAAATTTCCGACTACTTCCTCCA
GTCGTTTCCAGAAGGCTTTACTTACGAGAGAAACATTTCGTTATCAAGATGGAG
GAACTGCAATTGTTAAATCTGATATAAGCTTGAAGATGGTAAATTCATAGT
GAATGTAGACTTCAAAGCGAAGGATCTACGTCGCATGGGACCAGTCATGCAG
35 CAAGACATCGTGGGTATGCAGCCATCGTATGAGTCAATGTACACCAATGTCAC
TTCAGTTATAGGGGAATGTATAATAGCATTCAAACCTCAAACCTGGCAAGCATT
TCACTTACCACATGAGGACAGTTTACAAATCAAAGAAGCCAGTGGAAACTATG
CCATTGTATCATTTTCATCCAGCATCGCTCGTTAAGACCAATGTGGACACAGC
CAGTGGTTACGTTGTGCAACACGAGACAGCAATTGCAGCGCATTCTACAATCA
40 AAAAAATTGAAGGCTCTTTACCATAGATACCTGTACACAATTATTCTATGCAC
GTAGCATTTTTTTTGGAAATATAAGTGGTATTGTTCAATAAAATATTAAATATAT
GCTTTTGCAAAAAAAAAAAAAAAAAAAAAAAAAA.

-47-

An exemplary mutein is set forth in SEQ ID No. 33, and humanized codon are set forth in SEQ ID No. 26.

Also contemplated are the coding portion of the sequence of nucleotides
5 that hybridize under moderate or high stringency to the sequence of nucleotides set forth above, particularly when using probes provided herein. Probes derived from this nucleic acid that can be used in methods provided herein to isolate GFPs from any *Renilla reniformis* species are provided. In an exemplary embodiment, nucleic acid encoding *Renilla reniformis* GFP is provided. This
10 nucleic acid encodes the sequence of amino acids set forth above.

GFPs, including the *Renilla reniformis* protein provided herein, are activated by blue light to emit green light and thus may be used in the absence of luciferase and in conjunction with an external light source with novelty items (see U.S. Patent Nos. 5,876,995, 6,152,358 and 6,113,886) and in conjunction
15 with bioluminescence generating system for novelty items (see U.S. Patent Nos. 5,876,995, 6,152,358 and 6,113,886), for tumor diagnosis (see, allowed co-pending U.S. application Serial No. 08/908,909) and in biochips (see, U.S. application Serial No. 08/990,103, which is published as International PCT application No. WO 98/26277).

20 *Renilla reniformis* GFP is intended for use in any of the novelty items and combinations, such as the foods, including beverages, greeting cards, and toys, including bubble making toys, particularly bubble-making compositions or mixtures. Also of particular interest are the use of these proteins in cosmetics, particularly face paints or make-up, hair colorants or hair conditioners, mousses
25 or other such products and skin creams. Such systems are particularly of interest because no luciferase is needed to activate the photoprotein and because the proteins are non-toxic and safe to apply to the skin, hair, eyes and to ingest. These fluorescent proteins may also be used in addition to bioluminescence generating systems to enhance or create an array of different
30 colors. Transgenic animals and plants that express the *Renilla reniformis* GFP-encoding nucleic acid are also provided. Such animals and plants, include transgenic fish, transgenic worms for use, for example, as lures for fishing;

-48-

transgenic animals, such as monkeys and rodents for research in which a marker gene is used, and transgenic animals as novelty items and to produce glowing foods, such as ham, eggs, chicken, and other meats; transgenic plants in which the *Renilla reniformis* is a marker, and also transgenic plants that are novelty items, particularly ornamental plants, such as glowing orchids, roses and other flowering plants.

The *Renilla reniformis* GFP may be used alone or in combination with bioluminescence generating systems to produce an array of colors. They may be used in combinations such that the color of, for example, a beverage changes over time, or includes layers of different colors. The cloning and expression of *Renilla reniformis* GFP and uses thereof are described below.

C. BIOLUMINESCENCE GENERATING SYSTEMS AND COMPONENTS

The following is a description of bioluminescence generating systems and the components thereof. The *Renilla reniformis* GFP provided herein can be used alone for a variety of applications, and with any compatible bioluminescence generating systems.

A bioluminescence-generating system refers to the components that are necessary and sufficient to generate bioluminescence. These include a luciferase, luciferin and any necessary co-factors or conditions. Virtually any bioluminescent system known to those of skill in the art will be amenable to use in the apparatus, systems, combinations and methods provided herein. Factors for consideration in selecting a bioluminescent-generating system, include, but are not limited to: the targeting agent used in combination with the bioluminescence; the medium in which the reaction is run; stability of the components, such as temperature or pH sensitivity; shelf life of the components; sustainability of the light emission, whether constant or intermittent; availability of components; desired light intensity; color of the light; and other such factors. Such bioluminescence generating systems are known (see those described in U.S. Patent Nos. 5,876,995, 6,152,358 and 6,113,886).

1. General description

In general, bioluminescence refers to an energy-yielding chemical reaction in which a specific chemical substrate, a luciferin, undergoes oxidation,

-49-

catalyzed by an enzyme, a luciferase. Bioluminescent reactions are easily maintained, requiring only replenishment of exhausted luciferin or other substrate or cofactor or other protein, in order to continue or revive the reaction.

Bioluminescence generating reactions are well-known to those of skill in this art

- 5 and any such reaction may be adapted for use in combination with articles of manufacture as described herein.

There are numerous organisms and sources of bioluminescence generating systems, and some representative genera and species that exhibit bioluminescence are set forth in the following table (reproduced in part from

- 10 Hastings in (1995) *Cell Physiology: Source Book*, N. Sperelakis (ed.), Academic Press, pp 665-681):

TABLE 2
Representative luminous organism

15	Type of Organism	Representative genera
	Bacteria	Photobacterium Vibrio Xenorhabdus
	Mushrooms	Panus, Armillaria Pleurotus
	Dinoflagellates	Gonyaulax Pyrocystis Noctiluca
20	Cnidaria (coelenterates) Jellyfish Hydroid Sea Pansy	Aequorea Obelia Renilla
	Ctenophores	Mnemiopsis Beroe
25	Annelids Earthworms Marine polychaetes Syllid fireworm	Diplocardia Chaetopterus, Phyxotrix Odontosyllis

-50-

	Type of Organism	Representative genera
	Molluscs Limpet Clam Squid	Latia Pholas Heteroteuthis Heterocarpus
5	Crustacea Ostracod	Vargula (Cypridina)
10	Shrimp (euphausids) Decapod Copepods	Meganyctiphanes Acanthophyra Oplophorus Gnathophausia Sergestes
15	Insects Coleopterids (beetles) Firefly Click beetles Railroad worm Diptera (flies)	Photinus, Photuris Pyrophorus Phengodes, Phrixothrix Arachnocampa
20	Echinoderms Brittle stars Sea cucumbers	Ophiopsila Laetmogone
	Anthozoans	Renilla
	Chordates Tunicates	Pyrosoma
25	Fish Cartilaginous Bony	Squalus
30	Ponyfish Flashlight fish Angler fish Midshipman Lantern fish Shiny loosejaw Hatchet fish and other fish	Leiognathus Photoblepharon Cryptopsaras Porichthys Benia Aristostomias Agyropelecus Pachystomias Malacosteus
35	Midwater fish	Cyclothone Neoscopelus Tarletonbeania

-51-

Other bioluminescent organisms contemplated for use herein are *Gonadostomias*, *Gaussia* (copepods), *Watensia*, *Halisturia*, Vampire squid, *Glyphus*, Myctophids (fish), *Vinciguerria*, *Howella*, *Florenciella*, *Chaudiodus*, *Melanocostus* and Sea Pens.

5 It is understood that a bioluminescence generating system may be isolated from natural sources, such as those in the above Table, or may be produced synthetically. In addition, for uses herein, the components need only be sufficiently pure so that mixture thereof, under appropriate reaction conditions, produces a glow so that cells and tissues can be visualized during a
10 surgical procedure.

 Thus, in some embodiments, a crude extract or merely grinding up the organism may be adequate. Generally, however, substantially pure components are used. Also, components may be synthetic components that are not isolated from natural sources. DNA encoding luciferases is available (see, *e.g.*, SEQ ID
15 Nos. 1-13) and has been modified (see, *e.g.*, SEQ ID Nos. 3 and 10-13) and synthetic and alternative substrates have been devised. The DNA listed herein is only representative of the DNA encoding luciferases that is available.

 Any bioluminescence generating system, whether synthetic or isolated from natural sources, such as those set forth in Table 2, elsewhere herein or
20 known to those of skill in the art, is intended for use in the combinations, systems and methods provided herein. Chemiluminescence systems per se, which do not rely on oxygenases (luciferases) are not encompassed herein.

(a) Luciferases

 Luciferases refer to any compound that, in the presence of any necessary
25 activators, catalyze the oxidation of a bioluminescence substrate (luciferin) in the presence of molecular oxygen, whether free or bound, from a lower energy state to a higher energy state such that the substrate, upon return to the lower energy state, emits light. For purposes herein, luciferase is broadly used to encompass enzymes that act catalytically to generate light by oxidation of a substrate and
30 also photoproteins, such as aequorin, that act, though not strictly catalytically (since such proteins are exhausted in the reaction), in conjunction with a substrate in the presence of oxygen to generate light. These luciferases,

-52-

including photoproteins, such as aequorin, are herein also included among the luciferases. These reagents include the naturally-occurring luciferases (including photoproteins), proteins produced by recombinant DNA, and mutated or modified variants thereof that retain the ability to generate light in the presence of an
5 appropriate substrate, co-factors and activators or any other such protein that acts as a catalyst to oxidize a substrate, whereby light is produced.

Generically, the protein that catalyzes or initiates the bioluminescent reaction is referred to as a luciferase, and the oxidizable substrate is referred to as a luciferin. The oxidized reaction product is termed oxyluciferin, and certain
10 luciferin precursors are termed etioluciferin. Thus, for purposes herein bioluminescence encompasses light produced by reactions that are catalyzed by (in the case of luciferases that act enzymatically) or initiated by (in the case of the photoproteins, such as aequorin, that are not regenerated in the reaction) a biological protein or analog, derivative or mutant thereof.

15 For clarity herein, these catalytic proteins are referred to as luciferases and include enzymes such as the luciferases that catalyze the oxidation of luciferin, emitting light and releasing oxyluciferin. Also included among luciferases are photoproteins, which catalyze the oxidation of luciferin to emit light but are changed in the reaction and must be reconstituted to be used again.
20 The luciferases may be naturally occurring or may be modified, such as by genetic engineering to improve or alter certain properties. As long as the resulting molecule retains the ability to catalyze the bioluminescent reaction, it is encompassed herein.

Any protein that has luciferase activity (a protein that catalyzes oxidation
25 of a substrate in the presence of molecular oxygen to produce light as defined herein) may be used herein. The preferred luciferases are those that are described herein or that have minor sequence variations. Such minor sequence variations include, but are not limited to, minor allelic or species variations and insertions or deletions of residues, particularly cysteine residues. Suitable
30 conservative substitutions of amino acids are known to those of skill in this art and may be made generally without altering the biological activity of the

-53-

resulting molecule. Such substitutions are preferably made in accordance with those set forth in TABLE 1 as described above.

- The luciferases may be obtained commercially, isolated from natural sources, expressed in host cells using DNA encoding the luciferase, or obtained
- 5 in any manner known to those of skill in the art. For purposes herein, crude extracts obtained by grinding up selected source organisms may suffice. Since large quantities of the luciferase may be desired, isolation of the luciferase from host cells is preferred. DNA for such purposes is widely available as are modified forms thereof.
- 10 Examples of luciferases include, but are not limited to, those isolated from the ctenophores *Mnemiopsis* (mnemiopsin) and *Beroe ovata* (berovin), those isolated from the coelenterates *Aequorea* (aequorin), *Obelia* (obelin), *Pelagia*, the *Renilla* luciferase, the luciferases isolated from the mollusca *Pholas* (pholasin), the luciferases isolated from fish, such as *Aristostomias*, *Pachystomias* and
- 15 *Porichthys* and from the ostracods, such as *Cypridina* (also referred to as *Vargula*). Preferred luciferases for use herein are the Aequorin protein, *Renilla* luciferase and *Cypridina* (also called *Vargula*) luciferase (see, *e.g.*, SEQ ID Nos. 1, 2, and 4-13). Also, preferred are luciferases which react to produce red and/or near infrared light. These include luciferases found in species of
- 20 *Aristostomias*, such as *A. scintillans*, *Pachystomias*, *Malacosteus*, such as *M. niger*.

(b) Luciferins

- The substrates for the reaction or for inclusion in the conjugates include any molecule(s) with which the luciferase reacts to produce light. Such
- 25 molecules include the naturally-occurring substrates, modified forms thereof, and synthetic substrates (see, *e.g.*, U.S. Patent Nos. 5,374,534 and 5,098,828). Exemplary luciferins include those described herein, as well as derivatives thereof, analogs thereof, synthetic substrates, such as dioxetanes (see, *e.g.*, U.S. Patent Nos. 5,004,565 and 5,455,357), and other compounds that are
- 30 oxidized by a luciferase in a light-producing reaction (see, *e.g.*, U.S. Patent Nos. 5,374,534, 5,098,828 and 4,950,588). Such substrates also may be identified empirically by selecting compounds that are oxidized in bioluminescent reactions.

-54-

(c) **Activators**

The bioluminescent generating systems also require additional components discussed herein and known to those of skill in the art. All bioluminescent reactions require molecular oxygen in the form of dissolved or
5 bound oxygen. Thus, molecular oxygen, dissolved in water or in air or bound to a photoprotein, is the activator for bioluminescence reactions. Depending upon the form of the components, other activators include, but are not limited to, ATP (for firefly luciferase), flavin reductase (bacterial systems) for regenerating FMNH₂ from FMN, and Ca²⁺ or other suitable metal ion (aequorin).

10 Most of the systems provided herein will generate light when the luciferase and luciferin are mixed and exposed to air or water. The systems that use photoproteins that have bound oxygen, such as aequorin, however, will require exposure to Ca²⁺ (or other suitable metal ion), which can be provided in the form of an aqueous composition of a calcium salt. In these instances,
15 addition of a Ca²⁺ (or other suitable metal ion) to a mixture of luciferase (aequorin) and luciferin (such as coelenterazine) will result in generation of light. The *Renilla* system and other Anthozoa systems also require Ca²⁺ (or other suitable metal ion).

If crude preparations are used, such as ground up *Cypridina* (shrimp) or
20 ground fireflies, it may be necessary to add only water. In instances in which fireflies (or a firefly or beetle luciferase) are used the reaction may only require addition ATP. The precise components will be apparent, in light of the disclosure herein, to those of skill in this art or may be readily determined empirically.

25 It is also understood that these mixtures will also contain any additional salts or buffers or ions that are necessary for each reaction to proceed. Since these reactions are well-characterized, those of skill in the art will be able to determine precise proportions and requisite components. Selection of components will depend upon the apparatus, article of manufacture and
30 luciferase. Various embodiments are described and exemplified herein; in view of such description, other embodiments will be apparent.

-55-

(d) Reactions

In all embodiments, all but one component, either the luciferase or luciferin, of a bioluminescence generating system will be mixed or packaged with or otherwise combined. Since the result to be achieved is the production of light visible to the naked eye for qualitative, not quantitative, diagnostic purposes, the precise proportions and amounts of components of the bioluminescence reaction need not be stringently determined or met. They must be sufficient to produce light. Generally, an amount of luciferin and luciferase sufficient to generate a visible glow is used; this amount can be readily determined empirically and is dependent upon the selected system and selected application. Where quantitative measurements are required, more precision may be required.

For purposes herein, such amount is preferably at least the concentrations and proportions used for analytical purposes by those of skill in the such arts. Higher concentrations may be used if the glow is not sufficiently bright. Alternatively, a microcarrier coupled to more than one luciferase molecule linked to a targeting agent may be utilized to increase signal output. Also because the conditions in which the reactions are used are not laboratory conditions and the components are subject to storage, higher concentration may be used to overcome any loss of activity. Typically, the amounts are 1 mg, preferably 10 mg and more preferably 100 mg, of a luciferase per liter of reaction mixture or 1 mg, preferably 10 mg, more preferably 100 mg. Compositions may contain at least about 0.01 mg/l, and typically 0.1 mg/l, 1 mg/l, 10 mg/l or more of each component on the item. The amount of luciferin is also between about 0.01 and 100 mg/l, preferably between 0.1 and 10 mg/l, additional luciferin can be added to many of the reactions to continue the reaction. In embodiments in which the luciferase acts catalytically and does not need to be regenerated, lower amounts of luciferase can be used. In those in which it is changed during the reaction, it also can be replenished; typically higher concentrations will be selected. Ranges of concentration per liter (or the amount of coating on substrate the results from contacting with such composition) of each component on the order of 0.1 to 20 mg, preferably 0.1 to 10 mg, more preferably between about 1 and 10 mg of each component will be sufficient. When preparing coated substrates, as

-56-

described herein, greater amounts of coating compositions containing higher concentrations of the luciferase or luciferin may be used.

Thus, for example, in presence of calcium, 5 mg of luciferin, such as coelenterazine, in one liter of water will glow brightly for at least about 10 to 20
5 minutes, depending on the temperature of the water, when about 10 mg of luciferase, such as aequorin photoprotein luciferase or luciferase from *Renilla*, is added thereto. Increasing the concentration of luciferase, for example, to 100 mg/l, provides a particularly brilliant display of light.

It is understood, that concentrations and amounts to be used depend
10 upon the selected bioluminescence generating system but these may be readily determined empirically. Proportions, particularly those used when commencing an empirical determination, are generally those used for analytical purposes, and amounts or concentrations are at least those used for analytical purposes, but the amounts can be increased, particularly if a sustained and brighter glow is
15 desired.

For purposes herein, *Renilla reniformis* GFP is added to the reaction in order to shift the spectrum of the generated light.

2. The *Renilla* system

Renilla, also known as soft coral sea pansies, are members of the class of
20 coelenterates Anthozoa, which includes other bioluminescent genera, such as *Cavarnularia*, *Ptilosarcus*, *Stylatula*, *Acanthoptilum*, and *Parazoanthus*. Bioluminescent members of the Anthozoa genera contain luciferases and luciferins that are similar in structure (see, e.g., Cormier *et al.* (1973) *J. Cell. Physiol.* 81:291-298; see, also Ward *et al.* (1975) *Proc. Natl. Acad. Sci. U.S.A.*
25 72:2530-2534). The luciferases and luciferins from each of these anthozoans crossreact with one another and produce a characteristic blue luminescence.

Renilla luciferase and the other coelenterate and ctenophore luciferases, such as the aequorin photoprotein, use imidazopyrazine substrates, particularly the substrates generically called coelenterazine (see, formulae (I) and (II) of
30 Section C.4.b, below). Other genera that have luciferases that use a coelenterazine include: squid, such as *Chiroteuthis*, *Eucleoteuthis*,

-57-

Onychoteuthis, *Watasenia*, cuttlefish, *Sepiolina*; shrimp, such as *Oplophorus*, *Acanthophyra*, *Sergestes*, and *Gnathophausia*; deep-sea fish, such as *Argyropelecus*, *Yarella*, *Diaphus*, *Gonadostomias* and *Neoscopelus*.

Renilla luciferase does not, however, have bound oxygen, and thus requires dissolved oxygen in order to produce light in the presence of a suitable luciferin substrate. Since *Renilla* luciferase acts as a true enzyme (*i.e.*, it does not have to be reconstituted for further use) the resulting luminescence can be long-lasting in the presence of saturating levels of luciferin. Also, *Renilla* luciferase is relatively stable to heat.

- 10 *Renilla* luciferases, DNA encoding *Renilla reniformis* luciferase, and use of the *Renilla reniformis* DNA to produce recombinant luciferase, as well as DNA encoding luciferase from other coelenterates, are well known and available (see, *e.g.*, SEQ ID No. 1, U.S. Patent Nos. 5,418,155 and 5,292,658; see, also, Prasher *et al.* (1985) *Biochem. Biophys. Res. Commun.* 126:1259-1268;
- 15 Cormier (1981) "*Renilla* and *Aequorea* bioluminescence" in *Bioluminescence and Chemiluminescence*, pp. 225-233; Charbonneau *et al.* (1979) *J. Biol. Chem.* 254:769-780; Ward *et al.* (1979) *J. Biol. Chem.* 254:781-788; Lorenz *et al.* (1981) *Proc. Natl. Acad. Sci. U.S.A.* 88: 4438-4442; Hori *et al.* (1977) *Proc. Natl. Acad. Sci. U.S.A.* 74:4285-4287; Hori *et al.* (1975) *Biochemistry*
- 20 14:2371-2376; Hori *et al.* (1977) *Proc. Natl. Acad. Sci. U.S.A.* 74:4285-4287; Inouye *et al.* (1975) *Jap. Soc. Chem. Lett.* 141-144; and Matthews *et al.* (1979) *Biochemistry* 16:85-91). The DNA encoding *Renilla reniformis* luciferase and host cells containing such DNA provide a convenient means for producing large quantities of *Renilla reniformis* enzyme, such as in those known to those of skill
- 25 in the art (see, *e.g.*, U.S. Patent Nos. 5,418,155 and 5,292,658, which describe recombinant production of *Renilla reniformis* luciferase).

- When used herein, the *Renilla* luciferase can be packaged in lyophilized form, encapsulated in a vehicle, either by itself or in combination with the luciferin substrate. Prior to use the mixture is contacted with an aqueous
- 30 composition, preferably a phosphate buffered saline pH 7-8; dissolved O₂ will activate the reaction. Final concentrations of luciferase in the glowing mixture will be on the order of 0.01 to 1 mg/l or more. Concentrations of luciferin will

-58-

be at least about 10^{-8} M, but 1 to 100 or more orders of magnitude higher to produce a long lasting bioluminescence.

In certain embodiments herein, about 1 to 10 mg, or preferably 2-5 mg, more preferably about 3 mg of coelenterazine will be used with about 100 mg of *Renilla* luciferase. The precise amounts, of course can be determined empirically, and, also will depend to some extent on the ultimate concentration and application. In particular, addition of about 0.25 ml of a crude extract from the bacteria that express *Renilla* to 100 ml of a suitable assay buffer and about 0.005 μ g was sufficient to produce a visible and lasting glow (see, U.S. Patent Nos. 5,418,155 and 5,292,658, which describe recombinant production of *Renilla reniformis* luciferase).

Lyophilized mixtures, and compositions containing the *Renilla* luciferase are also provided. The luciferase or mixtures of the luciferase and luciferin may also be encapsulated into a suitable delivery vehicle, such as a liposome, glass particle, capillary tube, drug delivery vehicle, gelatin, time release coating or other such vehicle. The luciferase may also be linked to a substrate, such as biocompatible materials.

3. Ctenophore systems

Ctenophores, such as *Mnemiopsis* (mnemiopsin) and *Beroe ovata* (berovin), and coelenterates, such as *Aequorea* (aequorin), *Obelia* (obelin) and *Pelagia*, produce bioluminescent light using similar chemistries (see, e.g., Stephenson *et al.* (1981) *Biochimica et Biophysica Acta* 678:65-75; Hart *et al.* (1979) *Biochemistry* 18:2204-2210; International PCT Application No. WO 94/18342, which is based on U.S. application Serial No. 08/017,116, U.S. Patent No. 5,486,455 and other references and patents cited herein). The *Aequorin* and *Renilla* systems are representative and are described in detail herein as exemplary and as among the presently preferred systems. The *Aequorin* and *Renilla* systems can use the same luciferin and produce light using the same chemistry, but each luciferase is different. The *Aequorin* luciferase aequorin, as well as, for example, the luciferases mnemiopsin and berovin, is a photoprotein that includes bound oxygen and bound luciferin, requires Ca^{2+} (or other suitable metal ion) to trigger the reaction, and must be regenerated for

-59-

repeated use; whereas, the *Renilla* luciferase acts as a true enzyme because it is unchanged during the reaction and it requires dissolved molecular oxygen.

4. The aequorin system

The aequorin system is well known (see, *e.g.*, Tsuji *et al.* (1986)

- 5 "Site-specific mutagenesis of the calcium-binding photoprotein aequorin," *Proc. Natl. Acad. Sci. USA* 83:8107-8111; Prasher *et al.* (1985) "Cloning and Expression of the cDNA Coding for Aequorin, a Bioluminescent Calcium-Binding Protein," *Biochemical and Biophysical Research Communications* 126:1259-1268; Prasher *et al.* (1986) *Methods in Enzymology* 133:288-297;
- 10 Prasher, *et al.* (1987) "Sequence Comparisons of cDNAs Encoding for Aequorin Isotypes," *Biochemistry* 26:1326-1332; Charbonneau *et al.* (1985) "Amino Acid Sequence of the Calcium-Dependent Photoprotein Aequorin," *Biochemistry* 24:6762-6771; Shimomura *et al.* (1981) "Resistivity to denaturation of the apoprotein of aequorin and reconstitution of the luminescent photoprotein from
- 15 the partially denatured apoprotein," *Biochem. J.* 199:825-828; Inouye *et al.* (1989) *J. Biochem.* 105:473-477; Inouye *et al.* (1986) "Expression of Apoequorin Complementary DNA in *Escherichia coli*," *Biochemistry* 25:8425-8429; Inouye *et al.* (1985) "Cloning and sequence analysis of cDNA for the luminescent protein aequorin," *Proc. Natl. Acad. Sci. USA* 82:3154-3158;
- 20 Prendergast, *et al.* (1978) "Chemical and Physical Properties of Aequorin and the Green Fluorescent Protein Isolated from *Aequorea forskalea*" *J. Am. Chem. Soc.* 100:3448-3453; European Patent Application 0 540 064 A1; European Patent Application 0 226 979 A2, European Patent Application 0 245 093 A1 and European Patent Application 0 245 093 B1; U.S. Patent No. 5,093,240; U.S.
- 25 Patent No. 5,360,728; U.S. Patent No. 5,139,937; U.S. Patent No. 5,422,266; U.S. Patent No. 5,023,181; U.S. Patent No. 5,162,227; and SEQ ID Nos. 5-13, which set forth DNA encoding the apoprotein; and a form, described in U.S. Patent No. 5,162,227, European Patent Application 0 540 064 A1 and Sealite Sciences Technical Report No. 3 (1994), is commercially available from Sealite,
- 30 Sciences, Bogart, GA as AQUALITE®).

This system is among the preferred systems for use herein. As will be evident, since the aequorin photoprotein includes noncovalently bound luciferin

-60-

and molecular oxygen, it is suitable for storage in this form as a lyophilized powder or encapsulated into a selected delivery vehicle. The system can be encapsulated into pellets, such as liposomes or other delivery vehicles. When used, the vehicles are contacted with a composition, even tap water, that

5 contains Ca^{2+} (or other suitable metal ion), to produce a mixture that glows.

a. **Aequorin and related photoproteins**

The photoprotein, aequorin, isolated from the jellyfish, *Aequorea*, emits light upon the addition of Ca^{2+} (or other suitable metal ion). The aequorin photoprotein, which includes bound luciferin and bound oxygen that is released

10 by Ca^{2+} , does not require dissolved oxygen. Luminescence is triggered by calcium, which releases oxygen and the luciferin substrate producing apoaequorin.

The bioluminescence photoprotein aequorin is isolated from a number of species of the jellyfish *Aequorea*. It is a 22 kilodalton (kD) molecular weight

15 peptide complex (see, e.g., Shimomura *et al.* (1962) *J. Cellular and Comp. Physiol.* 59:233-238; Shimomura *et al.* (1969) *Biochemistry* 8:3991-3997; Kohama *et al.* (1971) *Biochemistry* 10:4149-4152; and Shimomura *et al.* (1972) *Biochemistry* 11:1602-1608). The native protein contains oxygen and a heterocyclic compound coelenterazine, a luciferin, (see, below) noncovalently

20 bound thereto. The protein contains three calcium binding sites. Upon addition of trace amounts Ca^{2+} (or other suitable metal ion, such as strontium) to the photoprotein, it undergoes a conformational change that catalyzes the oxidation of the bound coelenterazine using the protein-bound oxygen. Energy from this oxidation is released as a flash of blue light, centered at 469 nm.

25 Concentrations of calcium ions as low as 10^{-6} M are sufficient to trigger the oxidation reaction.

Naturally-occurring apoaequorin is not a single compound but rather is a mixture of microheterogeneous molecular species. *Aequoria* jellyfish extracts contain as many as twelve distinct variants of the protein (see, e.g., Prasher *et al.* (1975) *Biochemistry* 26:1326-1332; Blinks *et al.* (1975) *Fed. Proc.* 34:474).

30 DNA encoding numerous forms has been isolated (see, e.g., SEQ ID Nos. 5-9 and 13).

-61-

The photoprotein can be reconstituted (see, *e.g.*, U.S. Patent No. 5,023,181) by combining the apoprotein, such as a protein recombinantly produced in *E. coli*, with a coelenterazine, such as a synthetic coelenterazine, in the presence of oxygen and a reducing agent (see, *e.g.*, Shimomura *et al.* (1975) *Nature* 256:236-238; Shimomura *et al.* (1981) *Biochemistry J.* 199:825-828), such as 2-mercaptoethanol, and also EDTA or EGTA (concentrations between about 5 to about 100 mM or higher for applications herein) tie up any Ca^{2+} to prevent triggering the oxidation reaction until desired. DNA encoding a modified form of the apoprotein that does not require 2-mercaptoethanol for reconstitution is also available (see, *e.g.*, U.S. Patent No. 5,093,240). The reconstituted photoprotein is also commercially available (sold, *e.g.*, under the trademark AQUALITE[®], which is described in U.S. Patent No. 5,162,227).

The light reaction is triggered by adding Ca^{2+} at a concentration sufficient to overcome the effects of the chelator and achieve the 10^{-6} M concentration. Because such low concentrations of Ca^{2+} can trigger the reaction, for use in the methods herein, higher concentrations of chelator may be included in the compositions of photoprotein. Accordingly, higher concentrations of added Ca^{2+} in the form of a calcium salt will be required. Precise amounts may be empirically determined. For use herein, it may be sufficient to merely add water to the photoprotein, which is provided in the form of a concentrated composition or in lyophilized or powdered form. Thus, for purposes herein, addition of small quantities of Ca^{2+} , such as those present in phosphate buffered saline (PBS) or other suitable buffers or the moisture on the tissue to which the compositions are contacted, should trigger the bioluminescence reaction.

Numerous isoforms of the aequorin apoprotein been identified isolated. DNA encoding these proteins has been cloned, and the proteins and modified forms thereof have been produced using suitable host cells (see, *e.g.*, U.S. Patent Nos. 5,162,227, 5,360,728, 5,093,240; see, also, Prasher *et al.* (1985) *Biophys. Biochem. Res. Commun.* 126:1259-1268; Inouye *et al.* (1986) *Biochemistry* 25:8425-8429). U.S. Patent No. 5,093,240; U.S. Patent No. 5,360,728; U.S. Patent No. 5,139,937; U.S. Patent No. 5,288,623; U.S. Patent No. 5,422,266, U.S. Patent No. 5,162,227 and SEQ ID Nos. 5-13, which set

-62-

forth DNA encoding the apoprotein; and a form is commercially available from Sealite, Sciences, Bogart, GA as AQUALITE®). DNA encoding apoaequorin or variants thereof is useful for recombinant production of high quantities of the apoprotein. The photoprotein is reconstituted upon addition of the luciferin, coelenterazine, preferably a sulfated derivative thereof, or an analog thereof, and molecular oxygen (see, *e.g.*, U.S. Patent No. 5,023,181). The apoprotein and other constituents of the photoprotein and bioluminescence generating reaction can be mixed under appropriate conditions to regenerate the photoprotein and concomitantly have the photoprotein produce light. Reconstitution requires the presence of a reducing agent, such as mercaptoethanol, except for modified forms, discussed below, that are designed so that a reducing agent is not required (see, *e.g.*, U.S. Patent No. 5,093,240).

For use herein, it is preferred aequorin is produced using DNA, such as that set forth in SEQ ID Nos. 5-13 and known to those of skill in the art or modified forms thereof. The DNA encoding aequorin is expressed in a host cell, such as *E. coli*, isolated and reconstituted to produce the photoprotein (see, *e.g.*, U.S. Patent Nos. 5,418,155, 5,292,658, 5,360,728, 5,422,266, 5,162,227).

Of interest herein, are forms of the apoprotein that have been modified so that the bioluminescent activity is greater than unmodified apoaequorin (see, *e.g.*, U.S. Patent No. 5,360,728, SEQ ID Nos. 10-12). Modified forms that exhibit greater bioluminescent activity than unmodified apoaequorin include proteins including sequences set forth in SEQ ID Nos. 10-12, in which aspartate 124 is changed to serine, glutamate 135 is changed to serine, and glycine 129 is changed to alanine, respectively. Other modified forms with increased bioluminescence are also available.

For use in certain embodiments herein, the apoprotein and other components of the aequorin bioluminescence generating system are packaged or provided as a mixture, which, when desired is subjected to conditions under which the photoprotein reconstitutes from the apoprotein, luciferin and oxygen (see, *e.g.*, U.S. Patent No. 5,023,181; and U.S. Patent No. 5,093,240). Particularly preferred are forms of the apoprotein that do not require a reducing agent, such as 2-mercaptoethanol, for reconstitution. These forms, described,

-63-

for example in U.S. Patent No. 5,093,240 (see, also Tsuji *et al.* (1986) *Proc. Natl. Acad. Sci. U.S.A.* 83:8107-8111), are modified by replacement of one or more, preferably all three cysteine residues with, for example serine.


Replacement may be effected by modification of the DNA encoding the aequorin apoprotein, such as that set forth in SEQ ID No. 5, and replacing the cysteine codons with serine.

The photoproteins and luciferases from related species, such as *Obelia* are also contemplated for use herein. DNA encoding the Ca^{2+} -activated photoprotein obelin from the hydroid polyp *Obelia longissima* is known and available (see, *e.g.*, Illarionov *et al.* (1995) *Gene* 153:273-274; and Bondar *et al.* (1995) *Biochim. Biophys. Acta* 1231:29-32). This photoprotein can also be activated by Mn^{2+} (see, *e.g.*, Vysotski *et al.* (1995) *Arch. Bioch. Biophys.* 316:92-93, Vysotski *et al.* (1993) *J. Biolumin. Chemilumin.* 8:301-305).

In general for use herein, the components of the bioluminescence are packaged or provided so that there is insufficient metal ions to trigger the reaction. When used, the trace amounts of triggering metal ion, particularly Ca^{2+} is contacted with the other components. For a more sustained glow, aequorin can be continuously reconstituted or can be added or can be provided in high excess.

20 b. Luciferin

The aequorin luciferin is coelenterazine and analogs therein, which include molecules including the structure (formula (I)):




in which R_1 is $\text{CH}_2\text{C}_6\text{H}_5$ or CH_3 ; R_2 is C_6H_5 , and R_3 is *p*- $\text{C}_6\text{H}_4\text{OH}$ or CH_3 or other such analogs that have activity. Preferred coelenterazine has the structure in which R_1 is *p*- $\text{CH}_2\text{C}_6\text{H}_4\text{OH}$, R_2 is C_6H_5 , and R_3

-64-

is $p\text{-C}_6\text{H}_4\text{OH}$, which can be prepared by known methods (see, e.g., Inouye *et al.* (1975) *Jap. Chem. Soc., Chemistry Ltrrs.* pp 141-144; and Hart *et al.* (1979) *Biochemistry* 18:2204-2210). Among the preferred analogs, are those that are modified, whereby the spectral frequency of the resulting light is shifted to


5 another frequency.

The preferred coelenterazine has the structure (II):



and sulfated derivatives thereof.

Another coelenterazine has formula (V):




(see, Hart *et al.* (1979)

35 *Biochemistry* 18:2204-2210). Using this derivative in the presence of luciferase all of the light is in the ultraviolet with a peak at 390 nm. Upon addition of GFP, all light emitted is now in the visible range with a peak at 509 nm accompanied by an about 200-fold increase in the amount of light emitted. Viewed with a cut-off filter of 470 nm, in the light yield in the absence of GFP would be about
40 zero, and would be detectable in the presence of GFP. This provides the basis for an immunoassay described in the EXAMPLES.

-65-

The reaction of coelenterazine when bound to the aequorin photoprotein with bound oxygen and in the presence of Ca^{2+} can be represented as follows:



15

The photoprotein aequorin (which contains apoaequorin bound to a coelenterate luciferin molecule) and *Renilla* luciferase, discussed below, can use the same coelenterate luciferin. The aequorin photoprotein catalyses the oxidation of coelenterate luciferin (coelenterazine) to oxyluciferin

20 (coelenteramide) with the concomitant production of blue light ($\lambda_{\text{max}} = 469 \text{ nm}$).

Importantly, the sulfate derivative of the coelenterate luciferin (lauryl-luciferin) is particularly stable in water, and thus may be used in a coelenterate-like bioluminescent system. In this system, adenosine diphosphate (ADP) and a

25 sulphate-kinase are used to convert the coelenterazine to the sulphated form. Sulfatase is then used to reconvert the lauryl-luciferin to the native coelenterazine. Thus, the more stable lauryl-luciferin is used in the item to be illuminated and the luciferase combined with the sulfatase are added to the luciferin mixture when illumination is desired.

30 Thus, the bioluminescent system of *Aequorea* is particularly suitable for use in the methods herein. The particular amounts and the manner in which the components are provided depends upon the type of neoplasia or specialty tissue to be visualized. This system can be provided in lyophilized form, that will glow upon addition of Ca^{2+} . It can be encapsulated, linked to microcarriers, such as

35 microbeads, or in as a compositions, such as a solution or suspension, preferably in the presence of sufficient chelating agent to prevent triggering the reaction.

-66-

The concentration of the aequorin photoprotein will vary and can be determined empirically. Typically concentrations of at least 0.1 mg/l, more preferably at least 1 mg/l and higher, will be selected. In certain embodiments, 1-10 mg luciferin/100 mg of luciferase will be used in selected volumes and at the desired concentrations will be used.

5. Crustacean, particularly *Cypridina* systems

The ostracods, such as *Vargula serratta*, *hilgendorffii* and *noctiluca* are small marine crustaceans, sometimes called sea fireflies. These sea fireflies are found in the waters off the coast of Japan and emit light by squirting luciferin and luciferase into the water, where the reaction, which produces a bright blue luminous cloud, occurs. The reaction involves only luciferin, luciferase and molecular oxygen, and, thus, is very suitable for application herein.

The systems, such as the *Vargula* bioluminescent systems, are particularly preferred herein because the components are stable at room temperature if dried and powdered and will continue to react even if contaminated. Further, the bioluminescent reaction requires only the luciferin/luciferase components in concentrations as low as 1:40 parts per billion to 1:100 parts per billion, water and molecular oxygen to proceed. An exhausted system can be renewed by addition of luciferin.

a. *Vargula* luciferase

The *Vargula* luciferase is water soluble and is among those preferred for use in the methods herein. *Vargula* luciferase is a 555-amino acid polypeptide that has been produced by isolation from *Vargula* and also using recombinant technology by expressing the DNA in suitable bacterial and mammalian host cells (see, e.g., Thompson *et al.* (1989) *Proc. Natl. Acad. Sci. U.S.A.* 86:6567-6571; Inouye *et al.* (1992) *Proc. Natl. Acad. Sci. U.S.A.* 89:9584-9587; Johnson *et al.* (1978) *Methods in Enzymology* LVII:331-349; Tsuji *et al.* (1978) *Methods Enzymol.* 57:364-72; Tsuji (1974) *Biochemistry* 13:5204-5209; Japanese Patent Application No. JP 3-30678 Osaka; and European Patent Application No. EP 0 387 355 A1).

-67-

(1) Purification from *Cypridina*

Methods for purification of *Vargula* (*Cypridina*) luciferase are well known. For example, crude extracts containing the active can be readily prepared by grinding up or crushing the *Vargula* shrimp. In other

5 embodiments, a preparation of *Cypridina hilgendorfi* luciferase can be prepared by immersing stored frozen *C. hilgendorfi* in distilled water containing, 0.5-5.0 M salt, preferably 0.5-2.0 M sodium or potassium chloride, ammonium sulfate, at 0-30° C, preferably 0-10° C, for 1-48 hr, preferably 10-24 hr, for extraction followed by hydrophobic chromatography and then ion exchange or affinity

10 chromatography (TORAY IND INC, Japanese patent application JP 4258288, published September 14, 1993; see, also, Tsuji *et al.* (1978) *Methods Enzymol.* 57:364-72 for other methods).

(2) Preparation by Recombinant Methods


The luciferase is preferably produced by expression of cloned DNA

15 encoding the luciferase (European Patent Application No. 0 387 355 A1; International PCT Application No. WO 95/001542; see, also SEQ ID No. 5, which sets forth the sequence from Japanese Patent Application No. JP 3-30678 and Thompson *et al.* (1989) *Proc. Natl. Acad. Sci. U.S.A.* 86:6567-6571) DNA encoding the luciferase or variants thereof is introduced into *E. coli*

20 using appropriate vectors and isolated using standard methods.

b. *Vargula* luciferin

The natural luciferin is a substituted imidazopyrazine nucleus, such a compound of formula (III):



-68-

The luciferin can be isolated from ground dried *Vargula* by heating the extract, which destroys the luciferase but leaves the luciferin intact (see, *e.g.*, U.S. Patent No. 4,853,327).

5 Analogous thereof and other compounds that react with the luciferase in a light producing reaction also may be used.

Other bioluminescent organisms that have luciferases that can react with the *Vargula* luciferin include, the genera *Apogon*, *Parapriacanthus* and *Porichthys*.

c. Reaction

10 The luciferin upon reaction with oxygen forms a dioxetanone intermediate (which includes a cyclic peroxide similar to the firefly cyclic peroxide molecule intermediate). In the final step of the bioluminescent reaction, the peroxide breaks down to form CO₂ and an excited carbonyl. The excited molecule then emits a blue to blue-green light.

15 The optimum pH for the reaction is about 7. For purposes herein, any pH at which the reaction occurs may be used. The concentrations of reagents are those normally used for analytical reactions or higher (see, *e.g.*, Thompson *et al.* (1990) *Gene* 96:257-262). Typically concentrations of the luciferase between 0.1 and 10 mg/l, preferably 0.5 to 2.5 mg/l will be used. Similar concentrations
20 or higher concentrations of the luciferin may be used.

6. Insect bioluminescent systems including fireflies, click beetles, and other insect system

The biochemistry of firefly bioluminescence was the first bioluminescent system to be characterized (see, *e.g.*,

25 Wienhausen *et al.* (1985) *Photochemistry and Photobiology* 42:609-611; McElroy *et al.* (1966) in *Molecular Architecture in cell Physiology*, Hayashi *et al.*, eds. Prentice Hall, Inc., Englewood Cliffs, NJ, pp. 63-80) and it is commercially available (*e.g.*, from Promega Corporation, Madison, WI, see, *e.g.*, Leach *et al.* (1986) *Methods in Enzymology* 133:51-70, esp. Table 1). Luciferases from
30 different species of fireflies are antigenically similar. These species include members of the genera *Photinus*, *Photurins* and *Luciola*. Further, the bioluminescent reaction produces more light at 30°C than at 20°C, the luciferase is

-69-

stabilized by small quantities of bovine albumin serum, and the reaction can be buffered by tricine.

a. Luciferase

DNA clones encoding luciferases from various insects and the use to
5 produce the encoded luciferase is well known. For example, DNA clones that
encode luciferase from *Photinus pyralis*, *Luciola cruciata* (see, *e.g.*, de Wet *et al.*
(1985) *Proc. Natl. Acad. Sci. U.S.A.* 82:7870-7873; de We *et al.* (1986)
Methods in Enzymology 133:3; U.S. Patent No. 4,968,613, see, also SEQ ID
No. 3) are available. The DNA has also been expressed in *Saccharomyces* (see,
10 *e.g.*, Japanese Application No. JP 63317079, published December 26, 1988,
KIKKOMAN CORP) and in tobacco.

In addition to the wild-type luciferase modified insect luciferases have
been prepared. For example, heat stable luciferase mutants, DNA-encoding the
mutants, vectors and transformed cells for producing the luciferases are
15 available. A protein with 60% amino acid sequence homology with luciferases
from *Photinus pyralis*, *Luciola mingrelica*, *L. cruciata* or *L. lateralis* and having
luciferase activity is available (see, *e.g.*, International PCT Application No. WO
95/25798). It is more stable above 30° C than naturally-occurring insect
luciferases and may also be produced at 37° C or above, with higher yield.


20 Modified luciferases can generate light at different wavelengths
(compared with native luciferase), and thus, may be selected for their color-
producing characteristics. For example, synthetic mutant beetle luciferase(s) and
DNA encoding such luciferases that produce bioluminescence at a wavelength
different from wild-type luciferase are known (Promega Corp, International PCT
25 Application No. WO 95/18853, which is based on U.S. application Serial No.
08/177,081). The mutant beetle luciferase has an amino acid sequence differing
from that of the corresponding wild-type *Luciola cruciata* (see, *e.g.*, U.S. Patent
Nos. 5,182,202, 5,219,737, 5,352,598, see, also SEQ ID No.3) by a
substitution(s) at one or two positions. The mutant luciferase produces a
30 bioluminescence with a wavelength of peak intensity that differs by at least 1
nm from that produced by wild-type luciferases.

-70-

- Other mutant luciferases can be produced. Mutant luciferases with the amino acid sequence of wild-type luciferase, but with at least one mutation in which valine is replaced by isoleucine at the amino acid number 233, valine by isoleucine at 239, serine by asparagine at 286, glycine by serine at 326, histidine by tyrosine at 433 or proline by serine at 452 are known (see, e.g., U.S. Patent Nos. 5,219,737, and 5,330,906). The luciferases are produced by expressing DNA-encoding each mutant luciferase in *E. coli* and isolating the protein. These luciferases produce light with colors that differ from wild-type. The mutant luciferases catalyze luciferin to produce red (λ 609 nm and 612 nm), orange (λ 595 and 607 nm) or green (λ 558 nm) light. The other physical and chemical properties of mutant luciferase are substantially identical to native wild type-luciferase. The mutant luciferase has the amino acid sequence of *Luciola cruciata* luciferase with an alteration selected from Ser 286 replaced by Asn, Gly 326 replaced by Ser, His 433 replaced by Tyr or Pro 452 replaced by Ser.
- Thermostable luciferases are also available (see, e.g., U.S. Patent No. 5,229,285; see, also International PCT Application No. WO 95/25798, which provides *Photinus* luciferase in which the glutamate at position 354 is replaced with lysine and *Luciola* luciferase in which the glutamate at 356 is replaced with lysine).
- These mutant luciferases as well as the wild type luciferases can be used in combination with the GFPs provided herein particularly in instances when a variety of colors are desired or when stability at higher temperatures is desired.


b. Luciferin

The firefly luciferin is a benzothiazole:



Analogues of this luciferin and synthetic firefly luciferins are also known to those of skill in art (see, e.g., U.S. Patent No. 5,374,534 and 5,098,828). These include compounds of formula (IV) (see, U.S. Patent No. 5,098,828):

-71-



in which:

- 10 R^1 is hydroxy, amino, linear or branched C_1 - C_{20} alkoxy, C_2 - C_{20} alkenyloxy, an L-amino acid radical bond via the α -amino group, an oligopeptide radical with up to ten L-amino acid units linked via the α -amino group of the terminal unit;

- R^2 is hydrogen, H_2PO_3 , HSO_3 , unsubstituted or phenyl substituted linear
15 or branched C_1 - C_{20} alkyl or C_2 - C_{20} alkenyl, aryl containing 6 to 18 carbon atoms, or R^3 -C(O)-; and

- R^3 is an unsubstituted or phenyl substituted linear or branched C_1 - C_{20} alkyl or C_2 - C_{20} alkenyl, aryl containing 6 to 18 carbon atoms, a nucleotide radical with 1 to 3 phosphate groups, or a glycosidically attached mono- or
20 disaccharide, except when formula (IV) is a D-luciferin or D-luciferin methyl ester.

Modified luciferins that have been modified to produce light of shifted frequencies are known to those of skill in the art.

c. Reaction

- 25 The reaction catalyzed by firefly luciferases and related insect luciferases requires ATP, Mg^{2+} as well as molecular oxygen. Luciferin must be added exogenously. Firefly luciferase catalyzes the firefly luciferin activation and the subsequent steps leading to the excited product. The luciferin reacts with ATP to form a luciferyl adenylate intermediate. This intermediate then reacts with
30 oxygen to form a cyclic luciferyl peroxy species, similar to that of the coelenterate intermediate cyclic peroxide, which breaks down to yield CO_2 and an excited state of the carbonyl product. The excited molecule then emits a yellow light; the color, however, is a function of pH. As the pH is lowered the color of the bioluminescence changes from yellow-green to red.

-72-

Different species of fireflies emit different colors of bioluminescence so that the color of the reaction will be dependent upon the species from which the luciferase is obtained. Additionally, the reaction is optimized at pH 7.8.

5 Addition of ATP and luciferin to a reaction that is exhausted produces additional light emission. Thus, the system, once established, is relatively easily maintained. Therefore, it is highly suitable for use herein in embodiments in which a sustained glow is desired.

7. Other systems

Numerous other systems are known and have been described in detail for
10 example in U.S. Patent Nos. 5,876,995, 6,152,358 and 6,113,886).

a. Bacterial systems

Luminous bacteria typically emit a continuous light, usually blue-green. When strongly expressed, a single bacterium may emit 10^4 to 10^5 photons per second. Bacterial bioluminescence systems include, among others, those
15 systems found in the bioluminescent species of the genera *Photobacterium*, *Vibrio* and *Xenorhabdus*. These systems are well known and well characterized (see, e.g., Baldwin *et al.* (1984) *Biochemistry* 23:3663-3667; Nicoli *et al.* (1974) *J. Biol. Chem.* 249:2393-2396; Welches *et al.* (1981) *Biochemistry* 20:512-517; Engebrecht *et al.* (1986) *Methods in Enzymology* 133:83-99; Frackman *et al.*
20 (1990) *J. of Bacteriology* 172:5767-5773; Miyamoto *et al.* (1986) *Methods in Enzymology* 133:70; U.S. Patent No. 4,581,335).

(1) Luciferases

Bacterial luciferase, as exemplified by luciferase derived from *Vibrio harveyi* (EC 1.14.14.3, alkanol reduced-FMN-oxygen oxidoreductase 1-hydroxylating,
25 luminescing), is a mixed function oxidase, formed by the association of two different protein subunits α and β . The α -subunit has an apparent molecular weight of approximately 42,000 kDa and the β -subunit has an apparent molecular weight of approximately 37,000 kDa (see, e.g., Cohn *et al.* (1989) *Proc. Natl. Acad. Sci. U.S.A.* 90:102-123). These subunits associate to form a
30 2-chain complex luciferase enzyme, which catalyzes the light emitting reaction of bioluminescent bacteria, such as *Vibrio harveyi* (U.S. Patent No. 4,581,335; Belas *et al.* (1982) *Science* 218:791-793), *Vibrio fischeri* (Engebrecht *et al.*

-73-

(1983) *Cell* 32:773-781; Engebrecht *et al.* (1984) *Proc. Natl. Acad. Sci. U.S.A.* 81:4154-4158) and other marine bacteria.

Bacterial luciferase genes have been cloned (see, *e.g.*, U.S. Patent No. 5,221,623; U.S. Patent No. 4,581,335; European Patent Application No. EP 386 5 691 A). Plasmids for expression of bacterial luciferase, such as *Vibrio harveyi*, include pFIT001 (NRRL B-18080), pPALE001 (NRRL B-18082) and pMR19 (NRRL B-18081)) are known. For example the sequence of the entire *lux* regulon from *Vibrio fischeri* has been determined (Baldwin *et al.* (1984), *Biochemistry* 23:3663-3667; Baldwin *et al.* (1981) *Biochem.* 20:512-517; Baldwin *et al.* 10 (1984) *Biochem.* 23:3663-3667; see, also, *e.g.*, U.S. Patent Nos. 5,196,318, 5,221,623, and 4,581,335). This regulon includes *luxI* gene, which encodes a protein required for autoinducer synthesis (see, *e.g.*, Engebrecht *et al.* (1984) *Proc. Natl. Acad. Sci. U.S.A.* 81:4154-4158), the *luxC*, *luxD*, and *luxE* genes, which encode enzymes that provide the luciferase with an aldehyde substrate, 15 and the *luxA* and *luxB* genes, which encode the alpha and beta subunits of the luciferase.


Lux genes from other bacteria have also been cloned and are available (see, *e.g.*, Cohn *et al.* (1985) *J. Biol. Chem.* 260:6139-6146; U.S. Patent No. 5,196,524, which provides a fusion of the *luxA* and *luxB* genes from *Vibrio* 20 *harveyi*). Thus, luciferase alpha and beta subunit-encoding DNA is provided and can be used to produce the luciferase. DNA encoding the α (1065 bp) and β (984 bp) subunits, DNA encoding a luciferase gene of 2124 bp, encoding the alpha and beta subunits, a recombinant vector containing DNA encoding both subunits and a transformed *E. coli* and other bacterial hosts for expression and 25 production of the encoded luciferase are available. In addition, bacterial luciferases are commercially available.

(2) Luciferins

Bacterial luciferins include:

30

-74-




- 15 in which the tetradecanal with reduced flavin mononucleotide are considered luciferin since both are oxidized during the light emitting reaction.

(3) Reactions

The bacterial systems require, in addition to reduced flavin, five polypeptides to complete the bioluminescent reaction: two subunits, α and β , of bacterial luciferin and three units of a fatty acid reductase system complex, which supplies the tetradecanal aldehyde. Examples of bacterial bioluminescent systems useful in the apparatus and methods provided herein include those derived from *Vibrio fischeri* and *Vibrio harveyi*. One advantage to this system is its ability to operate at cold temperatures; certain surgical procedures are performed by cooling the body to lower temperatures.

Bacterial luciferase catalyzes the flavin-mediated hydroxylation of a long-chain aldehyde to yield carboxylic acid and an excited flavin; the flavin decays to ground state with the concomitant emission of blue green light ($\lambda_{\text{max}} = 490 \text{ nm}$; see, e.g., Legocki *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 81:9080; see U.S. Patent No. 5,196,524):



The reaction can be initiated by contacting reduced flavin mononucleotide (FMNH₂) with a mixture of the bacterial luciferase, oxygen, and a long-chain aldehyde, usually n-decyl aldehyde.

DNA encoding luciferase from the fluorescent bacterium *Alteromonas hanedai* is known (CHISSO CORP; see, also, Japanese application JP 7222590,

-75-

published August 22, 1995). The reduced flavin mononucleotide (FMNH₂; luciferin) reacts with oxygen in the presence of bacterial luciferase to produce an intermediate peroxy flavin. This intermediate reacts with a long-chain aldehyde (tetradecanal) to form the acid and the luciferase-bound hydroxy flavin in its
5 excited state. The excited luciferase-bound hydroxy flavin then emits light and dissociates from the luciferase as the oxidized flavin mononucleotide (FMN) and water. *In vivo* FMN is reduced again and recycled, and the aldehyde is regenerated from the acid.


Flavin reductases have been cloned (see, *e.g.*, U.S. Patent No.
10 5,484,723; see, SEQ ID No. 14 for a representative sequence from this patent). These as well as NAD(P)H can be included in the reaction to regenerate FMNH₂ for reaction with the bacterial luciferase and long chain aldehyde. The flavin reductase catalyzes the reaction of FMN, which is the luciferase reaction, into FMNH₂; thus, if luciferase and the reductase are included in the reaction system,
15 it is possible to maintain the bioluminescent reaction. Namely, since the bacterial luciferase turns over many times, bioluminescence continues as long as a long chain aldehyde is present in the reaction system.

The color of light produced by bioluminescent bacteria also results from the participation of a protein blue-florescent protein (BFP) in the bioluminescence
20 reaction. This protein, which is well known (see, *e.g.*, Lee *et al.* (1978) *Methods in Enzymology LVII*:226-234), may also be added to bacterial bioluminescence reactions in order to cause a shift in the color.

b. Dinoflagellate bioluminescence generating systems

In dinoflagellates, bioluminescence occurs in organelles termed scintillons.
25 These organelles are outpocketings of the cytoplasm into the cell vacuole. The scintillons contain only dinoflagellate luciferase and luciferin (with its binding protein), other cytoplasmic components being somehow excluded. The dinoflagellate luciferin is a tetrapyrrole related to chlorophyll:

-76-



10

or an analog thereof.

The luciferase is a 135 kD single chain protein that is active at pH 6.5, but inactive at pH 8 (see, *e.g.*, Hastings (1981) *Bioluminescence and Chemiluminescence*, DeLuca *et al.*, eds. Academic Press, NY, pp.343-360).

15 Luminescent activity can be obtained in extracts made at pH 8 by simply shifting the pH from 8 to 6. This occurs in soluble and particulate fractions. Within the intact scintillon, the luminescent flash occurs for ~100 msec, which is the duration of the flash *in vivo*. In solution, the kinetics are dependent on dilution, as in any enzymatic reaction. At pH 8, the luciferin is bound to a protein
20 (luciferin binding protein) that prevents reaction of the luciferin with the luciferase. At pH 6, however, the luciferin is released and free to react with the enzyme.

D. ISOLATION AND IDENTIFICATION OF NUCLEIC ACIDS ENCODING LUCIFERASES AND GFPs

25 Nucleic acid encoding bioluminescent proteins are provided. Particularly, nucleic acid encoding *Renilla reniformis* GFP is provided.

1. Isolation of specimens of the genus *Renilla*

Specimens of *Renilla* are readily available from the oceans of the world, including the Gulf of Mexico, Pacific Ocean and Atlantic Ocean. *Renilla* typically
30 live on the ocean bottom at about 30 to 100 feet deep and can be easily collected by dragging. For example, specimens of *R. kollikeri* can be obtained off the coast of California or Baja, Mexico. Alternatively, live specimens of *Renilla* may be purchased from a commercial supplier (*e.g.*, Gulf Marine Incorporated, Panacea, Fla). Upon capture or receipt, the specimens are washed
35 thoroughly and may also be dissected to enrich for light-emitting tissues. The

-77-

whole organisms or dissected tissues are then snap frozen and stored in liquid nitrogen.

As described in detail in the examples below, the frozen tissues were used as a source to isolate nucleic acids encoding *Renilla mulleri* GFP and luciferase (*e.g.*, see SEQ ID No. 15 and SEQ ID No. 17, respectively).

2. Preparation of *Renilla* cDNA expression libraries

Renilla cDNA expression libraries may be prepared from intact RNA following the methods described herein or by other methods known to those of skill the art (*e.g.*, see Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.; U.S. Patent No. 5,292,658).

Typically, the preparation of cDNA libraries includes the isolation of polyadenylated RNA from the selected organism followed by single-strand DNA synthesis using reverse transcriptase, digestion of the RNA strand of the DNA/RNA hybrid and subsequent conversion of the single-stranded DNA to double stranded cDNA.

a. RNA isolation and cDNA synthesis

Whole *Renilla* or dissected *Renilla* tissues can be used a source of total cytoplasmic RNA for the preparation of *Renilla* cDNA. Total intact RNA can be isolated from crushed *Renilla* tissue, for example, by using a modification of methods generally known in the art (*e.g.*, see Chirgwin *et al.* (1970) *Biochemistry* 18:5294-5299). After isolating total cellular RNA, polyadenylated RNA species are then easily separated from the nonpolyadenylated species using affinity chromatography on oligodeoxythymidylate cellulose columns, (*e.g.*, as described by Aviv *et al.*, (1972) *Proc. Natl. Acad. Sci. U.S.A.* 69:1408).

The purified *Renilla* polyA-mRNA is then subjected to a cDNA synthesis reaction to generate a cDNA library from total polyA-mRNA. Briefly, reverse transcriptase is used to extend an annealed polydT primer to generate an RNA/DNA duplex. The RNA strand is then digested using an RNase, *e.g.*, RNase H, and following second-strand synthesis, the cDNA molecules are blunted-ended with S1 nuclease or other appropriate nuclease. The resulting double-stranded cDNA fragments can be ligated directly into a suitable expression

-78-

vector or, alternatively, oligonucleotide linkers encoding restriction endonuclease sites can be ligated to the 5'-ends of the cDNA molecules to facilitate cloning of the cDNA fragments.

b. Construction of cDNA expression libraries

5 The best characterized vectors for the construction of cDNA expression libraries are lambda vectors. Lambda-based vectors tolerate cDNA inserts of about 12 kb and provide greater ease in library screening, amplification and storage compared to standard plasmid vectors. Presently preferred vectors for the preparation of *Renilla* cDNA expression libraries are the Lambda, Uni-Zap, 10 Lambda-Zap II or Lambda-ZAP Express/*EcoRI/XhoI* vectors, which are known to those of skill in the art (*e.g.*, see U.S. Pat. No. 5,128,256), and are also commercially available (Stratagene, La Jolla, CA).

Generally, the Lambda-Zap vectors combine the high efficiency of a bacteriophage lambda vector systems with the versatility of a plasmid system. 15 Fragments cloned into these vectors can be automatically excised using a helper phage and recircularized to generate subclones in the pBK-derived phagemid. The pBK phagemid carries the neomycin-resistance gene for selection in bacteria and G418 selection in eukaryotic cells or may contain the β -lactamase resistance gene. Expression of the recombinant polypeptide is under the control of the *lacZ* 20 promoter in bacteria and the CMV promoter in eukaryotes.

More specifically, these lambda-based vectors are composed of an initiator-terminator cassette containing the plasmid system, *e.g.*, a pBK Bluescript derivative (Stratagene, San Diego), bracketed by the right and left arm of the bacteriophage lambda. The lambda arms allow for efficient packaging of 25 replicated DNA whereas the excisable initiator-terminator cassette allows for easy cloning of the cDNA fragments and the generation of a plasmid library without the need for additional subcloning.

When used herein, cDNA fragments are inserted into the multiple cloning site contained within the initiator-terminator cassette of the Lambda-Zap vector 30 to create a set of cDNA expression vectors. The set of cDNA expression vectors is allowed to infect suitable *E. coli* cells, followed by co-infection with a filamentous helper phage. Within the cell, trans-acting proteins encoded by the

-79-

helper phage, *e.g.*, the gene II protein of M13, recognize two separate domains positioned within the lambda arms of the vector and introduce single-stranded nicks flanking the initiator-terminator cassette. Upon a subsequent round of DNA synthesis, a new DNA strand is synthesized that displaces the existing nick strand liberating the initiator-terminator cassette. The displaced strand is then circularized, packaged as filamentous phage by the helper proteins and excreted from the cell. The BK plasmid containing the cDNA is recovered by infecting an F' strain of *E. coli* and plating the infected cells on solid medium supplemented with kanamycin for the selection of pBK-containing cells.

10 The *Renilla* cDNA expression library can be screened using a variety of methods known to those of skill in the art. For example, identification of *Renilla* GFP may be achieved using a functional screening method employing blue light and observing colonies visually for emission of green fluorescence or by observing light emission using one or more bandpass filter.

15 **3. Cloning of *Renilla reniformis* Green Fluorescent Protein**

Renilla reniformis GFP has 233 amino acids compared to GFPs from animals that contain luciferase-GFP bioluminescent systems *Renilla mulleri*, *Ptilosarcus* and *Aequorea Victoria*. Other such GFPs have 238 amino acids. At the amino acid level, *Renilla reniformis* is respectively 53, 51 and 19% identical to the GFPs from these animals. The extent of identity of *Renilla reniformis* GFP to the half dozen cloned anthozoan coral GFPs, which do not contain associated luciferases, ranges from 32 to 38%. The overall identity among these GFPs is surprisingly low for a protein evolved from a common ancestor. These relationships are depicted as a phylogenetic tree (Figure 1).

25 Most surprising is the finding that the *Renilla reniformis* GFP is much more closely related to *Ptilosarcus* GFP (77% identity) than to *Renilla reniformis* GFP (53%). It is unclear why the sequence relatedness between these 3 GFPs does not follow traditional taxonomy. Given the sequence differences at the amino acid level, coding DNA sequences are surprisingly well conserved. *Renilla reniformis* GFP DNA is 56 and 59% identical to *Renilla mulleri* and *Ptilosarcus* GFP DNA.

-80-

Thus cloning *Renilla reniformis* GFP clone suggests why many groups may have failed in attempts to clone this gene by traditional methods. An attempt to sequence the entire protein by Edman degradation was difficult from the outset because the GFP was refractory to most attempts at specific
5 proteolysis. Although over 80% of the protein was eventually accurately sequenced, a 30 amino acid region (110-139 of SEQ ID No. 27) had not be sequenced (as well as other regions, including amino acids 41-43, 65-71; SEQ ID No. 27). This 30 amino acid region apparently is degraded by the proteolytic methods used into very small fragments that are difficult to isolate and
10 sequence; proper ordering of sequenced fragments was also difficult.

The cloned DNA fragments can be replicated in bacterial cells, such as *E. coli*. A preferred DNA fragment also includes a bacterial origin of replication, to ensure the maintenance of the DNA fragment from generation to generation of the bacteria. In this way, large quantities of the DNA fragment can be produced
15 by replication in bacteria. Preferred bacterial origins of replication include, but are not limited to, the f1-ori and col E1 origins of replication. Preferred hosts contain chromosomal copies of DNA encoding T7 RNA polymerase operably linked to an inducible promoter, such as the lacUV promoter (see, U.S. Patent No. 4,952,496). Such hosts include, but are not limited to, lysogens *E. coli*
20 strains HMS174(DE3)pLysS, BL21(DE3)pLysS, HMS174(DE3) and BL21(DE3). Strain BL21(DE3) is preferred. The pLys strains provide low levels of T7 lysozyme, a natural inhibitor of T7 RNA polymerase.

For expression and for preparation of muteins, such as temperature sensitive muteins, eukaryotic cells, among them, yeast cells, such as
25 *Saccharomyces* are preferred.

Nucleic acid encoding fusion proteins of the luciferases and GFPs are also provided. The resulting fusion proteins are also provided. Nucleic acids that encode luciferase and GFPs as polycistronic mRNA or under the control of separate promoters are also provided. Methods of use thereof are also provided.
30

The GFP cloned from *Renilla* has spectral properties that make it extremely useful. These properties include very high quantum efficiency, high

-81-

molar absorbency and efficient use with universally available fluorescein filters (*e.g.*, Endo GFP filter set sold by Chroma). It is known that *Renilla reniformis* GFP is sixfold brighter than the wild-type *Aequorea* GFP on a molar basis, and three to fourfold brighter than the brightest mutant.

- 5 The *Renilla mullerei* GFP encoded by the nucleic acid clones provided herein exhibits similar functional characteristics, and the spectra appear identical with those from native *reniformis* GFP. Sequence comparison among the GFPs isolated from *Aequorea victoria*, *Renilla mullerei*, and *Ptilosarcus* reveal that the chromophore sequences of *R. mullerei* and *Ptilosarcus* are identical, and
10 differ from *A. victoria*. These sequence differences point to protein sites that can be modified without affecting the essential fluorescence properties and also provide a means to identify residues that change these properties.

4. Isolation and identification of DNA encoding *Renilla mulleri* GFP

- Methods for identification and cloning of GPFs from *Renilla* have been
15 described (see, published International PCT application No. WO 99/49019, and copending allowed U.S. application Serial No. 09/277,716). Nucleic acid encoding *Renilla mulleri* has been isolated. Briefly, a *R. mulleri* λ Uni-Zap cDNA expression plasmid library was prepared, transformed into competent *E. coli* cells and plated onto modified L-broth plates containing carbon black to absorb
20 background fluorescence. Transformants were sprayed with a solution containing IPTG to induce expression of the recombinant *Renilla* GFP from the heterologous cDNA. To identify GFP expressing clones, transformants were placed in blue light, preferably 470 to 490 nm light, and colonies that emitted green fluorescence were isolated and grown in pure culture.
25 The nucleotide sequence of the cDNA insert of a green fluorescent transformant was determined (*e.g.*, see SEQ ID No. 15). The 1,079 cDNA insert encodes a 238 amino acid polypeptide that is only 23.5 % identical to *A. victoria* GFP. The recombinant protein exhibits excitation and emission spectra similar to those reported for live *Renilla* species.

-82-

5. Isolation and identification of DNA encoding *Renilla mulleri* luciferase

The above-described *R. mulleri* cDNA expression library was also used to clone DNA encoding a *R. mulleri* luciferase. Single colony transformants were grown on modified L-broth plates containing carbon black and expression from the heterologous DNA was induced with IPTG, essentially as described above. After allowing time for expression, the transformants were sprayed with coelenterazine and screened for those colonies that emit blue light. Light-emitting colonies were isolated and grown in pure culture.

The nucleotide sequence of the cDNA insert contained in the light-emitting transformant was determined. The 1,217 cDNA insert encodes a 311 amino acid polypeptide. The recombinant protein exhibits excitation and emission spectra similar to those reported for live *Renilla* species.

E. RECOMBINANT EXPRESSION OF PROTEINS

1. DNA encoding *Renilla* proteins

As described above, DNA encoding a *Renilla* GFP or *Renilla* luciferase can be isolated from natural sources, synthesized based on *Renilla* sequences provided herein or isolated as described herein.

In preferred embodiments, the DNA fragment encoding a *Renilla* GFP has the sequence of amino acids set forth in SEQ ID No. 27, encoded by nucleic acid, such as that set forth SEQ ID Nos. 23-26 and 27.

A DNA molecule encoding a *Renilla* luciferase has the sequence of amino acids set forth in SEQ ID No. 18. In more preferred embodiments, the DNA fragment encodes the sequence of amino acids encoded by nucleotides 31-963 of the sequence of nucleotides set forth in SEQ ID No. 17.

2. DNA constructs for recombinant production of *Renilla reniformis* GFP and other proteins

DNA is introduced into a plasmid for expression in a desired host. In preferred embodiments, the host is a bacterial host. The sequences of nucleotides in the plasmids that are regulatory regions, such as promoters and operators, are operationally associated with one another for transcription of the sequence of nucleotides that encode a *Renilla* GFP or luciferase. The sequence

-83-

of nucleotides encoding the FGF mutein may also include DNA encoding a secretion signal, whereby the resulting peptide is a precursor of the *Renilla* GFP.

In preferred embodiments the DNA plasmids also include a transcription terminator sequence. The promoter regions and transcription terminators are
5 each independently selected from the same or different genes.

A wide variety of multipurpose vectors suitable for the expression of heterologous proteins are known to those of skill in the art and are commercially available. Expression vectors containing inducible promoters or constitutive promoters that are linked to regulatory regions are preferred. Such promoters
10 include, but are not limited to, the T7 phage promoter and other T7-like phage promoters, such as the T3, T5 and SP6 promoters, the *trp*, *lpp*, *tet* and *lac* promoters, such as the *lacUV5*, from *E. coli*; the SV40 promoter; the P10 or polyhedron gene promoter of baculovirus/insect cell expression systems, retroviral long-terminal repeats and inducible promoters from other eukaryotic
15 expression systems.

Particularly preferred vectors for recombinant expression of *Renilla mulleri* in prokaryotic organisms are *lac*- and T7 promoter-based vectors, such as the well known Bluescript vectors, which are commercially available (Stratagene, La Jolla, CA).

20 3. Host organisms for recombinant production of *Renilla* proteins

Host organisms include those organisms in which recombinant production of heterologous proteins have been carried out, such as, but not limited to, bacteria (for example, *E. coli*), yeast (for example, *Saccharomyces cerevisiae* and *Pichia pastoris*), fungi, baculovirus/insect systems, amphibian cells, mammalian
25 cells, plant cells and insect cells. Presently preferred host organisms are strains of bacteria or yeast. Most preferred host organisms are strains of *E. coli* or *Saccharomyces cerevisiae*.

4. Methods for recombinant production of *Renilla* proteins

The DNA encoding a *Renilla* GFP or *Renilla mulleri* luciferase is introduced
30 into a plasmid in operative linkage to an appropriate promoter for expression of polypeptides in a selected host organism. The DNA molecule encoding the *Renilla* GFP or luciferase may also include a protein secretion signal that

-84-

functions in the selected host to direct the mature polypeptide into the periplasm or culture medium. The resulting *Renilla* GFP or luciferase can be purified by methods routinely used in the art, including methods described hereinafter in the Examples.

5 Methods of transforming suitable host cells, preferably bacterial cells, and more preferably *E. coli* cells, as well as methods applicable for culturing said cells containing a gene encoding a heterologous protein, are generally known in the art. See, for example, Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.

10 Once the *Renilla*-encoding DNA molecule has been introduced into the host cell, the desired *Renilla* GFP is produced by subjecting the host cell to conditions under which the promoter is induced, whereby the operatively linked DNA is transcribed. The cellular extracts of lysed cells containing the protein
15 may be prepared and the resulting "clarified lysate" was employed as a source of recombinant *Renilla* GFP or *Renilla mulleri* luciferase. Alternatively, the lysate may be subjected to additional purification steps (e.g., ion exchange chromatography or immunoaffinity chromatography) to further enrich the lysate or provide a homogeneous source of the purified enzyme (see *e.g.*, U.S. Patent
20 Nos. 5,292,658 and 5,418,155).

5. Recombinant cells expressing heterologous nucleic acid encoding *Renilla* GFP

25 Cells, vectors and methods are described with respect to *Renilla*. The same cells, vectors and methods may be used for expressing luciferases and other GFPs from species including *Gaussia*, *Pleuromamma* and *Ptilosarcus*.

Recombinant cells containing heterologous nucleic acid encoding a *Renilla reniformis* GFP are provided. In preferred embodiments, the recombinant cells express the encoded *Renilla* GFP which is functional and non-toxic to the cell.

30 In certain embodiments, the recombinant cells may also include heterologous nucleic acid encoding a component of a bioluminescence-generating system, preferably a photoprotein or luciferase. In preferred embodiments, the nucleic acid encoding the bioluminescence-generating system

-85-

component is isolated from the species *Aequorea*, *Vargula* or *Renilla*. In more preferred embodiments, the bioluminescence-generating system component is a *Renilla mulleri* luciferase having the amino acid sequence set forth in SEQ ID No. 18.

- 5 Recombinant host cells containing heterologous nucleic acid encoding a *Renilla mulleri* luciferase are also provided. In preferred embodiments, the heterologous nucleic acid encodes the sequence of amino acids as set forth in SEQ ID No. 18. In more preferred embodiments, the heterologous nucleic acid encodes the sequence of nucleotides set forth in SEQ ID No. 17.
- 10 Exemplary cells include bacteria (e.g., *E. coli*), plant cells, cells of mammalian origin (e.g., COS cells, mouse L cells, Chinese hamster ovary (CHO) cells, human embryonic kidney (HEK) cells, African green monkey cells and other such cells known to those of skill in the art), amphibian cells (e.g., *Xenopus laevis* oocytes), yeast cells (e.g., *Saccharomyces cerevisiae*, *Pichia pastoris*), and
- 15 the like. Exemplary cells for expressing injected RNA transcripts include *Xenopus laevis* oocytes. Eukaryotic cells that are preferred for transfection of DNA are known to those of skill in the art or may be empirically identified, and include HEK293 (which are available from ATCC under accession #CRL 1573); Ltk⁻ cells (which are available from ATCC under accession #CCL1.3); COS-7
- 20 cells (which are available from ATCC under accession #CRL 1651); and DG44 cells (dhfr CHO cells; see, e.g., Urlaub et al. (1986) *Cell. Molec. Genet.* 12: 555). Presently preferred cells include strains of bacteria and yeast.

- The recombinant cells that contain the heterologous DNA encoding the *Renilla* GFP are produced by transfection with DNA encoding a *Renilla* GFP or
- 25 luciferase or by introduction of RNA transcripts of DNA encoding a *Renilla* proteins using methods well known to those of skill in the art. The DNA may be introduced as a linear DNA fragment or may be included in an expression vector for stable or transient expression of the encoding DNA. The sequences set forth herein for *Renilla reniformis* GFP are presently preferred (see SEQ ID Nos
- 30 23-25 and 27; see, also SEQ ID No. 26, which sets forth human optimized codons).

-86-

Heterologous DNA may be maintained in the cell as an episomal element or may be integrated into chromosomal DNA of the cell. The resulting recombinant cells may then be cultured or subcultured (or passaged, in the case of mammalian cells) from such a culture or a subculture thereof. Also, DNA may
5 be stably incorporated into cells or may be transiently expressed using methods known in the art.

The recombinant cells can be used in a wide variety of cell-based assay methods, such as those methods described for cells expressing wild type or modified *A. victoria* GFPs or GFP fusion proteins (*e.g.*, see U.S. Patent No.
10 5,625,048; International patent application Publication Nos. WO 95/21191; WO 96/23810; WO 96/27675; WO 97/26333; WO 97/28261; WO 97/41228; and WO 98/02571).

F. COMPOSITIONS AND CONJUGATES

Compositions and conjugates and methods of use are described with
15 reference to *Renilla* proteins and nucleic acids. The same compositions and methods for preparation and use thereof are intended for use with other luciferases, such as *Pleuromamma* and *Ptilosarcus* proteins and nucleic acids.

1. *Renilla* GFP compositions

Compositions containing a *Renilla* GFP or GFP peptide are provided. The
20 compositions can take any of a number of forms, depending on the intended method of use therefor. In certain embodiments, for example, the compositions contain a *Renilla* GFP or GFP peptide, preferably *Renilla mulleri* GFP or *Renilla reniformis* GFP peptide, formulated for use in luminescent novelty items, immunoassays, FRET and FET assays. The compositions may also be used in
25 conjunction with multi-well assay devices containing integrated photodetectors, such as those described herein.

Compositions that contain a *Renilla mulleri* GFP or GFP peptide and at least one component of a bioluminescence-generating system, preferably a luciferase, luciferin or a luciferase and a luciferin, are provided. In preferred
30 embodiments, the luciferase/luciferin bioluminescence- generating system is selected from those isolated from: an insect system, a coelenterate system, a ctenophore system, a bacterial system, a mollusk system, a crustacea system, a

-87-

fish system, an annelid system, and an earthworm system. Presently preferred bioluminescence- generating systems are those isolated from *Renilla*, *Aequorea*, and *Vargula*.

5 In more preferred embodiments, the bioluminescence-generating system component is a *Renilla mulleri* luciferase having the amino acid sequence set forth in SEQ ID No. 18 or a *Renilla reniformis* luciferase. These compositions can be used in a variety of methods and systems, such as included in conjunction with diagnostic systems for the *in vivo* detection of neoplastic tissues and other tissues, such as those methods described in detail below.

10 These methods and products include any known to those of skill in the art in which luciferase is used, including, but not limited to U.S. application Serial No. 08/757,046, 08/597,274 and 08/990,103, U.S. Patent No. 5,625,048; International patent application Publication Nos. WO 95/21191; WO 96/23810; WO 96/27675; WO 97/26333; WO 97/28261; WO 97/41228; and
15 WO 98/02571).

2. *Renilla* luciferase compositions

DNA encoding the *Renilla mulleri* luciferase or *Renilla reniformis* luciferase is used to produce the encoded luciferase, which has diagnostic applications as well as use as a component of the bioluminescence generating systems as
20 described herein, such as in beverages, and methods of diagnosis of neoplasia and in the diagnostic chips described herein. These methods and products include any known to those of skill in the art in which luciferase is used, including, but not limited to, U.S. application Serial No. 08/757,046, 08/597,274 and 08/990,103, U.S. Patent No. 5,625,048; International patent
25 application Publication Nos. WO 95/21191; WO 96/23810; WO 96/27675; WO 97/26333; WO 97/28261; WO 97/41228; and WO 98/02571).

In other embodiments, the *Renilla* luciferase and the remaining components may be packaged as separate compositions, that, upon mixing, glow. For example, a composition containing *Renilla* luciferase may be provided
30 separately from, and for use with, a separate composition containing a bioluminescence substrate and bioluminescence activator. In another instance, luciferase and luciferin compositions may be separately provided and the

-88-

bioluminescence activator may be added after, or simultaneously with, mixing of the other two compositions.

3. Conjugates

Conjugates are provided herein for a variety of uses. Among them are for
5 targeting to tumors for visualization of the tumors, particularly *in situ* during surgery. A general description of these conjugates and the uses thereof is described in allowed U.S. application Serial No. 08/908,909. In practice, prior to a surgical procedure, the conjugate is administered via any suitable route, whereby the targeting agent binds to the targeted tissue by virtue of its specific
10 interaction with a tissue-specific cell surface protein. During surgery the tissue is contacted, with the remaining component(s), typically by spraying the area or local injection, and any tissue to which conjugate is bound will glow. The glow should be sufficient to see under dim light or, if necessary, in the dark.

The conjugates that are provided herein contain a targeting agent, such
15 as a tissue specific or tumor specific monoclonal antibody or fragment thereof linked either directly or via a linker to a targeted agent, a *Renilla* GFP, *Renilla* or *Gaussia* luciferase and other luciferases (including photoproteins or luciferase enzymes) or a luciferin. The targeted agent may be coupled to a microcarrier. The linking is effected either chemically, by recombinant expression of a fusion
20 protein in instances when the targeted agent is a protein, and by combinations of chemical and recombinant expression. The targeting agent is one that will preferentially bind to a selected tissue or cell type, such as a tumor cell surface antigen or other tissue specific antigen.

Methods for preparing conjugates are known to those of skill in the art.
25 For example, aequorin that is designed for conjugation and conjugates containing such aequorin have been produced (see, *e.g.*, International PCT application No. WO 94/18342; see, also Smith *et al.* (1995) in *American Biotechnology Laboratory*). Aequorin has been conjugated to an antibody molecule by means of a sulfhydryl-reacting binding agent (Stultz *et al.* (1992) Use of Recombinant
30 Biotinylated Apoaequorin from *Escherichia coli*. *Biochemistry* 31, 1433-1442). Such methods may be adapted for use herein to produce the luciferase coupled to protein or other such molecules, which are useful as targeting agents.

-89-

Vargula luciferase has also been linked to other molecules (see, *e.g.*, Japanese application No. JP 5064583, March 19, 1993). Such methods may be adapted for use herein to produce luciferase coupled to molecules that are useful as targeting agents.

- 5 The conjugates can be employed to detect the presence of or quantitate a particular antigen in a biological sample by direct correlation to the light emitted from the bioluminescent reaction.

- As an alternative, a component of the bioluminescence generating system may be modified for linkage, such as by addition of amino acid residues that are
10 particularly suitable for linkage to the selected substrate. This can be readily effected by modifying the DNA and expressing such modified DNA to produce luciferase with additional residues at the N- or C-terminus.

- Methods for preparing conjugates are known to those of skill in the art. For example, aequorin that is designed for conjugation and conjugates containing
15 such aequorin have been produced (see, *e.g.*, International PCT application No. WO 94/18342; see, also Smith *et al.* (1995) in *American Biotechnology Laboratory*). Aequorin has been conjugated to an antibody molecule by means of a sulfhydryl-reacting binding agent (Stultz *et al.* (1992) Use of Recombinant Biotinylated Apoequorin from *Escherichia coli*. *Biochemistry* 31, 1433-1442).
20 Such methods may be adapted for use herein to produce aequorin coupled to protein or other such molecules, which are useful as targeting agents.

- Vargula* luciferase has also been linked to other molecules (see, *e.g.*, Japanese application No. JP 5064583, March 19, 1993). Such methods may be adapted for use herein to produce aequorin coupled to protein or other such molecules,
25 which are useful as targeting agents. The bioluminescence generating reactions are used with the *Renilla reniformis* GFP provided herein.

a. Linkers

- Any linker known to those of skill in the art may be used herein. Other linkers are suitable for incorporation into chemically produced conjugates.
30 Linkers that are suitable for chemically linked conjugates include disulfide bonds, thioether bonds, hindered disulfide bonds, and covalent bonds between free reactive groups, such as amine and thiol groups. These bonds are produced

-90-

using heterobifunctional reagents to produce reactive thiol groups on one or both of the polypeptides and then reacting the thiol groups on one polypeptide with reactive thiol groups or amine groups to which reactive maleimido groups or thiol groups can be attached on the other. Other linkers include, acid cleavable
5 linkers, such as bismaleimideoxy propane, acid labile-transferrin conjugates and adipic acid dihydrazide, that would be cleaved in more acidic intracellular compartments; cross linkers that are cleaved upon exposure to UV or visible light and linkers, such as the various domains, such as C_H1, C_H2, and C_H3, from the constant region of human IgG₁ (see, Batra *et al.* (1993) *Molecular Immunol.*
10 30:379-386). In some embodiments, several linkers may be included in order to take advantage of desired properties of each linker.

Chemical linkers and peptide linkers may be inserted by covalently coupling the linker to the TA and the targeted agent. The heterobifunctional agents, described below, may be used to effect such covalent coupling. Peptide
15 linkers may also be linked by expressing DNA encoding the linker and TA, linker and targeted agent, or linker, targeted agent and TA as a fusion protein.

Flexible linkers and linkers that increase solubility of the conjugates are contemplated for use, either alone or with other linkers are contemplated herein.

Numerous heterobifunctional cross-linking reagents that are used to form
20 covalent bonds between amino groups and thiol groups and to introduce thiol groups into proteins, are known to those of skill in this art (see, *e.g.*, the PIERCE CATALOG, ImmunoTechnology Catalog & Handbook, 1992-1993, which describes the preparation of and use of such reagents and provides a commercial source for such reagents; see, also, *e.g.*, Cumber *et al.* (1992) *Bioconjugate*
25 *Chem.* 3:397-401; Thorpe *et al.* (1987) *Cancer Res.* 47:5924-5931; Gordon *et al.* (1987) *Proc. Natl. Acad. Sci.* 84:308-312; Walden *et al.* (1986) *J. Mol. Cell Immunol.* 2:191-197; Carlsson *et al.* (1978) *Biochem. J.* 173: 723-737; Mahan *et al.* (1987) *Anal. Biochem.* 162:163-170; Wawryznaczak *et al.* (1992) *Br. J. Cancer* 66:361-366; Fattom *et al.* (1992) *Infection & Immun.* 60:584-589).
30 These reagents may be used to form covalent bonds between the TA and targeted agent. These reagents include, but are not limited to: N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP; disulfide linker); sulfosuccinimidyl 6-(3-(2-

-91-

pyridyldithio)propionamido)hexanoate (sulfo-LC-SPDP); succinimidylloxycarbonyl- α -methyl benzyl thiosulfate (SMBT, hindered disulfate linker); succinimidyl 6-(3-(2-pyridyldithio) propionamido)hexanoate (LC-SPDP); sulfosuccinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate (sulfo-SMCC); succinimidyl 3-(2-pyridyldithio)butyrate (SPDB; hindered disulfide bond linker); sulfosuccinimidyl 2-(7-azido-4-methylcoumarin-3-acetamide) ethyl-1,3'-dithiopropionate (SAED); sulfo-succinimidyl 7-azido-4-methylcoumarin-3-acetate (SAMCA); sulfosuccinimidyl 6-(α -methyl- α -(2-pyridyldithio)toluamido)hexanoate (sulfo-LC-SMPT); 1,4-di-(3'-(2'-pyridyldithio)propionamido)butane (DPDPB); 4-succinimidylloxycarbonyl- α -methyl- α -(2-pyridylthio)toluene (SMPT, hindered disulfate linker); sulfo-succinimidyl 6-(α -methyl- α -(2-pyridyldithio)toluamido)hexanoate (sulfo-LC-SMPT); *m*-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS); *m*-maleimidobenzoyl-N-hydroxysulfosuccinimide ester (sulfo-MBS); N-succinimidyl(4-iodoacetyl)aminobenzoate (SIAB; thioether linker); sulfosuccinimidyl(4-iodoacetyl)amino benzoate (sulfo-SIAB); succinimidyl 4-(*p*-maleimidophenyl)butyrate (SMPB); sulfosuccinimidyl 4-(*p*-maleimidophenyl)butyrate (sulfo-SMPB); azidobenzoyl hydrazide (ABH).

Acid cleavable linkers, photocleavable and heat sensitive linkers may also be used, particularly where it may be necessary to cleave the targeted agent to permit it to be more readily accessible to reaction.

Acid cleavable linkers include, but are not limited to, bismaleimideoxypropane; and adipic acid dihydrazide linkers (see, *e.g.*, Fattom *et al.* (1992) *Infection & Immun.* 60:584-589) and acid labile transferrin conjugates that contain a sufficient portion of transferrin to permit entry into the intracellular transferrin cycling pathway (see, *e.g.*, Welhöner *et al.* (1991) *J. Biol. Chem.* 266:4309-4314).

Photocleavable linkers are linkers that are cleaved upon exposure to light (see, *e.g.*, Goldmacher *et al.* (1992) *Bioconj. Chem.* 3:104-107, which linkers are herein incorporated by reference), thereby releasing the targeted agent upon exposure to light. Photocleavable linkers that are cleaved upon exposure to light are known (see, *e.g.*, Hazum *et al.* (1981) in *Pept., Proc. Eur. Pept. Symp.*, 16th, Brunfeldt, K (Ed), pp. 105-110, which describes the use of a

-92-

- nitrobenzyl group as a photocleavable protective group for cysteine; Yen *et al.* (1989) *Makromol. Chem* 190:69-82, which describes water soluble photocleavable copolymers, including hydroxypropylmethacrylamide copolymer, glycine copolymer, fluorescein copolymer and methylrhodamine copolymer;
- 5 Goldmacher *et al.* (1992) *Bioconj. Chem.* 3:104-107, which describes a cross-linker and reagent that undergoes photolytic degradation upon exposure to near UV light (350 nm); and Senter *et al.* (1985) *Photochem. Photobiol* 42:231-237, which describes nitrobenzyloxycarbonyl chloride cross linking reagents that produce photocleavable linkages), thereby releasing the targeted agent upon
- 10 exposure to light. Such linkers would have particular use in treating dermatological or ophthalmic conditions that can be exposed to light using fiber optics. After administration of the conjugate, the eye or skin or other body part can be exposed to light, resulting in release of the targeted moiety from the conjugate. Such photocleavable linkers are useful in connection with diagnostic
- 15 protocols in which it is desirable to remove the targeting agent to permit rapid clearance from the body of the animal.

b. Targeting Agents

- Targeting agents include any agent that will interact with and localize the targeted agent cells in a tumor or specialized tissue (targeted tissue). Such
- 20 agents include any agent that specifically interacts with a cell surface protein or receptor that is present at sufficiently higher concentrations or amounts on the targeted tissue, whereby, when contacted with an appropriate bioluminescence generating reagent and activators produces light. These agents include, but are not limited to, growth factors, preferentially modified to not internalize,
- 25 methotrexate, and antibodies, particularly, antibodies raised against tumor specific antigens. A plethora of tumor-specific antigens have been identified from a number of human neoplasms.

c. Anti-tumor Antigen Antibodies

- Polyclonal and monoclonal antibodies may be produced against selected
- 30 antigens. Alternatively, many such antibodies are presently available. An exemplary list of antibodies and the tumor antigen for which each has been directed against is provided in U.S. application Serial No.08/908,909, which is incorporated

-93-

by reference in its entirety. It is contemplated that any of the antibodies listed may be conjugated with a bioluminescence generating component following the methods provided herein.

Among the preferred antibodies for use in the methods herein are those of human origin or, more preferably, are humanized monoclonal antibodies. These are preferred for diagnosis of humans.

d. Preparation of the conjugates

The methods for preparation of the conjugates for use in the tumor diagnostic methods can be used for preparation of the fusion proteins and conjugated proteins for use in the BRET system described below. Any method for linking proteins may be used. For example, methods for linking a luciferase to an antibody is described in U.S. Patent No. 5,486,455. As noted above, the targeting agent and luciferin or luciferase may be linked directly, such as through covalent bonds, *i.e.*, sulfhydryl bonds or other suitable bonds, or they may be linked through a linker. There may be more than one luciferase or luciferin per targeting agent, or more than one targeting agent per luciferase or luciferin.

Alternatively, an antibody, or F(Ab)₂ antigen-binding fragment thereof or other protein targeting agent may be fused (directly or via a linking peptide) to the luciferase using recombinant DNA technology. For example, the DNA encoding any of the anti-tumor antibodies of Table 3 may be ligated in the same translational reading frame to DNA encoding any of the above-described luciferases, *e.g.*, SEQ ID NOs. 1-14 and inserted into an expression vector. The DNA encoding the recombinant antibody-luciferase fusion may be introduced into an appropriate host, such as bacteria or yeast, for expression.

4. Formulation of the compositions for use in the diagnostic systems

In most embodiments, the *Renilla* GFPS and components of the diagnostic systems provided herein, such as *Renilla* luciferase, are formulated into two compositions: a first composition containing the conjugate; and a second composition containing the remaining components of the bioluminescence generating system. The compositions are formulated in any manner suitable for administration to an animal, particularly a mammal, and more particularly a human. Such formulations include those suitable for topical, local, enteric,

-94-

parenteral, intracystal, intracutaneous, intravitreal, subcutaneous, intramuscular, or intravenous administration.

For example, the conjugates, which in preferred embodiments, are a targeting agent linked to a luciferase (or photoprotein) are formulated for systemic or local administration. The remaining components are formulated in a separate second composition for topical or local application. The second composition will typically contain any other agents, such as spectral shifters that will be included in the reaction. It is preferred that the components of the second composition are formulated in a time release manner or in some other manner that prevents degradation and/or interaction with blood components.

a. The first composition: formulation of the conjugates

As noted above, the conjugates either contain a luciferase or luciferin and a targeting agents. The preferred conjugates are formed between a targeting agent and a luciferase, particularly the *Gaussia*, *Renilla mulleri* or *Pleuromamma* luciferase. The conjugates may be formulated into pharmaceutical compositions suitable for topical, local, intravenous and systemic application. Effective concentrations of one or more of the conjugates are mixed with a suitable pharmaceutical carrier or vehicle. The concentrations or amounts of the conjugates that are effective requires delivery of an amount, upon administration, that results in a sufficient amount of targeted moiety linked to the targeted cells or tissue whereby the cells or tissue can be visualized during the surgical procedure. Typically, the compositions are formulated for single dosage administration. Effective concentrations and amounts may be determined empirically by testing the conjugates in known *in vitro* and *in vivo* systems, such as those described here; dosages for humans or other animals may then be extrapolated therefrom.

Upon mixing or addition of the conjugate(s) with the vehicle, the resulting mixture may be a solution, suspension, emulsion or the like. The form of the resulting mixture depends upon a number of factors, including the intended mode of administration and the solubility of the conjugate in the selected carrier or vehicle. The effective concentration is sufficient for targeting a sufficient

-95-

amount of targeted agent to the site of interest, whereby when combined with the remaining reagents during a surgical procedure the site will glow. Such concentration or amount may be determined based upon *in vitro* and/or *in vivo* data, such as the data from the mouse xenograft model for tumors or rabbit
5 ophthalmic model. If necessary, pharmaceutically acceptable salts or other derivatives of the conjugates may be prepared.

Pharmaceutical carriers or vehicles suitable for administration of the conjugates provided herein include any such carriers known to those skilled in the art to be suitable for the particular mode of administration.

10 In addition, the conjugates may be formulated as the sole pharmaceutically ingredient in the composition or may be combined with other active ingredients.

The conjugates can be administered by any appropriate route, for example, orally, parenterally, intravenously, intradermally, subcutaneously, or topically, in liquid, semi-liquid or solid form and are formulated in a manner
15 suitable for each route of administration. Intravenous or local administration is presently preferred. Tumors and vascular proliferative disorders, will typically be visualized by systemic, intradermal or intramuscular, modes of administration.

The conjugate is included in the pharmaceutically acceptable carrier in an amount sufficient to produce detectable tissue and to not result in undesirable
20 side effects on the patient or animal. It is understood that number and degree of side effects depends upon the condition for which the conjugates are administered. For example, certain toxic and undesirable side effects are tolerated when trying to diagnose life-threatening illnesses, such as tumors, that would not be tolerated when diagnosing disorders of lesser consequence.

25 The concentration of conjugate in the composition will depend on absorption, inactivation and excretion rates thereof, the dosage schedule, and amount administered as well as other factors known to those of skill in the art. Typically an effective dosage should produce a serum concentration of active ingredient of from about 0.1 ng/ml to about 50-1000 μ g/ml, preferably 50-100
30 μ g/ml. The pharmaceutical compositions typically should provide a dosage of from about 0.01 mg to about 100 - 2000 mg of conjugate, depending upon the conjugate selected, per kilogram of body weight per day. Typically, for

-96-

intravenous administration a dosage of about between 0.05 and 1 mg/kg should be sufficient. Local application for, such as visualization of ophthalmic tissues or local injection into joints, should provide about 1 ng up to 1000 μ g, preferably about 1 μ g to about 100 μ g, per single dosage administration. It is understood
5 that the amount to administer will be a function of the conjugate selected, the indication, and possibly the side effects that will be tolerated. Dosages can be empirically determined using recognized models.

The active ingredient may be administered at once, or may be divided into a number of smaller doses to be administered at intervals of time. It is
10 understood that the precise dosage and duration of administration is a function of the disease condition being diagnosed and may be determined empirically using known testing protocols or by extrapolation from *in vivo* or *in vitro* test data. It is to be noted that concentrations and dosage values may also vary with the severity of the condition to be alleviated. It is to be further understood that
15 for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed compositions.

20 Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application can include any of the following components: a sterile diluent, such as water for injection, saline solution, fixed oil, polyethylene glycol, glycerine, propylene glycol or other synthetic solvent; antimicrobial agents, such as benzyl alcohol and methyl parabens; antioxidants, such as ascorbic acid and
25 sodium bisulfite; chelating agents, such as ethylenediaminetetraacetic acid (EDTA); buffers, such as acetates, citrates and phosphates; and agents for the adjustment of tonicity such as sodium chloride or dextrose. Parental preparations can be enclosed in ampules, disposable syringes or multiple dose vials made of glass, plastic or other suitable material.

30 If administered intravenously, suitable carriers include physiological saline or phosphate buffered saline (PBS), and solutions containing thickening and solubilizing agents, such as glucose, polyethylene glycol, and polypropylene

-97-

glycol and mixtures thereof. Liposomal suspensions may also be suitable as pharmaceutically acceptable carriers. These may be prepared according to methods known to those skilled in the art.

- The conjugates may be prepared with carriers that protect them against rapid elimination from the body, such as time release formulations or coatings. Such carriers include controlled release formulations, such as, but not limited to, implants and microencapsulated delivery systems, and biodegradable, biocompatible polymers, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, polyorthoesters, polylactic acid and others.
- 10 The conjugates may be formulated for local or topical application, such as for topical application to the skin and mucous membranes, such as in the eye, in the form of gels, creams, and lotions and for application to the eye or for intracisternal or intraspinal application. Such solutions, particularly those intended for ophthalmic use, may be formulated as 0.01% -10% isotonic
- 15 solutions, pH about 5-7, with appropriate salts. The ophthalmic compositions may also include additional components, such as hyaluronic acid. The conjugates may be formulated as aerosols for topical application (see, *e.g.*, U.S. Patent Nos. 4,044,126, 4,414,209, and 4,364,923).

- Also, the compositions for activation of the conjugate *in vivo* during
- 20 surgical procedures may be formulated as an aerosol. These compositions contain the activators and also the remaining bioluminescence generating agent, such as luciferin, where the conjugate targets a luciferase, or a luciferase, where the conjugate targets a luciferin, such as coelenterazine.

- If oral administration is desired, the conjugate should be provided in a
- 25 composition that protects it from the acidic environment of the stomach. For example, the composition can be formulated in an enteric coating that maintains its integrity in the stomach and releases the active compound in the intestine. Oral compositions will generally include an inert diluent or an edible carrier and may be compressed into tablets or enclosed in gelatin capsules. For the purpose
- 30 of oral administration, the active compound or compounds can be incorporated with excipients and used in the form of tablets, capsules or troches.

-98-

Pharmaceutically compatible binding agents and adjuvant materials can be included as part of the composition.

Tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder, such as
5 microcrystalline cellulose, gum tragacanth and gelatin; an excipient such as starch and lactose, a disintegrating agent such as, but not limited to, alginic acid and corn starch; a lubricant such as, but not limited to, magnesium stearate; a glidant, such as, but not limited to, colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; and a flavoring agent such as peppermint, methyl
10 salicylate, and fruit flavoring.

When the dosage unit form is a capsule, it can contain, in addition to material of the above type, a liquid carrier such as a fatty oil. In addition, dosage unit forms can contain various other materials which modify the physical form of the dosage unit, for example, coatings of sugar and other enteric agents.
15 The conjugates can also be administered as a component of an elixir, suspension, syrup, wafer, chewing gum or the like. A syrup may contain, in addition to the active compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings and flavors.

The active materials can also be mixed with other active materials that do
20 not impair the desired action, or with materials that supplement the desired action, such as cis-platin for treatment of tumors.

Finally, the compounds may be packaged as articles of manufacture containing packaging material, one or more conjugates or compositions as provided herein within the packaging material, and a label that indicates the
25 indication for which the conjugate is provided.

b. The second composition

The second composition will include the remaining components of the bioluminescence generating reaction. In preferred embodiments in which these components are administered systemically, the remaining components include
30 the luciferin or substrate, and optionally additional agents, such as spectral shifters, particularly the GFPs provided herein. These components, such as the luciferin, can be formulated as described above for the conjugates. In some

-99-

embodiments, the luciferin or luciferase in this composition will be linked to a protein carrier or other carrier to prevent degradation or dissolution into blood cells or other cellular components.

For embodiments, in which the second composition is applied locally or topically, they can be formulated in a spray or aerosol or other suitable means for local or topical application.

In certain embodiments described herein, all components, except an activator are formulated together, such as by encapsulation in a time release formulation that is targeted to the tissue. Upon release the composition will have been localized to the desired site, and will begin to glow.

In practice, the two compositions can be administered simultaneously or sequentially. Typically, the first composition, which contains the conjugate is administered first, generally an hour or two before the surgery, and the second composition is then administered, either pre-operatively or during surgery.

The conjugates that are provided herein contain a targeting agent, such as a tissue specific or tumor specific monoclonal antibody or fragment thereof linked either directly or via a linker to a targeted agent, a luciferase (including photoproteins or luciferase enzymes) or a luciferin. The targeted agent may be coupled to a microcarrier. The linking is effected either chemically, by recombinant expression of a fusion protein in instances when the targeted agent is a protein, and by combinations of chemical and recombinant expression. The targeting agent is one that will preferentially bind to a selected tissue or cell type, such as a tumor cell surface antigen or other tissue specific antigen.

Methods for preparing conjugates are known to those of skill in the art. For example, aequorin that is designed for conjugation and conjugates containing such aequorin have been produced (see, *e.g.*, International PCT application No. WO 94/18342; see, also Smith *et al.* (1995) in *American Biotechnology Laboratory*). Aequorin has been conjugated to an antibody molecule by means of a sulfhydryl-reacting binding agent (Stultz *et al.* (1992) Use of Recombinant Biotinylated Apoaequorin from *Escherichia coli* (*Biochemistry* 31:1433-1442)). Such methods may be adapted for use herein to produce aequorin coupled to protein or other such molecules, which are useful as targeting agents.

-100-

Vargula luciferase has also been linked to other molecules (see, *e.g.*, Japanese application No. JP 5064583, March 19, 1993). Such methods may be adapted for use herein to produce aequorin coupled to protein or other such molecules, which are useful as targeting agents.

5 Aequorin-antibody conjugates have been employed to detect the presence of or quantitate a particular antigen in a biological sample by direct correlation to the light emitted from the bioluminescent reaction.

 As an alternative, the *Renilla* GFP or *Renilla mulleri* or *Gaussia* luciferase or a component of the bioluminescence generating system may be modified for
10 linkage, such as by addition of amino acid residues that are particularly suitable for linkage to the selected substrate. This can be readily effected by modifying the DNA and expressing such modified DNA to produce luciferase with additional residues at the N- or C-terminus.

 Selection of the system depends upon factors such as the desired color
15 and duration of the bioluminescence desired as well as the particular item. Selection of the targeting agent primarily depends upon the type and characteristics of neoplasia or tissue to be visualized and the setting in which visualization will be performed.

 The *Renilla reniformis* GFP is added to one or both compositions to act as
20 a spectral shifter.

c. Practice of the reactions in combination with targeting agents

 The particular manner in which each bioluminescence system will be combined with a selected targeting agent will be a function of the agent and the
25 neoplasia or tissue to be visualized. In general, however, a luciferin, *Renilla* GFP, *Renilla mulleri*, *Pleuromamma* or *Gaussia* luciferase or other luciferase, of the reaction will be conjugated to the targeting agent, administered to an animal prior to surgery. During the surgery, the tissues of interest are contacted with the remaining component(s) of a bioluminescence generating system. Any tissue
30 to which or with which the targeting agent reacts will glow.

-101-

Any color of visible light produced by a bioluminescence generating system is contemplated for use in the methods herein. Preferably the visible light is a combination of blue, green and/or red light of varying intensities and wavelengths. For visualizing neoplasia or specialty tissues through mammalian
5 tissues or tumors deeply embedded in tissue, longer wavelengths of visible light, *i.e.*, red and near infrared light, is preferred because wavelengths of near infrared light of about 700-1300 nm are known to penetrate soft tissue and bone (*e.g.*, see U.S. Patent No. 4,281,645).

In other embodiments, the conjugate can be applied to the tissues during
10 surgery, such as by spraying a sterile solution over the tissues, followed by application of the remaining components. Tissues that express the targeted antigen will glow.

The reagents may be provided in compositions, such as suspensions, as powders, as pastes or any in other suitable sterile form. They may be provided
15 as sprays, aerosols, or in any suitable form. The reagents may be linked to a matrix, particularly microbeads suitable for *in vivo* use and of size that they pass through capillaries. Typically all but one or more, though preferably all but one, of the components necessary for the reaction will be mixed and provided together; reaction will be triggered contacting the mixed component(s) with the
20 remaining component(s), such as by adding Ca^{2+} , FMN with reductase, FMNH₂, ATP, air or oxygen.

In preferred embodiments the luciferase or luciferase/luciferin will be provided in combination with the targeting agent before administration to the patient. The targeting agent conjugate will then be contacted *in vivo* with the
25 remaining components. As will become apparent herein, there are a multitude of ways in which each system may be combined with a selected targeting agent.

G. COMBINATIONS

Renilla reniformis GFP can be used in combination with articles of manufacture to produce novelty items. The *Renilla reniformis* GFP can be used
30 with a bioluminescence generating system. Such items and methods for preparation are described in detail in U.S. Patent Nos. 5,876,995, 6,152,358 and 6,113,886) These novelty items, which are articles of manufacture, are

-102-

designed for entertainment, recreation and amusement, and include, but are not limited to: toys, particularly squirt guns, toy cigarettes, toy "Halloween" eggs, footbags and board/card games; finger paints and other paints, slimy play material; textiles, particularly clothing, such as shirts, hats and sports gear suits, threads and yarns; bubbles in bubble making toys and other toys that produce bubbles; balloons; figurines; personal items, such as bath powders, body lotions, gels, powders and creams, nail polishes, cosmetics including make-up, toothpastes and other dentifrices, soaps, body paints, and bubble bath; items such as fishing lures, inks, paper; foods, such as gelatins, icings and frostings; fish food containing luciferins and transgenic fish, particularly transgenic fish that express a luciferase; plant food containing a luciferin or luciferase, preferably a luciferin for use with transgenic plants that express luciferase; and beverages, such as beer, wine, champagne, soft drinks, and ice cubes and ice in other configurations; fountains, including liquid "fireworks" and other such jets or sprays or aerosols of compositions that are solutions, mixtures, suspensions, powders, pastes, particles or other suitable form.

Any article of manufacture that can be combined with a bioluminescence-generating system as provided herein and thereby provide entertainment, recreation and/or amusement, including use of the items for recreation or to attract attention, such as for advertising goods and/or services that are associated with a logo or trademark is contemplated herein. Such uses may be in addition to or in conjunction with or in place of the ordinary or normal use of such items. As a result of the combination, the items glow or produce, such as in the case of squirt guns and fountains, a glowing fluid or spray of liquid or particles. H. Exemplary uses of *Renilla reniformis* GFPs and encoding nucleic acid molecules

1. Methods for diagnosis of neoplasms and other tissues

Methods for diagnosis and visualization of tissues *in vivo* or *in situ*, preferably neoplastic tissue, using compositions containing a *Renilla mulleri* or *Ptilosarcus* GFP and/or a *Renilla mulleri*, *Pleuromamma* or *Gaussia* luciferase are provided. For example, the *Renilla mulleri* GFP protein can be used in conjunction with diagnostic systems that rely on bioluminescence for visualizing

-103-

tissues *in situ*, such as those described in co-pending application Serial No. 08/908,909. The systems are particularly useful for visualizing and detecting neoplastic tissue and specialty tissue, such as during non-invasive and invasive procedures. The systems include compositions containing conjugates that

5 include a tissue specific, particularly a tumor-specific, targeting agent linked to a targeted agent, such as a *Renilla reniformis* GFP, a luciferase or luciferin. The systems also include a second composition that contains the remaining components of a bioluminescence generating reaction and/or the GFP. In some

10 embodiments, all components, except for activators, which are provided *in situ* or are present in the body or tissue, are included in a single composition.

In particular, the diagnostic systems include two compositions. A first composition that contains conjugates that, in preferred embodiments, include antibodies directed against tumor antigens conjugated to a component of the bioluminescence generating reaction, a luciferase or luciferin, preferably a

15 luciferase are provided. In certain embodiments, conjugates containing tumor-specific targeting agents are linked to luciferases or luciferins. In other embodiments, tumor-specific targeting agents are linked to microcarriers that are coupled with, preferably more than one of the bioluminescence generating components, preferably more than one luciferase molecule.

20 The second composition contains the remaining components of a bioluminescence generating system, typically the luciferin or luciferase substrate. In some embodiments, these components, particularly the luciferin are linked to a protein, such as a serum albumin, or other protein carrier. The carrier and time release formulations, permit systemically administered components to travel to

25 the targeted tissue without interaction with blood cell components, such as hemoglobin that deactivates the luciferin or luciferase.

2. Methods of diagnosing diseases

Methods for diagnosing diseases, particularly infectious diseases, using chip methodology, a luciferase/luciferin bioluminescence-generating system,

30 including a *Gaussia*, *Pleuromamma* or *Renilla mulleri* luciferase plus a *Renilla reniformis* GFP, are provided. In particular, the chip includes an integrated photodetector that detects the photons emitted by the bioluminescence-

-104-

generating system as shifted by the GFP. This chip device, which is described in copending U.S. application Serial No. 08/990,103, which is published as International PCT application No. WO 98/26277, includes an integrated photodetector that detects the photons emitted by the bioluminescence

5 generating system. The method may be practiced with any suitable chip device, including self-addressable and non-self addressable formats, that is modified as described herein for detection of generated photons by the bioluminescence generating systems. The chip device provided herein is adaptable for use in an array format for the detection and identification of infectious agents in biological

10 specimens.

To prepare the chip, a suitable matrix for chip production is selected, the chip is fabricated by suitably derivatizing the matrix for linkage of macromolecules, and including linkage of photodiodes, photomultipliers CCD (charge coupled device) or other suitable detector, for measuring light

15 production; attaching an appropriate macromolecule, such as a biological molecule or anti-ligand, *e.g.*, a receptor, such as an antibody, to the chip, preferably to an assigned location thereon. Photodiodes are presently among the preferred detectors, and specified herein. It is understood, however, that other suitable detectors may be substituted therefor.

20 In one embodiment, the chip is made using an integrated circuit with an array, such as an X-Y array, of photodetectors, such as that described in copending U.S. application Serial No. 08/990,103. The surface of circuit is treated to render it inert to conditions of the diagnostic assays for which the chip is intended, and is adapted, such as by derivatization for linking molecules, such as

25 antibodies. A selected antibody or panel of antibodies, such as an antibody specific for particularly bacterial antigen, is affixed to the surface of the chip above each photodetector. After contacting the chip with a test sample, the chip is contacted with a second antibody linked to the GFP, such as the *Renilla* GFP, to form a chimeric antibody-GFP fusion protein or an antibody linked to a

30 component of a bioluminescence generating system, such as a *Pleuromamma*, *Gaussia* or *R. mulleri* luciferase. The antibody is specific for the antigen. The remaining components of the bioluminescence generating reaction are added,

-105-

and, if any of the antibodies linked to a component of a bioluminescence generating system are present on the chip, light will be generated and detected by the adjacent photodetector. The photodetector is operatively linked to a computer, which is programmed with information identifying the linked
5 antibodies, records the event, and thereby identifies antigens present in the test sample.

3. Methods for generating *Renilla mulleri* luciferase, *Pleuromamma* luciferase and *Gaussia* luciferase fusion proteins with *Renilla reniformis* GFP.

10 Methods for generating GFP and luciferase fusion proteins are provided. The methods include linking DNA encoding a gene of interest, or portion thereof, to DNA encoding a *Renilla reniformis* GFP and a luciferase in the same translational reading frame. The encoded-protein of interest may be linked in-frame to the amino- or carboxyl-terminus of the GFP or luciferase. The DNA
15 encoding the chimeric protein is then linked in operable association with a promoter element of a suitable expression vector. Alternatively, the promoter element can be obtained directly from the targeted gene of interest and the promoter-containing fragment linked upstream from the GFP or luciferase coding sequence to produce chimeric GFP proteins.

20 For example, a chimeric fusion containing the luciferase, preferably a *Renilla* luciferase, more preferably a *Renilla reniformis* luciferase, and *Renilla reniformis* GFP encoding DNA linked to the N-terminal portion of a cellulose binding domain is provided.

4. Cell-based assays for identifying compounds

25 Methods for identifying compounds using recombinant cells that express heterologous DNA encoding a *Renilla reniformis* GFP under the control of a promoter element of a gene of interest are provided. The recombinant cells can be used to identify compounds or ligands that modulate the level of transcription from the promoter of interest by measuring GFP-mediated fluorescence.
30 Recombinant cells expressing chimeric GFPs may also be used for monitoring gene expression or protein trafficking, or determining the cellular localization of

-106-

the target protein by identifying localized regions of GFP-mediated fluorescence within the recombinant cell.

I. KITS

Kits may be prepared containing the *Renilla reniformis* GFP or the
5 encoding nucleic acid molecules (see, SEQ ID Nos. 23-26) with or without components of a bioluminescence generating system for use in diagnostic and immunoassay methods and with the novelty items, including those described herein.

In one embodiment, the kits contain appropriate reagents and an article of
10 manufacture for generating bioluminescence in combination with the article. These kits, for example, can be used with a bubble-blowing or producing toy or with a squirt gun. These kits can also include a reloading or charging cartridge.

In another embodiment, the kits are used for detecting and visualizing neoplastic tissue and other tissues and include a first composition that contains
15 the *Renilla reniformis* GFP and a selected luciferase, such as a *Renilla mulleri*, *Renilla reniformis* or *Gaussia* luciferase, and a second that contains the activating composition, which contains the remaining components of the bioluminescence generating system and any necessary activating agents.

In other embodiments, the kits are used for detecting and identifying
20 diseases, particularly infectious diseases, using multi-well assay devices and include a multi-well assay device containing a plurality of wells, each having an integrated photodetector, to which an antibody or panel of antibodies specific for one or more infectious agents are attached, and composition containing a secondary antibody, such as an antibody specific for the infectious agent that is
25 linked, for example, to a *Renilla reniformis* GFP protein, a chimeric antibody-*Renilla reniformis* GFP fusion protein, F(Ab)₂ antibody fragment-*Renilla reniformis* GFP fusion protein or to such conjugates containing the, for example, *Gaussia* or *Renilla mulleri* or *reniformis*, luciferase. A second composition containing the remaining components of a bioluminescence generating system, such as
30 system that emits a wavelength of light within the excitation range of the GFP, such as species of *Renilla* or *Aequorea*, for exciting the *Renilla* luciferase, which

-107-

produces green light that is detected by the photodetector of the device to indicate the presence of the agent.

In further embodiments, the kits contain the components of the diagnostic systems. The kits comprise compositions containing the conjugates, preferably *Renilla* GFP and a *Gaussia*, or *Pleuromamma* or *Renilla mulleri* luciferase and remaining bioluminescence generating system components. The first composition in the kit typically contains the targeting agent conjugated to a GFP or luciferase. The second composition, contains at least the luciferin (substrate) and/or luciferase. Both compositions are formulated for systemic, local or topical application to a mammal. In alternative embodiments, the first composition contains the luciferin linked to a targeting agent, and the second composition contains the luciferase or the luciferase and a GFP.

In general, the packaging is non-reactive with the compositions contained therein and where needed should exclude water and or air to the degree those substances are required for the luminescent reaction to proceed.

Diagnostic applications may require specific packaging. The bioluminescence generating reagents may be provided in pellets, encapsulated as micro or macro-capsules, linked to matrices, preferably biocompatible, more preferably biodegradable matrices, and included in or on articles of manufacture, or as mixtures in chambers within an article of manufacture or in some other configuration. For example, a composition containing luciferase conjugate will be provided separately from, and for use with, a separate composition containing a bioluminescence substrate and bioluminescence activator.

Similarly, the *Renilla reniformis* GFP and selected luciferase and/or luciferin, such as a *Pleuromamma*, *Renilla mulleri* or *Gaussia* luciferase or luciferin, may be provided in a composition that is a mixture, suspension, solution, powder, paste or other suitable composition separately from or in combination with the remaining components, but in the absence of an activating component. Upon contacting the conjugate, which has been targeted to a selected tissue, with this composition the reaction commences and the tissue glows. In preferred embodiments, the tissue glows green emitting light near 510 nm. The luciferase, GFP and bioluminescence substrate, for example, are

-108-

packaged to exclude water and/or air, the bioluminescence activator. Upon administration and release at the targeted site, the reaction with salts or other components at the site, including air in the case of surgical procedures, will activate the components. In some embodiments, it is desirable to provide at least the GFPs or one component of the bioluminescence generating system linked to a matrix substrate, which can then be locally or systemically administered.

Suitable dispensing and packaging apparatus and matrix materials are known to those of skill in the art, and preferably include all such apparatus described in U.S. Patent Nos. see U.S. Patent Nos. 5,876,995, 6,152,358 and 6,113,886.

J. Muteins

Muteins of the *Renilla reniformis* GFP are provided herein. Muteins in which conservative amino acid changes that do not alter its ability to act as an acceptor of energy generated by a *Renilla* luciferase/substrate reaction are provided. Also provided are muteins with altered properties, including muteins with altered spectral properties, muteins with altered surface properties that reduce multimerization, including dimerization.

1. Mutation of GFP surfaces to disrupt multimerization

Figure 5 depicts the three anthozoan fluorescent protein for which a crystal structure exists, another available commercially from Clontech as dsRed (also known as drFP583, as in this alignment) (Wall *et al.* (2000); *Nature Struct. Biol.* 7:1133-1138; Yarbrough *et al.* (2001) *Proc. Natl. Acad. Sci. U.S.A.* 98: 462-467). A dark gray background depicts amino acid conservation, and a light gray background depicts shared physiochemical properties. These crystal structures and biochemical characterization (Baird *et al.* (2000) *Proc. Natl. Acad. Sci. U.S.A.* 97: 11984-11989) show that dsRed exists as an obligate tetramer *in vitro*. Evidence also exists that dsRed multimerizes in living cells (Baird *et al.* (2000) *Proc. Natl. Acad. Sci. U.S.A.* 97: 11984-11989). Sedimentation and native gel electrophoresis studies indicate that *Ptilosarcus* and *Renilla mullerei* GFPs also form tetramers *in vitro* and multimerize *in vivo*. *Ptilosarcus* and *Renilla mullerei* GFPs diverge strongly in amino acid sequence from dsRed (39% and

-109-

38% identical, respectively). Computational polypeptide threading algorithms predict that these GFPs fold into essentially the same structure as dsRed, and also the much more sequence divergent *Aequorea victoria* GFP. *Renilla reniformis* GFP is similarly related in sequence to dsRed, *Ptilosarcus* and *Renilla mullerei* GFPs (37%, 51% and 53% identical, respectively), and thus is extremely likely to form similar multimers. Multimerization is undesirable for many applications that use GFP as the reporting moiety in chimeric protein fusion. Hence muteins in which the capacity to multimerize is reduced are provided. Thus provided are mutations of *Renilla reniformis* GFP that disrupt the formation of GFP multimers. Such mutations may also be effected in the *Ptilosarcus* and *Renilla mullerei* and other GFPs (see Figure 6).

Two interaction surfaces within the dsRed tetramer, one primarily hydrophobic (residues marked by X) and one primarily hydrophilic (residues marked by O) have been described (see, Wall *et al.* (2000); *Nature Struct. Biol.* 7:1133-1138). In general, the corresponding residues vary considerably between the 4 GFPs in a complex way, although the physicochemical properties of the amino acids are often conserved. There are a few clusters of conserved residues, especially between *Ptilosarcus* and *Renilla mullerei* GFPs, in keeping with their 77% overall identity.

The scheme provided herein for disruption focuses on altering surface amino acid side chains so that the surfaces acquire or retain a hydrophilic character, and are also altered in their stereo-chemistry (the sizes of the side chains are altered). These GFP surface regions roughly map to the β -sheet secondary structures that comprise the GFP β -barrel tertiary structure. It is thus essential that the secondary structure in any surface mutants be retained, so that the choice of amino acid side chain substitutions is governed by this consideration.

It is also desirable to introduce mutations that alter charge. For example, such mutations are those in which R, H and K residues have been replaced with D, such that the hydrophobic and hydrophilic surfaces now each contain 3 mutated residues (SEQ ID No. 33; Lys to Asp at amino acids 108, 127 and 226, Arg to Asp at amino acids 131 and 199; His to Asp at amino acid 172).

-110-

Site directed mutagenesis techniques are used to introduce amino acid side chains that are amenable to aqueous solvation, and that significantly alter surface stereochemistry. Disruption of interacting surfaces involves loss-of-function mutagenesis. It is thus contemplated that altering only a few residues, perhaps even one, is sufficient.

2. Use of advantageous GFP surfaces with substituted fluorophores

Other surfaces of GFPs may be key determinants of GFP usefulness as reporters in living systems. A GFP surface may adventitiously interact with vital cellular components, thereby contributing to GFP-induced cytotoxicity. Anthozoan GFPs from bioluminescent luciferase-GFP systems serve fundamentally different biological functions than do anthozoan GFPs from coral and anemones. The *Renilla reniformis* GFP is present in low quantity and functions as a resonance energy acceptor in response to a dynamic neural network that enables a startled animal to emit light flashes. A coral GFP-like protein is present in large quantity and apparently is used primarily as a passive pigment; it may not have evolved to dynamically interact with sensitive cellular machinery. These two classes of anthozoan fluorescent proteins thus may have surfaces with markedly different biological properties.

Figure 4 exemplifies the site for substitution for inserting fluorophores into the background of *Ptilosarcus*, *Renilla mullerei* and *Renilla reniformis* GFPs. In particular, the 20 amino acid region that lies between two highly conserved prolines with the corresponding 20 amino acid region from any other anthozoan GFP (the underlined regions corresponds to amino acids 56-75 of SEQ ID No. 27 *Renilla reniformis* GFP; amino acids 59-78 of SEQ ID No. 16 *Renilla mulleri* GFP; and amino acids 59-78 of SEQ ID No. 32 for *Ptilosarcus* GFP) is replaced or modified. These 20 residues comprise the bulk of a polypeptide region that threads along the interior of the β -barrel structure that is characteristic of anthozoan GFPs (Wall *et al.* (2000) *Nature Struct. Biol.* 7:1133-1138; Yarbrough *et al.* (2001) *Proc. Natl. Acad. Sci. U.S.A.* 98: 462-467); replacement or modification alters spectral properties.

-111-

K. Transgenic plants and animals

As discussed above, transgenic animals and plants that contain the nucleic acid encoding the *Renilla reniformis* GFP are provided. Methods for producing transgenic plants and animals that express a GFP are known (see, 5 *e.g.*, U.S. Patent No. 6,020,538).

Among the transgenic plants and animals provided are those that are novelty items, such as animals with eyes or fingernails or tusks or hair that glows fluorescently. Transgenic food animals, such as chickens and cows and pigs are contemplated from which glowing meat and eggs (green eggs and ham) 10 can be obtained; glowing worms can serve as fishing lures. In addition, the *Renilla reniformis* can serve as a reporter to detect that a heterologous gene linked to the GFP gene is incorporated into the animal's genome or becomes part of the genome in some or all cells. The *Renilla reniformis* can similarly be used as a reporter for gene therapy. The GFP can be introduced into plants to make 15 transgenic ornamental plants that glow, such as orchids and roses and other flowering plants. Also the GFP can be used as a marker in plants, such as by linking it to a promoter, such as Fos that responds to secondary messages to assess signal transduction. The GFP can be linked to adenylycyclase causing the plants to emit different spectral frequencies as the levels of adenylycyclase 20 change.

L. Bioluminescence Resonance Energy Transfer (BRET) System

In nature, coelenterazine-using luciferases emit broadband blue-green light (max. ~480 nm). Bioluminescence Resonance Energy Transfer (BRET) is a natural phenomenon first inferred from studies of the hydrozoan *Obelia* (Morin & 25 Hastings (1971) *J. Cell Physiol.* 77:313-18), whereby the green bioluminescent emission observed *in vivo* was shown to be the result of the luciferase non-radiatively transferring energy to an accessory green fluorescent protein (GFP). BRET was soon thereafter observed in the hydrozoan *Aequorea victoria* and the anthozoan *Renilla reniformis*. Although energy transfer *in vitro* between purified 30 luciferase and GFP has been demonstrated in *Aequorea* (Morise *et al.* (1974) *Biochemistry* 13:2656-62) and *Renilla* (Ward & Cormier (1976) *J. Phys. Chem.* 80:2289-91) systems, a key difference is that in solution efficient radiationless

-112-

- energy transfer occurs only in *Renilla*, apparently due to the pre-association of one luciferase molecule with one GFP homodimer (Ward & Cormier (1978) Photochem. Photobiol. 27:389-96). The blue (486 nm) luminescent emission of *Renilla* luciferase can be completely converted to narrow band green emission (508 nm) upon addition of proper amounts of *Renilla* GFP (Ward & Cormier (1976) *J. Phys. Chem.* 80:2289-91). GFPs accept energy from excited states of luciferase-substrate complexes and re-emit the light as narrow-band green light (~510 nm). By virtue of the non-radiative energy transfer, the quantum yield of the luciferase is increased.
- 10 Luciferases and fluorescent proteins have many well-developed and valuable uses as protein tags and transcriptional reporters; BRET increases the sensitivity and scope of these applications. A GFP increases the sensitivity of the luciferase reporter by raising the quantum yield. A single luciferase fused (or chemically linked) to several spectrally distinct GFPs provides for the
- 15 simultaneous use of multiple luciferase reporters, activated by addition of a single luciferin. By preparing two fusion proteins (or chemical conjugates), each containing a GFP having a different emission wavelength fused to identical luciferases, two or more reporters can be used with a single substrate addition. Thus multiple events may be monitored or multiple assays run using a single
- 20 reagent addition. Such a reporter system is self-ratioing if the distribution of luciferin is uniform or reproducible.

- The ability to conveniently monitor several simultaneous macromolecular events within a cell is a major improvement over current bioluminescent technology. BRET also enables completely new modes of reporting by exploiting
- 25 changes in association or orientation of the luciferase and fluorescent protein. By making fusion proteins, the luciferase-GFP acceptor pair may be made to respond to changes in association or conformation of the fused moieties and hence serves as a sensor.

- Energy transfer between two fluorescent proteins (FRET) as a
- 30 physiological reporter has been reported (Miyawaki *et al.* (1997) *Nature* 388:882-7), in which two different GFPs were fused to the carboxyl and amino termini of calmodulin. Changes in calcium ion concentration caused a sufficient

-113-

conformational change in calmodulin to alter the level of energy transfer between the GFP moieties. The observed change in donor emission was ~10% while the change in ratio was ~1.8.

Figure 2, reproduced from allowed copending application U.S. application
5 Serial No. 09/277,716, illustrates the underlying principle of Bioluminescent Resonance Energy Transfer (BRET) using GFPs and luciferase, preferably cognate luciferase, and its use as sensor: A) in isolation, a luciferase, preferably an anthozoan luciferase, emits blue light from the coelenterazine-derived chromophore; B) in isolation, a GFP, preferably an anthozoan GFP that binds to
10 the luciferase, that is excited with blue-green light emits green light from its integral peptide based fluorophore; C) when the luciferase and GFP associate as a complex *in vivo* or *in vitro*, the luciferase non-radiatively transfers its reaction energy to the GFP fluorophore, which then emits the green light; D) any
15 molecular interaction that disrupts the luciferase-GFP complex can be quantitatively monitored by observing the spectral shift from green to blue light. Hence, the interaction or disruption thereof serves as a sensor.

The similar use of a luciferase-GFP pair in the presence of substrate luciferin has important advantages. First, there is no background and no
20 excitation of the acceptor from the primary exciting light. Second, because the quantum yield of the luciferase is greatly enhanced by nonradiative transfer to GFP, background from donor emission is less, and the signal from the acceptor relatively greater. Third, the wavelength shift from the peak emission of luciferase (~480 nm) to that of the GFP (typically 508-510 nm) is large, minimizing signal overlap. All three factors combine to increase the signal-to-
25 noise ratio. The concentration of the GFP acceptor can be independently ascertained by using fluorescence.

For some applications, *in vitro* crosslinked or otherwise *in vitro* modified versions of the native proteins is contemplated. The genetically encoded fusion
30 proteins have many great advantages: A) *In vivo* use - unlike chemistry-based luminescence or radioactivity-based assays, fusion proteins can be genetically incorporated into living cells or whole organisms. This greatly increases the range of possible applications; B) Flexible and precise modification - many

-114-

different response modifying elements can be reproducibly and quantitatively incorporated into a given luciferase-GFP pair; C) Simple purification - only one reagent would need to be purified, and its purification could be monitored via the fluorescent protein moiety. Ligand-binding motifs can be incorporated to

5 facilitate affinity purification methods.

1. Design of sensors based on BRET

Resonance energy transfer between two chromophores is a quantum mechanical process that is exquisitely sensitive to the distance between the donor and acceptor chromophores and their relative orientation in space (Wu &

10 Brand (1994) *Anal. Biochem.* 218:1-13). Efficiency of energy transfer is inversely proportional to the 6th power of chromophore separation. In practice, the useful distance range is about 10 to 100 Å, which has made resonance energy transfer a very useful technique for studying the interactions of biological macromolecules. A variety of fluorescence-based FRET biosensors have been

15 constructed, initially employing chemical fluors conjugated to proteins or membrane components, and more recently, using pairs of spectrally distinct GFP mutants (Giuliano & Taylor (1998) *Trends Biotech.* 16:99-146; Tsien (1998) *Annu. Rev. Biochem.* 67:509-44).

Although these genetically encoded GFP bioluminescence-based

20 biosensors have advantages over less convenient and less precise chemical conjugate-based biosensors, all share a limitation in their design: it is generally difficult to construct a biosensor in which energy transfer is quantitative when the chromophores are in closest apposition. It is almost impossible to arbitrarily manipulate the complex stereochemistry of proteins so that conjugated or

25 intrinsic chromophores are *stably positioned with minimal separation and optimal orientation*. The efficiency of such biosensors are also often limited by stoichiometric imbalances between resonance energy donor and acceptor; the donor and acceptor macromolecules must be quantitatively complexed to avoid background signal emanating from uncomplexed chromophores. These

30 limitations in general design become important when biosensors must be robust, convenient and cheap. Developing technologies such as high throughput screening for candidate drugs (using high throughput screening (HTS)

-115-

protocols), biochips and environmental monitoring systems would benefit greatly from modular biosensors where the signal of a rare target "hit" (e.g., complex formation between two polypeptides) is unambiguously (statistically) distinguishable from the huge excess of "non-hits"). Current genetically

- 5 encoded FRET and bioluminescence-based biosensors display hit signals that very often are less than two-fold greater than non-hit signals, and are at best a few-fold greater (Xu *et al.* (1999) *Proc. Natl. Acad. Sci USA* 96: 151-156; Miyawaki *et al.* (1997) *Nature* 388:882-7).

- To solve these problems, the anthozoan GFPs, particularly the *Renilla*
10 GFPs, provided herein can be used in combination with its cognate luciferases. Anthozoan luciferases-GFP complexes provide a "scaffold" upon which protein domains that confer the biological properties specific to a given biosensor can be linked. Although one can construct many useful two component biosensors based on this scaffold, in a biosensor contemplated herein, independent protein
15 domains that potentially complex with one another are respectively fused to the luciferase and the GFP.

- There are many possible variations on this theme. For example, in a three component system either the luciferase or GFP can be fused to a ligand-binding domain from a protein of interest or other target peptide or other moiety of
20 interest. If the design of the fusion protein is correct, binding of a small molecule or protein ligand then prevents the luciferase-GFP association. The resulting combination of elements is a BRET-based biosensor; the change in spectral properties in the presence and absence of the ligand serves as sensor. More complex protein fusions can be designed to create two component and
25 even single component BRET biosensors for a multitude of uses.

- The nucleic acids, and the constructs and plasmids herein, permit preparation of a variety of configurations of fusion proteins that include an anthozoan GFP, in this case *Renilla reniformis*, preferably with a *Renilla* luciferase, more preferably with the *Renilla reniformis* luciferase. The nucleic
30 acid encoding the GFP can be fused adjacent to the nucleic acid encoding the luciferase or separated therefrom by insertion of nucleic acid encoding, for example, a ligand-binding domain of a protein of interest. The GFP and

-116-

luciferase will be bound. Upon interaction of the ligand-binding domain with the a test compound or other moiety, the interaction of the GFP and luciferase will be altered thereby changing the emission signal of the complex. If necessary the GFP and luciferase can be modified to fine tune the interaction to make it more
5 sensitive to conformational changes or to temperature or other parameters.

2. BRET Sensor Architectures

Figure 3 depicts some exemplary BRET sensor architectures. The upper left panel depicts the luciferase-GFP scaffold, the basis for the representative BRET sensor architectures shown here. The depicted single polypeptide fusion
10 constructs place the luciferase and GFP at the polypeptide termini, bracketing interacting protein domains of choice. The luciferase and GFP can alternatively be placed centrally within the polypeptide, between interacting protein domains (not shown). This alternative arrangement is advantageous for one step protein interaction-based cloning schemes, where cDNA fragments encoding potential
15 protein targets can be ligated onto one end of the construct.

Single polypeptide sensors that detect conformational changes within protein targets or the association-dissociation of protein targets are well-suited for the detection of physiological signals, such as those mediated by phosphorylation or other modification of targets, or by binding of regulatory
20 ligands, such as hormones, to targets. Sensors based on interference are best suited to assaying the presence of small molecules or proteins independent of any regulatory context. Quantitative assays of metabolites, such as a sugar and allergens, are among those contemplated.

Since *in vivo* and *in vitro* luciferase-to-GFP energy transfer can be nearly
25 100% efficient, binding interactions between the luciferase and GFP must be sufficient to establish an optimal spatial relationship between donor and acceptor chromophores. Optimization of the luciferase-GFP energy transfer module is important in building effective BRET sensors. In a single polypeptide sensor it is crucial that the luciferase-GFP interaction be weak relative to interactions
30 between target domains, thus the need for an optimized energy transfer module. In practice, either the luciferase or GFP surface can be randomly mutagenized, and an optimized luciferase-GFP scaffold then selected by screening for either

-117-

blue or green emission at two near physiological temperatures (thermal endpoint-selection) using current robotic systems. This disruption of BRET is readily achievable because loss-of-function mutants (weakened luciferase-GFP binding) are orders of magnitude more frequent than gain-of-function mutants.

5 With an optimized energy transfer scaffold in hand, thermal endpoint-selection can then be used, if necessary, to optimize the interactions between the target domains incorporated into a sensor. This second round of thermal endpoint-selection may be especially important for the construction of interference sensors because it is essential that such sensors be able to “open
10 and close” at near physiological temperatures to sense interference. Thermal endpoint-selection can also be used to weaken the binding affinity of the analyte to the interference sensor, making it possible to thermally wash off the analyte and reuse the sensor, a great advantage for biochip-based applications.

3. Advantages of BRET sensors

15 There are many advantages to the BRET sensors provided herein. For example, BRET sensors are self-ratioing. The reporter and target are integrated into single polypeptide. This ensures 1:1:1 stoichiometry among luciferase, GFP and target (or a 1:N:1 stoichiometry if more than one, typically a homodimer, GFP can be bound to a luciferase). GFP fluorescence allows
20 absolute quantitation of sensor. The null state gives signal that verifies sensor functionality. Quantifiable null state facilitates disruption-of-BRET sensors (DBRET). BRET sensors have better signal-to-noise ratio than GFP FRET sensors because there is no cellular autofluorescence, no excitation of the acceptor from the primary exciting light, the quantum yield of luciferase greatly enhanced by
25 non-radiative energy transfer to GFP, and there is minimal signal overlap between emission of the luciferase and emission of the GFP. Also, anthozoan GFPs have 6-fold higher extinction coefficients than Aequorea GFP.

 The BRET sensors can be used for hit identification and downstream evaluation in *in vitro* screening assays in *in vitro* or *in vivo* or *in situ*, including in
30 cultured cells and tissues and animals. The BRET sensors can be created by thermal endpoint-selection, which is suited to

-118-

DBRET (Disruption-of-BRET) and reduces need for knowledge of target 3D structure and functional dynamics. Existing screening robotics can be used to optimize biosensors. BRET sensors benefit from vast genetic diversity anthozoans have evolved efficient luciferase-GFP energy transfer systems and the components can be mixed and matched. Highly efficient heterologous luciferases may be substituted for less active luciferases. For example, a copepod luciferase active site can be fused to an anthozoan luciferase GFP-binding domain. There are many diverse coelenterazine-using luciferases.

BRET sensors are modular so that an optimized sensor scaffold may be used with different targets. Also the BRET acceptor may be varied to give shifted emissions, facilitating multiple simultaneous readouts. The anthozoan GFPs can be mutated, GFPs or other proteins can be modified with different chemical fluors, high throughput screening (HTS) fluor-modified FRET acceptors can be adapted, and the BRET donor (luciferase) may be varied, such as by using an Aequorin (Ca⁺⁺ activated) photoprotein, or a firefly luciferase (requires ATP and a firefly luciferin) to give conditional activation. The sensor scaffold can be incorporated into a variety of immobilization motifs, including free format plates, which can reduce reagent volumes, reusable microtiter plates, miniature columns and biochips. Finally, BRET sensors are inexpensive and reproducible reagents because they can be produced by standardized protein production and can incorporate purification tags. Genetically encoded reporters more reproducible than chemically modified reporters. Linear translation of BRET modules ensures sensor integrity.

The following example is included for illustrative purposes only and is not intended to limit the scope of the invention.

EXAMPLE

Specimens of the sea pansy *Renilla reniformis* were collected from inshore waters off the coast of Georgia. To prepare the sea pansies for isolation of mRNA, about 25 or so at time were placed on a large bed of dry ice. They were flipped with a spatula to flip them over to prevent them from freezing. Oddly, the entire animal illuminated when it came in contact with the dry ice.

RECTIFIED SHEET (RULE91)
ISA/EP

-119-

The brightest and greenest were culled, placed in a bag and back into sea water at about 65-70° C for two hours. This process of dry ice, culling and sea water treatment was repeated three times over a 6 hour period. In addition, the process was performed at night. After they with exhausted were the last
5 chilling, the culled animals were frozen solid. A cDNA library was prepared from the frozen animals.

The animals that were selected this way were frozen in liquid nitrogen, and shipped to Stratagene, Inc. (La Jolla, Ca), a commercial vendor whose business includes the construction of custom cDNA libraries under contract to
10 prepare the library. Purified polyA-mRNA was prepared, and a cDNA synthesis reaction was performed, appending a 3' *XhoI* site and a 5' *EcoRI* restriction site to the cDNA. The cDNA was inserted by ligation between the *EcoRI* and *XhoI* sites of the Uni-ZAP Lambda phage cDNA cloning vector.

The resulting unamplified library contained approximately 1.6×10^8
15 primary plaques, which after amplification gave a titer of 3.5-7.5 pfb (plaque forming units)/ml. Insert sizes ranged from 0.9 to 3.0 kb, with an average size around 1.5 kb. Two mass excisions were performed to give pBluescript phagemid containing the cDNA inserts; each excision from about 8×10^6 plaques gave rise to about 4.8×10^9 cfu (colony forming units)/ml. Phagemids were
20 transfected into the SOLR strain of *E. coli*.

Screening was performed by plating (using an artist's airbrush) approximately 200,000 colonies to each of 40 cafeteria trays containing LB agar medium incorporating 0.4% carbon black to absorb background fluorescence. After 24 hours growth at 30° C in a humidified incubator, GFP expressing
25 colonies were identified by illuminating the plates using a 250 Watt quartz halogen fiber optics light (Cuda Products Corp) with an EGFP bandpass excitation filter (Chroma), and viewing colonies through a GFP bandpass emission filter. Approximately 10 fluorescent colonies were picked, DNA isolated from minipreps, and the DNA transformed into the XL-10 Gold *E. coli* strain
30 (Stratagene). Analysis by restriction digestion resolved three distinguishable sizes of insert. DNA was prepared from a clone of each size class and sent to

-120-

SeqWright LLB (Houston, TX) for sequencing. Sequencing data were reported to Prolume on 1-25-99.

Three independent cDNA clones of *Renilla reniformis* GFP were isolated (SEQ ID Nos 23-25). Each cDNA is full length as judged by identical 5' termini
5 and each encodes an identical protein of 233 amino acids (see SEQ ID No. 27). Compared to the primary clone (Clone 1), the coding sequence of Clone 2 differs by 4 silent mutations. Clones 2 and 3 also contain small differences in the 5' and 3' untranslated regions of the cDNA. This nucleic acid has been inserted into expression vector, and the encoded protein produced.

10 Since modifications may be apparent to those of skill in the art, it is intended that the invention be limited only by the appended claims.

-121-

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule encoding a *Renilla reniformis* green fluorescent protein, comprising a sequence of nucleotides that encodes the protein of SEQ ID No. 27 or a green fluorescent protein encoded by a *Renilla*
5 *reniformis* having at least 80% sequence identity thereto.
2. An isolated nucleic acid molecule of claim 1 that encodes a protein having at least 90% sequence identity to the protein of SEQ ID No. 27.
3. The isolated nucleic acid molecule of claim 1, comprising a sequence of nucleotides selected from the group consisting of:
10 (a) the coding portion of the sequence of nucleotides set forth in any of SEQ ID Nos. 23-25;
(b) a sequence of nucleotides that hybridizes under high stringency to the sequence of nucleotides of (a); and
(c) a sequence of nucleotides comprising degenerate codons of (a) or (b).
- 15 4. The isolated nucleic acid molecule of claim 1, wherein the nucleic acid is DNA.
5. The isolated nucleic acid molecule of claim 1, wherein the nucleic acid is RNA.
6. A nucleic acid probe or primer, comprising at least 14 contiguous
20 nucleotides selected from the sequence of nucleotides set of claim 1.
7. The probe or primer of claim 6, comprising at least 16 contiguous nucleotides selected from the sequence of nucleotides in claim 1.
8. The probe or primer of claim 7, comprising at least 30 contiguous nucleotides.
- 25 9. A plasmid, comprising the sequence of nucleotides of claim 1.
10. The plasmid of claim 8 that is an expression vector, comprising:
a promoter element;
a cloning site for the introduction of nucleic acid; and
a selectable marker;
30 wherein the nucleic acid encoding the cloning site is positioned between nucleic acids encoding the promoter element and the green fluorescent protein

-122-

and wherein the nucleic acid encoding the green fluorescent protein is operatively linked to the promoter element.

11. The plasmid of claim 9, further comprising a sequence of nucleotides encoding a luciferase.
- 5 12. A recombinant host cell, comprising the plasmid of claim 9.
13. The cell of claim 12, wherein the cell is selected from the group consisting of a bacterial cell, a yeast cell, a fungal cell, a plant cell, an insect cell and an animal cell.
14. An isolated substantially purified *Renilla reniformis* green
- 10 fluorescent protein (GFP) encoded by the nucleic acid molecule of claim 1.
15. A mutein of the GFP of claim 14 that exhibits altered spectral properties.
16. A mutein of the GFP of claim 14 that exhibits a reduced tendency to form multimers.
- 15 17. A composition, comprising the green fluorescent protein of claim 14 and at least one component of a bioluminescence generating system.
18. The composition of claim 17, wherein the bioluminescence generating system is selected from those isolated from: an insect system, a coelenterate system, a ctenophore system, a bacterial system, a mollusk
- 20 system, a crustacea system, a fish system, an annelid system, and an earthworm system.
19. The composition of claim 17, wherein the bioluminescence generating system is selected from those isolated from: fireflies, *Mnemiopsis*, *Beroe ovata*, *Aequorea*, *Obelia*, *Vargula*, *Pelagia*, *Renilla*, *Pholas* *Aristostomias*,
- 25 *Pachystomias*, *Porichthys*, *Cypridina*, *Aristostomias*, *Pachystomias*, *Malacosteus*, *Gonadostomias*, *Gaussia*, *Watensia*, *Halisturia*, Vampire squid, *Glyphus*, Mycotophids, *Vinciguerria*, *Howella*, *Florenciella*, *Chaudiodus*, *Melanocostus*, Sea Pens, *Chiroteuthis*, *Eucleoteuthis*, *Onychoteuthis*, *Watasenia*, cuttlefish, *Sepiolina*, *Oplophorus*, *Acanthophyra*, *Sergestes*, *Gnathophausia*, *Argyropelecus*,
- 30 *Yarella*, *Diaphus*, *Gonadostomias* and *Neoscopelus*.
20. A mutein of claim 15, comprising substitution in amino acids at amino acids 56-75 of SEQ ID No. 27, whereby spectral properties are altered.

-123-

21. The composition of claim 20, wherein the bioluminescence generating system is selected from those isolated from *Aequorea*, *Obelia*, *Vargula* and *Renilla*.
22. A reporter gene construct, comprising the nucleic acid of claim 1.
- 5 23. A combination, comprising:
an article of manufacture; and
a *Renilla reniformis* green fluorescent protein (GFP) encoded by a nucleic acid molecule of claim 1.
- 10 24. The combination of claim 23, further comprising
at least one component of a bioluminescence generating system, whereby the combination is a novelty item.
25. The combination of claim 24, wherein the component of the bioluminescence generating system comprises a luciferase.
- 15 26. The combination of claim 24, wherein the component of the bioluminescence generating system comprises a luciferin.
27. The combination of claim 23, wherein the article of manufacture is selected from among toys, fountains, personal care items, fairy dust, foods, textile and paper products.
- 20 28. The combination of claim 27, wherein the article of manufacture is selected from among toy guns, pellet guns, greeting cards, fingerpaints, foot bags, slimy play material, clothing, bubble making toys and bubbles therefor, balloons, bath powders, body lotions, gels, body powders, body creams, toothpastes, mouthwashes, soaps, body paints, bubble bath, board game toys, fishing lures, egg-shaped toys, toy cigarettes, dolls, sparklers, magic wand toys,
25 wrapping paper, gelatins, icings, frostings, fairy dust, beer, ornamental transgenic plants, wine, champagne, milk, soft drinks, ice cubes, ice, dry ice, soaps, body paints and bubble bath.
29. The combination of claim 28 that is a transgenic ornamental plant.
30. The combination of claim 28 that is a toy.
- 30 31. The combination of claim 28 that is a food.
32. The combination of claim 28 that is a cosmetic.
33. The combination of claim 28 that is a beverage.

RECTIFIED SHEET (RULE91)
ISA/EP

-124-

34. The combination of claim 24, wherein the article of manufacture is selected from among toys, fountains, personal care items, fairy dust, foods, textile, transgenic ornamental plant and paper products.

35. The combination of claim 34, wherein the article of manufacture is
5 selected from among toy guns, pellet guns, greeting cards, fingerpaints, foot bags, slimy play material, clothing, bubble making toys and bubbles therefor, balloons, bath powders, body lotions, gels, body powders, body creams, toothpastes, mouthwashes, soaps, body paints, bubble bath, board game toys, fishing lures, egg-shaped toys, toy cigarettes, dolls, sparklers, magic wand toys,
10 wrapping paper, gelatins, icings, frostings, fairy dust, beer, wine, champagne, soft drinks, ice cubes, ice, dry ice, soaps, body paints and bubble bath.

36. An antibody that specifically binds to *Renilla reniformis* or a molecule or derivative of the antibody containing the binding domain thereof.

37. The antibody of claim 36 that is a monoclonal antibody.

15 38. A nucleic acid construct, comprising a nucleotide sequence encoding a luciferase and a sequence of nucleotides of claim 1 that encodes a *Renilla reniformis* fluorescent protein (GFP).

39. The nucleic acid construct of claim 38, wherein the luciferase is a *Renilla mulleri* luciferase, a *Gaussia* luciferase or a *Pleuromamma* luciferase.

20 40. The nucleic acid construct of claim 39, wherein the *Gaussia* luciferase is a *Gaussia princeps* luciferase.

41. The nucleic acid construct of claim 38, wherein the luciferase is encoded by:

25 a sequence of nucleotides set forth in SEQ ID No. 17, SEQ ID No. 19, or SEQ ID No. 28;

a sequence of nucleotides encoding the amino acid sequence set forth in SEQ ID No. 18, SEQ ID No. 20 or SEQ ID No. 29; and

30 a sequence of nucleotides that hybridizes under high stringency to the sequence of nucleotides set forth in SEQ ID No. 17, SEQ ID No. 19 or SEQ ID No. 28.

42. The nucleic acid construct of claim 38 that is DNA.

43. The nucleic acid construct of claim 38 that is RNA.

-125-

44. A plasmid, comprising the nucleic acid construct of claim 38.
45. The plasmid of claim 44, further comprising a sequence of nucleotides encoding:
- 5 a promoter element;
- a selectable marker;
- wherein, the sequence of nucleotides encoding the luciferase and GFP is operatively linked to the promoter element, whereby the luciferase and GFP are expressed.
46. The construct of claim 38, wherein the luciferase and the GFP are
- 10 encoded by a polycistronic message.
47. The construct of claim 38, wherein the encoded luciferase and fluorescent protein comprise a fusion protein.
48. The construct of claim 38, wherein the luciferase is *Renilla reniformis* luciferase.
- 15 49. A recombinant host cell, comprising the plasmid of claim 44.
50. The cell of claim 46, wherein the cell is selected from the group consisting of a bacterial cell, a yeast cell, a fungal cell, a plant cell, an insect cell and an animal cell.
51. An isolated substantially purified luciferase and GFP fusion protein,
- 20 wherein the GFP is a *Renilla reniformis* GFP and the fusion protein is encoded by the nucleic acid construct of claim 47.
52. The fusion protein of claim 51, wherein the luciferase is a *Renilla* luciferase.
53. The fusion protein of claim 51, wherein the luciferase is a *Renilla*
- 25 *reniformis* luciferase.
54. A composition, comprising the fusion protein of claim 48.
55. The composition of claim 54, further comprising at least one component of a bioluminescence generating system.
56. The composition of claim 55, wherein the component of the
- 30 bioluminescence generating system is a luciferin.
57. The nucleic acid construct of claim 47, wherein the sequence of nucleotides encoding the luciferase and GFP are not contiguous.

-126-

58. The nucleic acid construct of claim 57, comprising a sequence of nucleotides that encodes a ligand binding domain of a target protein.
59. A biosensor, comprising a GFP protein encoded by the nucleic acid molecule of claim 1 and a luciferase.
- 5 60. The biosensor of claim 59, wherein the luciferase is a *Renilla* luciferase.
61. A biosensor of claim 59, further comprising a modulator.
62. A biosensor, comprising a fusion protein of claim 51.
63. The biosensor of claim 62, wherein the GFP and luciferase in the
10 fusion protein are not contiguous.
64. A bioluminescence resonance energy transfer (BRET) system, comprising:
- (a) a GFP encoded by the nucleic molecule of claim 1;
- (b) a luciferase from which the GFP can accept energy when
15 the GFP and luciferase associate; and
- (c) a luciferin or other substrate of the luciferase.
65. The BRET system of claim 64, further comprising one or more modulators.
66. The BRET system of claim 65, wherein the GFP and luciferase are
20 each attached to a different modulator, or each are attached to the same modulator.
67. The BRET system of claim 65, wherein a conformational change in a modulator causes an increase in the proximity of the luciferase and GFP.
68. The BRET system of claim 65, wherein a conformational change in
25 a modulator causes a decrease in the proximity of the luciferase and GFP.
69. The BRET system of claim 65, wherein the luciferase is *Renilla reniformis* luciferase.
70. A microelectronic device, comprising:
- a substrate;
- 30 a plurality of micro-locations defined on the substrate, wherein each micro-location is for linking a macromolecule;

-127-

- an independent photodetector integrated at or adjacent to each micro-location and optically coupled to each micro-location, each photodetector being configured to generate a sensed signal responsive to the photons of light emitted at the corresponding micro-location when a light-emitting chemical
- 5 reaction occurs at that micro-location, each photodetector being independent from the photodetectors optically coupled to the other micro-locations; and
- an electronic circuit coupled to each photodetector and configured to read the sensed signal generated by each photodetector and to generate output data signals therefrom that are indicative of the light emitted at each
- 10 micro-location by the light-emitting chemical reactions, whereby the device detects photons of light emitted by light-emitting chemical reactions, wherein:
- each micro-location is defined by a portion of the surface; and
- the micro-locations defined on the substrate each comprise a component of a bioluminescence generating system and a green fluorescent protein of claim
- 15 1, whereby photons of light are emitted when a reaction takes place at that micro-location.
71. The device of claim 70, wherein the micro-locations are provided as an array.
72. The device of claim 70, wherein the bioluminescence generating
- 20 system comprises a *Renilla* luciferase.
73. The device of claim 71, wherein the bioluminescence generating system comprises a *Renilla reniformis* luciferase.
74. A method of detecting and identifying analytes in a biological sample, comprising:
- 25 providing the microelectronic device of claims 70;
- attaching a macromolecule or plurality of different macromolecules to the surface at each micro-location on the device, wherein macromolecule is specific for binding to selected analyte that may be present in the biological sample;
- contacting the sample with the surface of the microelectronic device,
- 30 whereby any of the selected analytes that are present in the sample bind to the macromolecule attached to the surface at each micro-location;

-128-

exposing the surface of the microelectronic device to a second macromolecule or plurality thereof bound to the selected analyte already bound to the first macromolecule at each micro-location, wherein the second macromolecule comprises a component of a bioluminescence generating

5 reaction;

initiating the bioluminescence generating reaction by contacting the surface of the device with the remaining components of the bioluminescence generating reaction, wherein the wavelength of the resulting light is shifted by the *Renilla reniformis* GFP; and

10 detecting photons of light emitted by the GFP using a photodetector optically coupled to each micro-location, each photodetector generating a sensed signal representative of the bioluminescence generation at the respective micro-location.

75. A transgenic animal or plant that expresses the *Renilla reniformis*
15 nucleic acid of claim 1.

76. The transgenic animal or plant of claim 75, selected from among fish, worms, monkeys, rodents, goats, pigs, cows, sheep, horses, flowering plants, ornamental plants.

77. The transgenic animal or plant of claim 75 that is an orchid.




FIGURE 1



FIGURE 2A




FIGURE 2C




FIGURE 2B




FIGURE 2D

FIGURE 2

BRET Sensor Architectures




FIGURE 3

Utilization of advantageous GFP surfaces with substituted fluorophores

	60	*	80		
RM-GFP	:	GAPLPFAFDIVSPA	QYGNRTFTKYPNDIS--	:	83
Pt-GFP	:	GGPLPFAFDIVSIA	QYGNRTFTKYPDDIA--	:	83
RR-GFP	:	GAPLPFAFDIVSVA	FSYGNRAYTGYPEEIS--	:	80
GFP484	:	GAPLPFSYDILSNA	QYGNRALTKYPDDIA--	:	83
drFP583	:	GGPLPFAWDILSPQ	FQYGSKVYVKHPADIP--	:	80
asFP595	:	GGPLPFAFHILST	SCMYGSKTFIKYVSGIP--	:	77
dsFP483	:	GGPLPFGWHILCP	QFQYGKAFVHHPDNIH--	:	80
amFP486	:	GGPLAFSFDILST	VFKEYGNRCFTAYPTSMP--	:	82
zFP506	:	GGPLPFAEDILSAA	FNYGNRVFTEYPQDIV--	:	80
zFP538	:	GGPLPSEDILSAG	FKEYGDRIFTTEYPQDIV--	:	80

=====

FIGURE 4

```

R_reni form : ---MDLAKLGLKEVMPKINLEGLVGDHAFSMEGVGECNILEGTQEVKISVTKGAPLPFAFDIVSV : 63
R_mullerei : MSKQI.KNTC.Q...SY.V...I.NN.V.T...C.K...F.N.L.Q.R...H...P : 66
Ptilosarcu : MNRNV.KNT...I.SA.ASV...I.NN.V...F.K...V.F.N.LMQ.R...G...I : 66
drFP583 : ---RSS.NVI...F.RF.VRM...TNG.E.FI...E...RPY...HNT...LK...G...W...L.P : 63
* 20 40 60
*
R_reni form : AFSYGNRAYTGYEEISDYFLQSEPEEGTYERNIRYODGGTAIVKSDISLEDGKEIVNVDFKAKDL : 129
R_mullerei : Q...TF.K..ND...I...A.M...TL.E...LVEIR...N.IED...VYR.EY.GSNF : 132
Ptilosarcu : Q...TF.K..DD.A...V...A.F...L.FE...AIVDIR...D...D...HYK.EYRNGF : 132
drFP583 : Q.Q...SKV.VKH...AD.P...KKL...K...VMNFE...VVT.TQ.S.Q..C...YK.K.IGVNF : 129
* 80 100 120
*
R_reni form : RRMGPVMQDQDIVGMQPSYESMTNVTSVLIGECIIAFKIQTKGHFTYHMRVTYKSKPVEITMELVHF : 195
R_mullerei : PDD...KT.L...IE.F.A...M.NGVLV...V.LVY...NS...YYSK..K.LM...GV.KEF.S... : 198
Ptilosarcu : PSN...KA.L...E.F.VV...M.SGVLV...VDLVY...ES...NYYSK..K.F.R...GG.KEF.E... : 198
drFP583 : PSD...KKTMWEA...T.RL...PRDGLVK...IHK.L...KD...G.YLVEFKSI.MA...APVQL.G.YY : 195
* 140 160 180 2
*
R_reni form : IQHRLVKTNNVDTASGYVVOHETAIAAHSTIKKIEGSLP--- : 233
R_mullerei : ...E.Y.EDGGF-E...E...OMTS.G.PL...HEMV : 238
Ptilosarcu : .H...E.Y.EEG.F-E...E...QLT..G.PL...HEMV : 238
drFP583 : VDSKIDISHNEDYTI...E.Y...RTEGR.HLFL----- : 226
* 220
*

```

FIGURE 5

	20	40	60	80
Aequorea	---	---	---	---
R_mullerei	MSKGEELFTGVVPIILVELD	VNGHKFSVSGEGEDATYKGLTAFICTT	---	---
Ptilosarcu	MSQILKNTCLQEVMSYKVNLEG	VNNHVFVMEGCGKGNILFGLQVQIRVTK	---	---
R_reniiform	MNRNVLTNLTGLREIMSASAVEG	VNNHVFVMEGCGKGNILFGLQVQIRVTK	---	---
drFP583	---	MDLAKLGLKEVMPKINLEGT	VDGDAFVMEGVGEGNILEGTQVKSIVTK	---
drFP593	---	MRSRNVIKFPMKFKVRMEG	VNGHHEFFTEGEGCRPYEGHNTVVKLVTK	---
dsFP483	---	MSSCKNVIKFPMKFKVRMEG	VNGHHEFFTEGEGCRPYEGHNTVVKLVTK	---
cFP484	---	MSSCKSVIKFPMKFKVRMEG	VNGHHEFFTEGEGCRPYEGHNTVVKLVTK	---
asFP595	---	KALTMTMGVIRPDMKIKRMEG	VNGHHEFFTEGEGCRPYEGHNTVVKLVTK	---
amFP486	---	MALSNKFIQDDMKMTYHMDG	VNGHGYFKCTGKGEENPFEGQEMKIEVIE	---
zFP538	---	MAHSKHLKEENTMYHMEG	VNGHKKFVTGEGIGYFFKCKQ	---
zFP506	---	MAQSKHGLTKENTMYHMEG	VNGHKKFVTGEGIGYFFKCKQ	---
Aequorea	---	---	---	---
R_mullerei	RHDFKSAPEGVYQERLTFER	DGNNKTRAEVKEGEG	---	---
Ptilosarcu	SDYFIQSPFAGMTEPILRVED	GGVLEIRDSINLIE	---	---
R_reniiform	ADYFVQSPFAGMTEPILRVED	GGVLEIRDSINLIE	---	---
drFP583	---	SDYFLQSPFEGFTYERNTAY	QDGGTAIRKDSISLED	---
drFP593	---	PDYKLSFPFEGFTYERNTAY	QDGGTAIRKDSISLED	---
dsFP483	---	PDYKLSFPFEGFTYERNTAY	QDGGTAIRKDSISLED	---
cFP484	---	PDYKLSFPFEGFTYERNTAY	QDGGTAIRKDSISLED	---
asFP595	---	ADYFKQSPFEGFTYERNTAY	QDGGTAIRKDSISLED	---
amFP486	---	PDYFKQSPFEGFTYERNTAY	QDGGTAIRKDSISLED	---
zFP538	---	VDYFKNSCPAGYTWRSTLEF	DGNCILQVDSITVSVKENCYHSESEF	---
zFP506	---	VDYFKNSCPAGYTWRSTLEF	DGNCILQVDSITVSVKENCYHSESEF	---
Aequorea	---	---	---	---
R_mullerei	VNRKIRNIEDGSGVQVADHYC	QNTPIG	---	---
Ptilosarcu	GEVILVYKLSNGRNVMSCHMKT	LMKSGK	---	---
R_reniiform	GEVILVYKLSNGRNVMSCHMKT	LMKSGK	---	---
drFP583	---	GEVILVYKLSNGRNVMSCHMKT	LMKSGK	---
drFP593	---	GEVILVYKLSNGRNVMSCHMKT	LMKSGK	---
dsFP483	---	GEVILVYKLSNGRNVMSCHMKT	LMKSGK	---
cFP484	---	GEVILVYKLSNGRNVMSCHMKT	LMKSGK	---
asFP595	---	GEVILVYKLSNGRNVMSCHMKT	LMKSGK	---
amFP486	---	GEVILVYKLSNGRNVMSCHMKT	LMKSGK	---
zFP538	---	GEVILVYKLSNGRNVMSCHMKT	LMKSGK	---
zFP506	---	GEVILVYKLSNGRNVMSCHMKT	LMKSGK	---

D,E,H,K,R	N,Q,S,T	L,I,V,M,F,Y,W	A,G	C,P
polar charged	polar uncharged	non-polar	hydrophobic	small not grouped
			dimerization	hydrophilic
			surfaces	hydrophobic

FIGURE 6

-1/40-

SEQUENCE LISTING

<110> Bryan, Bruce
 Szent-Gyorgyi, Christopher
 Szczepaniak, William
 Prolume, LTD.

<120> RENILLA RENIFORMIS FLOURESCENT PROTEINS, NUCLEIC ACIDS ENCODING THE
 FLOURESCENT PROTEINS AND THE USE THEREOF IN DIAGNOSTICS, HIGH THROUGHPUT
 SCREENING AND NOVELTY ITEMS

<130> 24729-128

<140> Not Yet Assigned
 <141> Herewith

<150> 60/189,691
 <151> 2000-03-15

<150> 09/277,716
 <151> 1999-03-26

<150> 08/757,046
 <151> 1996-11-25

<150> 08/597,274
 <151> 1996-02-06

<150> 08/908,909
 <151> 1997-08-08

<150> 08/990,103
 <151> 1997-12-12

<160> 33

<170> FastSEQ for Windows Version 4.0

<210> 1
 <211> 1196
 <212> DNA
 <213> Renilla reniformis

<220>
 <221> CDS
 <222> (1)...(942)
 <223> Renilla reniformis luciferase

<400> 1
 agc tta aag atg act tcg aaa gtt tat gat cca gaa caa agg aaa cgg 48
 Ser Leu Lys Met Thr Ser Lys Val Tyr Asp Pro Glu Gln Arg Lys Arg
 1 5 10 15
 atg ata act ggt ccg cag tgg tgg gcc aga tgt aaa caa atg aat gtt 96
 Met Ile Thr Gly Pro Gln Trp Trp Ala Arg Cys Lys Gln Met Asn Val
 20 25 30
 ctt gat tca ttt att aat tat tat gat tca gaa aaa cat gca gaa aat 144
 Leu Asp Ser Phe Ile Asn Tyr Tyr Asp Ser Glu Lys His Ala Glu Asn
 35 40 45

-2/40-

gct gtt att ttt tta cat ggt aac gcg gcc tct tct tat tta tgg cga Ala Val Ile Phe Leu His Gly Asn Ala Ala Ser Ser Tyr Leu Trp Arg 50 55 60	192
cat gtt gtg cca cat att gag cca gta gcg cgg tgt att ata cca gat His Val Val Pro His Ile Glu Pro Val Ala Arg Cys Ile Ile Pro Asp 65 70 75 80	240
ctt att ggt atg ggc aaa tca ggc aaa tct ggt aat ggt tct tat agg Leu Ile Gly Met Gly Lys Ser Gly Lys Ser Gly Asn Gly Ser Tyr Arg 85 90 95	288
tta ctt gat cat tac aaa tat ctt act gca tgg ttg aac ttc tta att Leu Leu Asp His Tyr Lys Tyr Leu Thr Ala Trp Leu Asn Phe Leu Ile 100 105 110	336
tac caa aga aga tca ttt ttt gtc ggc cat gat tgg ggt gct tgt ttg Tyr Gln Arg Arg Ser Phe Phe Val Gly His Asp Trp Gly Ala Cys Leu 115 120 125	384
gca ttt cat tat agc tat gag cat caa gat aag atc aaa gca ata gtt Ala Phe His Tyr Ser Tyr Glu His Gln Asp Lys Ile Lys Ala Ile Val 130 135 140	432
cac gct gaa agt gta gta gat gtg att gaa tca tgg gat gaa tgg cct His Ala Glu Ser Val Val Asp Val Ile Glu Ser Trp Asp Glu Trp Pro 145 150 155 160	480
gat att gaa gaa gat att gcg ttg atc aaa tct gaa gaa gga gaa aaa Asp Ile Glu Glu Asp Ile Ala Leu Ile Lys Ser Glu Glu Gly Glu Lys 165 170 175	528
atg gtt ttg gag aat aac ttc ttc gtg gaa acc atg ttg cca tca aaa Met Val Leu Glu Asn Asn Phe Phe Val Glu Thr Met Leu Pro Ser Lys 180 185 190	576
atc atg aga aag tta gaa cca gaa gaa ttt gca gca tat ctt gaa cca Ile Met Arg Lys Leu Glu Pro Glu Glu Phe Ala Ala Tyr Leu Glu Pro 195 200 205	624
ttc aaa gag aaa ggt gaa gtt cgt cgt cca aca tta tca tgg cct cgt Phe Lys Glu Lys Gly Glu Val Arg Arg Pro Thr Leu Ser Trp Pro Arg 210 215 220	672
gaa atc ccg tta gta aaa ggt ggt aaa cct gac gtt gta caa att gtt Glu Ile Pro Leu Val Lys Gly Gly Lys Pro Asp Val Val Gln Ile Val 225 230 235 240	720
agg aat tat aat gct tat cta cgt gca agt gat gat tta cca aaa atg Arg Asn Tyr Asn Ala Tyr Leu Arg Ala Ser Asp Asp Leu Pro Lys Met 245 250 255	768
ttt att gaa tcg gat cca gga ttc ttt tcc aat gct att gtt gaa ggc Phe Ile Glu Ser Asp Pro Gly Phe Phe Ser Asn Ala Ile Val Glu Gly 260 265 270	816
gcc aag aag ttt cct aat act gaa ttt gtc aaa gta aaa ggt ctt cat Ala Lys Lys Phe Pro Asn Thr Glu Phe Val Lys Val Lys Gly Leu His 275 280 285	864

-3/40-

```

ttt tcg caa gaa gat gca cct gat gaa atg gga aaa tat atc aaa tcg      912
Phe Ser Gln Glu Asp Ala Pro Asp Glu Met Gly Lys Tyr Ile Lys Ser
    290                      295                      300

ttc gtt gag cga gtt ctc aaa aat gaa caa taattacttt gggttttttat      962
Phe Val Glu Arg Val Leu Lys Asn Glu Gln
    305                      310

ttacatttttt cccgggttta ataatatataa tgtcattttc aacaatttta ttttaactga 1022
atatttcaca gggaacattc atatatgttg attaatttag ctcgaacttt actctgtcat 1082
atcatttttg aatattacct ctttcaatga aactttataa acagtgggtc aattaattaa 1142
tatatattat aattacattt gttatgtaat aaactcggtt ttattataaa aaaa      1196

<210> 2
<211> 1822
<212> DNA
<213> Cypridina hilagendorfii luciferase

<220>
<221> CDS
<222> (1)...(1665)
<223> Cypridina hilgendorfii luciferase

<400> 2
atg aag cta ata att ctg tct att ata ttg gcc tac tgt gtc aca gtc      48
Met Lys Leu Ile Ile Leu Ser Ile Ile Leu Ala Tyr Cys Val Thr Val
    1                      5                      10                      15

aac tgc cag gat gca tgt cct gta gaa gct gaa gca ccg tca agt aca      96
Asn Cys Gln Asp Ala Cys Pro Val Glu Ala Glu Ala Pro Ser Ser Thr
    20                      25                      30

cca aca gtc cca aca tct tgt gaa gct aaa gaa gga gaa tgt atc gat      144
Pro Thr Val Pro Thr Ser Cys Glu Ala Lys Glu Gly Glu Cys Ile Asp
    35                      40                      45

acc aga tgc gca aca tgt aaa cga gac ata cta tca gac gga ctg tgt      192
Thr Arg Cys Ala Thr Cys Lys Arg Asp Ile Leu Ser Asp Gly Leu Cys
    50                      55                      60

gaa aat aaa cca ggg aag aca tgc tgt aga atg tgc cag tat gta att      240
Glu Asn Lys Pro Gly Lys Thr Cys Cys Arg Met Cys Gln Tyr Val Ile
    65                      70                      75                      80

gaa tcc aga gta gaa gct gct gga tat ttt aga acg ttt tac gcc aaa      288
Glu Ser Arg Val Glu Ala Ala Gly Tyr Phe Arg Thr Phe Tyr Ala Lys
    85                      90                      95

aga ttt aat ttt cag gaa cct ggt aaa tat gtg ctg gct cga gga acc      336
Arg Phe Asn Phe Gln Glu Pro Gly Lys Tyr Val Leu Ala Arg Gly Thr
    100                     105                     110

aag ggt ggc gac tgg tct gta acc ctc acc atg gag aat cta gat gga      384
Lys Gly Gly Asp Trp Ser Val Thr Leu Thr Met Glu Asn Leu Asp Gly
    115                     120                     125

cag aag gga gct gta ctg act aag aca aca ctg gag gta gta gga gac      432
Gln Lys Gly Ala Val Leu Thr Lys Thr Thr Leu Glu Val Val Gly Asp
    130                     135                     140

```

-4/40-

gta	ata	gac	att	act	caa	gct	act	gca	gat	cct	atc	aca	gtt	aac	gga	480
Val	Ile	Asp	Ile	Thr	Gln	Ala	Thr	Ala	Asp	Pro	Ile	Thr	Val	Asn	Gly	
145					150					155					160	
gga	gct	gac	cca	gtt	atc	gct	aac	ccg	ttc	aca	att	ggg	gag	gtg	acc	528
Gly	Ala	Asp	Pro	Val	Ile	Ala	Asn	Pro	Phe	Thr	Ile	Gly	Glu	Val	Thr	
				165					170					175		
att	gct	gtt	gtc	gaa	ata	ccc	ggc	ttc	aat	att	aca	gtc	atc	gaa	ttc	576
Ile	Ala	Val	Val	Glu	Ile	Pro	Gly	Phe	Asn	Ile	Thr	Val	Ile	Glu	Phe	
			180					185					190			
ttt	aaa	cta	atc	gtg	ata	gat	att	ctg	gga	gga	aga	tct	gtg	aga	att	624
Phe	Lys	Leu	Ile	Val	Ile	Asp	Ile	Leu	Gly	Gly	Arg	Ser	Val	Arg	Ile	
		195					200					205				
gct	cca	gac	aca	gca	aac	aaa	gga	ctg	ata	tct	ggg	atc	tgt	ggg	aat	672
Ala	Pro	Asp	Thr	Ala	Asn	Lys	Gly	Leu	Ile	Ser	Gly	Ile	Cys	Gly	Asn	
	210					215					220					
ctg	gag	atg	aat	gac	gct	gat	gac	ttt	act	aca	gac	gca	gat	cag	ctg	720
Leu	Glu	Met	Asn	Asp	Ala	Asp	Asp	Phe	Thr	Thr	Asp	Ala	Asp	Gln	Leu	
225					230					235				240		
gcg	atc	caa	ccc	aac	ata	aac	aaa	gag	ttc	gac	ggc	tgc	cca	ttc	tac	768
Ala	Ile	Gln	Pro	Asn	Ile	Asn	Lys	Glu	Phe	Asp	Gly	Cys	Pro	Phe	Tyr	
				245					250					255		
ggg	aat	cct	tct	gat	atc	gaa	tac	tgc	aaa	ggg	ctc	atg	gag	cca	tac	816
Gly	Asn	Pro	Ser	Asp	Ile	Glu	Tyr	Cys	Lys	Gly	Leu	Met	Glu	Pro	Tyr	
			260					265					270			
aga	gct	gta	tgt	cgt	aac	aat	atc	aac	ttc	tac	tat	tac	act	ctg	tcc	864
Arg	Ala	Val	Cys	Arg	Asn	Asn	Ile	Asn	Phe	Tyr	Tyr	Tyr	Thr	Leu	Ser	
		275					280					285				
tgc	gcc	ttc	gct	tac	tgt	atg	gga	gga	gaa	gaa	aga	gct	aaa	cac	gtc	912
Cys	Ala	Phe	Ala	Tyr	Cys	Met	Gly	Gly	Glu	Glu	Arg	Ala	Lys	His	Val	
	290					295					300					
ctt	ttc	gac	tat	gtt	gag	aca	tgc	gct	gca	ccg	gaa	acg	aga	gga	acg	960
Leu	Phe	Asp	Tyr	Val	Glu	Thr	Cys	Ala	Ala	Pro	Glu	Thr	Arg	Gly	Thr	
305				310						315					320	
tgt	gtt	tta	tca	gga	cat	act	ttc	tat	gac	aca	ttc	gac	aaa	gcc	aga	1008
Cys	Val	Leu	Ser	Gly	His	Thr	Phe	Tyr	Asp	Thr	Phe	Asp	Lys	Ala	Arg	
				325					330					335		
tat	caa	ttc	cag	ggc	cca	tgc	aaa	gag	ctt	ctg	atg	gcc	gca	gac	tgt	1056
Tyr	Gln	Phe	Gln	Gly	Pro	Cys	Lys	Glu	Leu	Leu	Met	Ala	Ala	Asp	Cys	
			340					345					350			
tac	tgg	aac	aca	tgg	gat	gta	aag	gtt	tca	cat	aga	gat	gtt	gag	tca	1104
Tyr	Trp	Asn	Thr	Trp	Asp	Val	Lys	Val	Ser	His	Arg	Asp	Val	Glu	Ser	
		355				360						365				
tac	act	gag	gta	gag	aaa	gta	aca	atc	agg	aaa	cag	tca	act	gta	gta	1152
Tyr	Thr	Glu	Val	Glu	Lys	Val	Thr	Ile	Arg	Lys	Gln	Ser	Thr	Val	Val	
	370					375					380					

-5/40-

gat ttg att gtg gat ggc aag cag gtc aag gtt gga gga gtg gat gta	1200
Asp Leu Ile Val Asp Gly Lys Gln Val Lys Val Gly Gly Val Asp Val	
385 390 395 400	
tct atc ccg tac agt tct gag aac aca tcc ata tac tgg cag gat gga	1248
Ser Ile Pro Tyr Ser Ser Glu Asn Thr Ser Ile Tyr Trp Gln Asp Gly	
405 410 415	
gac atc ctg acg acg gcc atc cta cct gaa gct ctt gtc gtt aag ttc	1296
Asp Ile Leu Thr Thr Ala Ile Leu Pro Glu Ala Leu Val Val Lys Phe	
420 425 430	
aac ttt aag cag ctc ctt gta gtt cat atc aga gat cca ttc gat gga	1344
Asn Phe Lys Gln Leu Leu Val Val His Ile Arg Asp Pro Phe Asp Gly	
435 440 445	
aag aca tgc ggc ata tgt ggt aac tat aat caa gat tca act gat gat	1392
Lys Thr Cys Gly Ile Cys Gly Asn Tyr Asn Gln Asp Ser Thr Asp Asp	
450 455 460	
ttc ttt gac gca gaa gga gca tgc gct ctg acc ccc aat ccc cca gga	1440
Phe Phe Asp Ala Glu Gly Ala Cys Ala Leu Thr Pro Asn Pro Pro Gly	
465 470 475 480	
tgt aca gag gag cag aaa cca gaa gct gag cga ctc tgc aat agt cta	1488
Cys Thr Glu Glu Gln Lys Pro Glu Ala Glu Arg Leu Cys Asn Ser Leu	
485 490 495	
ttt gat agt tct atc gac gag aaa tgt aat gtc tgc tac aag cct gac	1536
Phe Asp Ser Ser Ile Asp Glu Lys Cys Asn Val Cys Tyr Lys Pro Asp	
500 505 510	
cgt att gca cga tgt atg tac gag tat tgc ctg agg gga cag caa gga	1584
Arg Ile Ala Arg Cys Met Tyr Glu Tyr Cys Leu Arg Gly Gln Gln Gly	
515 520 525	
ttc tgt gac cat gct tgg gag ttc aaa aaa gaa tgc tac ata aag cat	1632
Phe Cys Asp His Ala Trp Glu Phe Lys Lys Glu Cys Tyr Ile Lys His	
530 535 540	
gga gac act cta gaa gta cca cct gaa tgc caa taaatgaaca aagatacaga	1685
Gly Asp Thr Leu Glu Val Pro Pro Glu Cys Gln	
545 550 555	
agctaagact actacagcag aagataaaaag agaagctgta gttcttcaaa aacagtatat	1745
tttgatgtac tcattgttta cttacataaaa aataaattgt tattatcata acgtaaagaa	1805
aaaaaaaaa aaaaaaa	1822
 <210> 3	
<211> 1644	
<212> DNA	
<213> <i>Luciola cruciata</i>	
 <220>	
<221> CDS	
<222> (1)...(1644)	
<223> <i>Luciola cruciata</i> luciferase	
 <400> 3	
atg gaa aac atg gaa aac gat gaa aat att gta gtt gga cct aaa ccg	48

-6/40-

Met 1	Glu	Asn	Met	Glu 5	Asn	Asp	Glu	Asn	Ile 10	Val	Val	Gly	Pro	Lys 15	Pro		
ttt	tac	cct	atc	gaa	gag	gga	tct	gct	gga	aca	caa	tta	cgc	aaa	tac		96
Phe	Tyr	Pro	Ile	Glu	Glu	Gly	Ser	Ala	Gly	Thr	Gln	Leu	Arg	Lys	Tyr		
			20					25					30				
atg	gag	cga	tat	gca	aaa	ctt	ggc	gca	att	gct	ttt	aca	aat	gca	gtt		144
Met	Glu	Arg	Tyr	Ala	Lys	Leu	Gly	Ala	Ile	Ala	Phe	Thr	Asn	Ala	Val		
		35					40					45					
act	ggt	gtt	gat	tat	tct	tac	gcc	gaa	tac	ttg	gag	aaa	tca	tgt	tgt		192
Thr	Gly	Val	Asp	Tyr	Ser	Tyr	Ala	Glu	Tyr	Leu	Glu	Lys	Ser	Cys	Cys		
	50					55					60						
cta	gga	aaa	gct	ttg	caa	aat	tat	ggt	ttg	gtt	gtt	gat	ggc	aga	att		240
Leu	Gly	Lys	Ala	Leu	Gln	Asn	Tyr	Gly	Leu	Val	Val	Asp	Gly	Arg	Ile		
65					70					75					80		
gcg	tta	tgc	agt	gaa	aac	tgt	gaa	gaa	ttt	ttt	att	cct	gta	ata	gcc		288
Ala	Leu	Cys	Ser	Glu	Asn	Cys	Glu	Glu	Phe	Phe	Ile	Pro	Val	Ile	Ala		
				85					90					95			
gga	ctg	ttt	ata	ggt	gta	ggt	gtt	gca	ccc	act	aat	gag	att	tac	act		336
Gly	Leu	Phe	Ile	Gly	Val	Gly	Val	Ala	Pro	Thr	Asn	Glu	Ile	Tyr	Thr		
			100					105					110				
tta	cgt	gaa	ctg	gtt	cac	agt	tta	ggt	atc	tct	aaa	cca	aca	att	gta		384
Leu	Arg	Glu	Leu	Val	His	Ser	Leu	Gly	Ile	Ser	Lys	Pro	Thr	Ile	Val		
		115					120					125					
ttt	agt	tct	aaa	aaa	ggc	tta	gat	aaa	gtt	ata	aca	gta	cag	aaa	aca		432
Phe	Ser	Ser	Lys	Lys	Gly	Leu	Asp	Lys	Val	Ile	Thr	Val	Gln	Lys	Thr		
	130					135					140						
gta	act	act	att	aaa	acc	att	gtt	ata	cta	gat	agc	aaa	gtt	gat	tat		480
Val	Thr	Thr	Ile	Lys	Thr	Ile	Val	Ile	Leu	Asp	Ser	Lys	Val	Asp	Tyr		
145					150					155					160		
cga	gga	tat	caa	tgt	ctg	gac	acc	ttt	ata	aaa	aga	aac	act	cca	cca		528
Arg	Gly	Tyr	Gln	Cys	Leu	Asp	Thr	Phe	Ile	Lys	Arg	Asn	Thr	Pro	Pro		
			165					170						175			
ggt	ttt	caa	gca	tcc	agt	ttc	aaa	act	gtg	gaa	gtt	gac	cgt	aaa	gaa		576
Gly	Phe	Gln	Ala	Ser	Ser	Phe	Lys	Thr	Val	Glu	Val	Asp	Arg	Lys	Glu		
		180					185					190					
caa	gtt	gct	ctt	ata	atg	aac	tct	tcg	ggt	tct	acc	ggt	ttg	cca	aaa		624
Gln	Val	Ala	Leu	Ile	Met	Asn	Ser	Ser	Gly	Ser	Thr	Gly	Leu	Pro	Lys		
		195					200					205					
ggc	gta	caa	ctt	act	cac	gaa	aat	aca	gtc	act	aga	ttt	tct	cat	gct		672
Gly	Val	Gln	Leu	Thr	His	Glu	Asn	Thr	Val	Thr	Arg	Phe	Ser	His	Ala		
	210					215					220						
aga	gat	ccg	att	tat	ggt	aac	caa	gtt	tca	cca	ggc	acc	gct	gtt	tta		720
Arg	Asp	Pro	Ile	Tyr	Gly	Asn	Gln	Val	Ser	Pro	Gly	Thr	Ala	Val	Leu		
225					230					235					240		
act	gtc	gtt	cca	ttc	cat	cat	ggt	ttt	ggt	atg	ttc	act	act	cta	ggg		768
Thr	Val	Val	Pro	Phe	His	His	Gly	Phe	Gly	Met	Phe	Thr	Thr	Leu	Gly		

-7/40-

245								250				255				
tat	tta	att	tgt	ggg	ttt	cgt	gtt	gta	atg	tta	aca	aaa	ttc	gat	gaa	816
Tyr	Leu	Ile	Cys	Gly	Phe	Arg	Val	Val	Met	Leu	Thr	Lys	Phe	Asp	Glu	
			260					265					270			
gaa	aca	ttt	tta	aaa	act	cta	caa	gat	tat	aaa	tgt	aca	agt	gtt	att	864
Glu	Thr	Phe	Leu	Lys	Thr	Leu	Gln	Asp	Tyr	Lys	Cys	Thr	Ser	Val	Ile	
		275					280					285				
ctt	gta	ccg	acc	ttg	ttt	gca	att	ctc	aac	aaa	agt	gaa	tta	ctc	aat	912
Leu	Val	Pro	Thr	Leu	Phe	Ala	Ile	Leu	Asn	Lys	Ser	Glu	Leu	Leu	Asn	
		290				295					300					
aaa	tac	gat	ttg	tca	aat	tta	gtt	gag	att	gca	tct	ggc	gga	gca	cct	960
Lys	Tyr	Asp	Leu	Ser	Asn	Leu	Val	Glu	Ile	Ala	Ser	Gly	Gly	Ala	Pro	
		305			310					315					320	
tta	tca	aaa	gaa	gtt	ggg	gaa	gct	gtt	gct	aga	cgc	ttt	aat	ctt	ccc	1008
Leu	Ser	Lys	Glu	Val	Gly	Glu	Ala	Val	Ala	Arg	Arg	Phe	Asn	Leu	Pro	
				325					330					335		
ggg	gtt	cgt	caa	ggg	tat	ggg	tta	aca	gaa	aca	aca	tct	gcc	att	att	1056
Gly	Val	Arg	Gln	Gly	Tyr	Gly	Leu	Thr	Glu	Thr	Thr	Ser	Ala	Ile	Ile	
			340					345					350			
att	aca	cca	gaa	gga	gac	gat	aaa	cca	gga	gct	tct	gga	aaa	gtc	gtg	1104
Ile	Thr	Pro	Glu	Gly	Asp	Asp	Lys	Pro	Gly	Ala	Ser	Gly	Lys	Val	Val	
		355					360					365				
ccg	ttg	ttt	aaa	gca	aaa	gtt	att	gat	ctt	gat	acc	aaa	aaa	tct	tta	1152
Pro	Leu	Phe	Lys	Ala	Lys	Val	Ile	Asp	Leu	Asp	Thr	Lys	Lys	Ser	Leu	
		370				375					380					
ggg	cct	aac	aga	cgt	gga	gaa	gtt	tgt	gtt	aaa	gga	cct	atg	ctt	atg	1200
Gly	Pro	Asn	Arg	Arg	Gly	Glu	Val	Cys	Val	Lys	Gly	Pro	Met	Leu	Met	
		385			390					395					400	
aaa	ggg	tat	gta	aat	aat	cca	gaa	gca	aca	aaa	gaa	ctt	att	gac	gaa	1248
Lys	Gly	Tyr	Val	Asn	Asn	Pro	Glu	Ala	Thr	Lys	Glu	Leu	Ile	Asp	Glu	
				405				410						415		
gaa	ggg	tgg	ctg	cac	acc	gga	gat	att	gga	tat	tat	gat	gaa	gaa	aaa	1296
Glu	Gly	Trp	Leu	His	Thr	Gly	Asp	Ile	Gly	Tyr	Tyr	Asp	Glu	Glu	Lys	
			420					425					430			
cat	ttc	ttt	att	gtc	gat	cgt	ttg	aag	tct	tta	atc	aaa	tac	aaa	gga	1344
His	Phe	Phe	Ile	Val	Asp	Arg	Leu	Lys	Ser	Leu	Ile	Lys	Tyr	Lys	Gly	
		435					440					445				
tac	caa	gta	cca	cct	gcc	gaa	tta	gaa	tcc	gtt	ctt	ttg	caa	cat	cca	1392
Tyr	Gln	Val	Pro	Pro	Ala	Glu	Leu	Glu	Ser	Val	Leu	Leu	Gln	His	Pro	
						455					460					
tct	atc	ttt	gat	gct	ggg	gtt	gcc	ggc	gtt	cct	gat	cct	gta	gct	ggc	1440
Ser	Ile	Phe	Asp	Ala	Gly	Val	Ala	Gly	Val	Pro	Asp	Pro	Val	Ala	Gly	
					470				475						480	
gag	ctt	cca	gga	gcc	gtt	gtt	gta	ctg	gaa	agc	gga	aaa	aat	atg	acc	1488
Glu	Leu	Pro	Gly	Ala	Val	Val	Val	Leu	Glu	Ser	Gly	Lys	Asn	Met	Thr	
				485					490					495		

-8/40-

gaa aaa gaa gta atg gat tat gtt gca agt caa gtt tca aat gca aaa	1536
Glu Lys Glu Val Met Asp Tyr Val Ala Ser Gln Val Ser Asn Ala Lys	
500 505 510	
cgt tta cgt ggt ggt gtt cgt ttt gtg gat gaa gta cct aaa ggt ctt	1584
Arg Leu Arg Gly Gly Val Arg Phe Val Asp Glu Val Pro Lys Gly Leu	
515 520 525	
act gga aaa att gac ggc aga gca att aga gaa atc ctt aag aaa cca	1632
Thr Gly Lys Ile Asp Gly Arg Ala Ile Arg Glu Ile Leu Lys Lys Pro	
530 535 540	
gtt gct aag atg	1644
Val Ala Lys Met	
545	
<210> 4	
<211> 1820	
<212> DNA	
<213> Vargula (cypridina)	
<220>	
<221> CDS	
<222> (1) ... (1665)	
<223> Vargula (cypridina) luciferase	
<400> 4	
atg aag ata ata att ctg tct gtt ata ttg gcc tac tgt gtc acc gac	48
Met Lys Ile Ile Ile Leu Ser Val Ile Leu Ala Tyr Cys Val Thr Asp	
1 5 10 15	
aac tgt caa gat gca tgt cct gta gaa gcg gaa ccg cca tca agt aca	96
Asn Cys Gln Asp Ala Cys Pro Val Glu Ala Glu Pro Pro Ser Ser Thr	
20 25 30	
cca aca gtt cca act tct tgt gaa gct aaa gaa gga gaa tgt ata gat	144
Pro Thr Val Pro Thr Ser Cys Glu Ala Lys Glu Gly Glu Cys Ile Asp	
35 40 45	
acc aga tgc gca aca tgt aaa cga gat ata cta tca gat gga ctg tgt	192
Thr Arg Cys Ala Thr Cys Lys Arg Asp Ile Leu Ser Asp Gly Leu Cys	
50 55 60	
gaa aat aaa cca ggg aag aca tgc tgt aga atg tgc cag tat gtg att	240
Glu Asn Lys Pro Gly Lys Thr Cys Cys Arg Met Cys Gln Tyr Val Ile	
65 70 75 80	
gaa tgc aga gta gaa gca gct ggt tat ttt aga acg ttt tac ggc aaa	288
Glu Cys Arg Val Glu Ala Ala Gly Tyr Phe Arg Thr Phe Tyr Gly Lys	
85 90 95	
aga ttt aat ttt cag gaa cct ggt aaa tat gtg ctg gct agg gga acc	336
Arg Phe Asn Phe Gln Glu Pro Gly Lys Tyr Val Leu Ala Arg Gly Thr	
100 105 110	
aag ggt ggc gat tgg tct gta acc ctc acc atg gag aat cta gat gga	384
Lys Gly Gly Asp Trp Ser Val Thr Leu Thr Met Glu Asn Leu Asp Gly	
115 120 125	

-9/40-

cag aag gga gct gtg ctg act aag aca aca ctg gag gtt gca gga gac Gln Lys Gly Ala Val Leu Thr Lys Thr Thr Leu Glu Val Ala Gly Asp 130 135 140	432
gta ata gac att act caa gct act gca gat cct atc aca gtt aac gga Val Ile Asp Ile Thr Gln Ala Thr Ala Asp Pro Ile Thr Val Asn Gly 145 150 155 160	480
gga gct gac cca gtt atc gct aac ccg ttc aca att ggt gag gtg acc Gly Ala Asp Pro Val Ile Ala Asn Pro Phe Thr Ile Gly Glu Val Thr 165 170 175	528
att gct gtt gtt gaa ata ccg ggc ttc aat atc aca gtc atc gaa ttc Ile Ala Val Val Glu Ile Pro Gly Phe Asn Ile Thr Val Ile Glu Phe 180 185 190	576
ttt aaa cta atc gtg att gat att ctg gga gga aga tct gtc aga att Phe Lys Leu Ile Val Ile Asp Ile Leu Gly Gly Arg Ser Val Arg Ile 195 200 205	624
gct cca gac aca gca aac aaa gga ctg ata tct ggt atc tgt ggt aat Ala Pro Asp Thr Ala Asn Lys Gly Leu Ile Ser Gly Ile Cys Gly Asn 210 215 220	672
ctg gag atg aat gac gct gat gac ttt act aca gat gca gat cag ctg Leu Glu Met Asn Asp Ala Asp Asp Phe Thr Thr Asp Ala Asp Gln Leu 225 230 235 240	720
gcg atc caa ccc aac ata aac aaa gag ttc gac ggc tgc cca ttc tat Ala Ile Gln Pro Asn Ile Asn Lys Glu Phe Asp Gly Cys Pro Phe Tyr 245 250 255	768
ggc aat cct tct gat atc gaa tac tgc aaa ggt ctg atg gag cca tac Gly Asn Pro Ser Asp Ile Glu Tyr Cys Lys Gly Leu Met Glu Pro Tyr 260 265 270	816
aga gct gta tgt cgt aac aat atc aac ttc tac tat tac act cta tcc Arg Ala Val Cys Arg Asn Asn Ile Asn Phe Tyr Tyr Tyr Thr Leu Ser 275 280 285	864
tgt gcc ttc gct tac tgt atg gga gga gaa gaa aga gct aaa cac gtc Cys Ala Phe Ala Tyr Cys Met Gly Gly Glu Glu Arg Ala Lys His Val 290 295 300	912
ctt ttc gac tat gtt gag aca tgc gct gcg ccg gaa acg aga gga acg Leu Phe Asp Tyr Val Glu Thr Cys Ala Ala Pro Glu Thr Arg Gly Thr 305 310 315 320	960
tgt gtt tta tca gga cat act ttc tat gac aca ttc gac aaa gca aga Cys Val Leu Ser Gly His Thr Phe Tyr Asp Thr Phe Asp Lys Ala Arg 325 330 335	1008
tat caa ttc cag ggc cca tgc aag gag att ctg atg gcc gca gac tgt Tyr Gln Phe Gln Gly Pro Cys Lys Glu Ile Leu Met Ala Ala Asp Cys 340 345 350	1056
tac tgg aac aca tgg gat gta aag gtt tca cat aga gac gtc gaa tca Tyr Trp Asn Thr Trp Asp Val Lys Val Ser His Arg Asp Val Glu Ser 355 360 365	1104
tac act gag gta gag aaa gta aca atc agg aaa cag tca act gta gta	1152

-10/40-

Tyr	Thr	Glu	Val	Glu	Lys	Val	Thr	Ile	Arg	Lys	Gln	Ser	Thr	Val	Val		
370						375					380						
gat	ctc	att	gtg	gat	ggc	aag	cag	gtc	aag	gtt	gga	gga	gtg	gat	gta	1200	
Asp	Leu	Ile	Val	Asp	Gly	Lys	Gln	Val	Lys	Val	Gly	Gly	Val	Asp	Val	400	
385					390					395							
tct	atc	ccg	tac	agc	tct	gag	aac	act	tcc	ata	tac	tggt	cag	gat	gga	1248	
Ser	Ile	Pro	Tyr	Ser	Ser	Glu	Asn	Thr	Ser	Ile	Tyr	Trp	Gln	Asp	Gly	415	
				405					410								
gac	atc	ctg	acg	acg	gcc	atc	cta	cct	gaa	gct	ctt	gtc	gtt	aag	ttc	1296	
Asp	Ile	Leu	Thr	Thr	Ala	Ile	Leu	Pro	Glu	Ala	Leu	Val	Val	Lys	Phe	430	
			420					425									
aac	ttt	aag	cag	ctc	ctt	gta	gtt	cat	atc	aga	gat	cca	ttc	gat	gca	1344	
Asn	Phe	Lys	Gln	Leu	Leu	Val	Val	His	Ile	Arg	Asp	Pro	Phe	Asp	Ala	445	
		435					440										
aag	aca	tgc	ggc	ata	tgt	ggc	aac	tat	aat	caa	gat	tca	act	gat	gat	1392	
Lys	Thr	Cys	Gly	Ile	Cys	Gly	Asn	Tyr	Asn	Gln	Asp	Ser	Thr	Asp	Asp	460	
		450				455											
ttc	ttt	gac	gca	gaa	gga	gca	tgc	gct	cta	acc	ccc	aac	ccc	cca	gga	1440	
Phe	Phe	Asp	Ala	Glu	Gly	Ala	Cys	Ala	Leu	Thr	Pro	Asn	Pro	Pro	Gly	480	
					470					475							
tgt	aca	gag	gaa	cag	aaa	cca	gaa	gct	gag	cga	ctt	tgc	aat	aat	ctc	1488	
Cys	Thr	Glu	Glu	Gln	Lys	Pro	Glu	Ala	Glu	Arg	Leu	Cys	Asn	Asn	Leu	495	
				485					490								
ttt	gat	tct	tct	atc	gac	gag	aaa	tgt	aat	gtc	tgc	tac	aag	cct	gac	1536	
Phe	Asp	Ser	Ser	Ile	Asp	Glu	Lys	Cys	Asn	Val	Cys	Tyr	Lys	Pro	Asp	510	
				500				505									
cggt	att	gcc	cga	tgt	atg	tac	gag	tat	tgc	ctg	agg	gga	caa	caa	gga	1584	
Arg	Ile	Ala	Arg	Cys	Met	Tyr	Glu	Tyr	Cys	Leu	Arg	Gly	Gln	Gln	Gly	525	
		515					520										
ttt	tgt	gac	cat	gct	tggt	gag	ttc	aag	aaa	gaa	tgc	tac	ata	aaa	cat	1632	
Phe	Cys	Asp	His	Ala	Trp	Glu	Phe	Lys	Lys	Glu	Cys	Tyr	Ile	Lys	His	540	
		530				535											
gga	gac	act	cta	gaa	gta	cca	cct	gaa	tgt	caa	taa	acgtaca	aagatacaga			1685	
Gly	Asp	Thr	Leu	Glu	Val	Pro	Pro	Glu	Cys	Gln							
					550					555							
agctaaggct	actacagcag	aagataaaaa	agaaactgta	gttccttcaa	aaaccgtgta											1745	
ttttatgtac	tcattgttta	attagagcaa	aataaattgt	tattatcata	acttaaacta											1805	
aaaaaaaaaa	aaaaa															1820	

<210> 5

<211> 958

<212> DNA

<213> Aequorea victoria

<220>

<221> CDS

<222> (115)...(702)

<223> Apoequorin-encoding gene

-11/40-

<400> 5
 gggggggggggg gggggggggggg gggggggggggg gggaatgcaa ttcacatctttg catcaaagaa 60
 ttacatcaaaa tctctagttg atcaactaaa ttgtctcgcac aacaacaagc aaac atg 117
 Met
 1

aca agc aaa caa tac tca gtc aag ctt aca tca gac ttc gac aac cca 165
 Thr Ser Lys Gln Tyr Ser Val Lys Leu Thr Ser Asp Phe Asp Asn Pro
 5 10 15

aga tgg att gga cga cac aag cat atg ttc aat ttc ctt gat gtc aac 213
 Arg Trp Ile Gly Arg His Lys His Met Phe Asn Phe Leu Asp Val Asn
 20 25 30

cac aat gga aaa atc tct ctt gag atg gtc tac aag gca tct gat 261
 His Asn Gly Lys Ile Ser Leu Asp Glu Met Val Tyr Lys Ala Ser Asp
 35 40 45

att gtc atc aat aac ctt gga gca aca cct gag caa gcc aaa cga cac 309
 Ile Val Ile Asn Asn Leu Gly Ala Thr Pro Glu Gln Ala Lys Arg His
 50 55 60 65

aaa gat gct gta gaa gcc ttc ttc gga gga gct gga atg aaa tat ggt 357
 Lys Asp Ala Val Glu Ala Phe Phe Gly Gly Ala Gly Met Lys Tyr Gly
 70 75 80

gtg gaa act gat tgg cct gca tat att gaa gga tgg aaa aaa ttg gct 405
 Val Glu Thr Asp Trp Pro Ala Tyr Ile Glu Gly Trp Lys Lys Leu Ala
 85 90 95

act gat gaa ttg gag aaa tac gcc aaa aac gaa cca acg ctc atc cgt 453
 Thr Asp Glu Leu Glu Lys Tyr Ala Lys Asn Glu Pro Thr Leu Ile Arg
 100 105 110

ata tgg ggt gat gct ttg ttt gat atc gtt gac aaa gat caa aat gga 501
 Ile Trp Gly Asp Ala Leu Phe Asp Ile Val Asp Lys Asp Gln Asn Gly
 115 120 125

gcc att aca ctg gat gaa tgg aaa gca tac acc aaa gct gct ggt atc 549
 Ala Ile Thr Leu Asp Glu Trp Lys Ala Tyr Thr Lys Ala Ala Gly Ile
 130 135 140 145

atc caa tca tca gaa gat tgc gag gaa aca ttc aga gtg tgc gat att 597
 Ile Gln Ser Ser Glu Asp Cys Glu Glu Thr Phe Arg Val Cys Asp Ile
 150 155 160

gat gaa agt gga caa ctc gat gtt gat gag atg aca aga caa cat tta 645
 Asp Glu Ser Gly Gln Leu Asp Val Asp Glu Met Thr Arg Gln His Leu
 165 170 175

gga ttt tgg tac acc atg gat cct gct tgc gaa aag ctc tac ggt gga 693
 Gly Phe Trp Tyr Thr Met Asp Pro Ala Cys Glu Lys Leu Tyr Gly Gly
 180 185 190

gct gtc ccc taagaagctc tacggtggtg atgcacccta ggaagatgat 742
 Ala Val Pro
 195

gtgattttga ataaaacact gatgaattca atcaaaattt tccaaatttt tgaacgattt 802
 caatcgtttg tgttgatttt tgtaattagg aacagattaa atcgaatgat tagttgtttt 862
 tttaatcaac agaacttaca aatcgaaaaa gtataaaaaaa aaaaaaaaaa aaaaaaaaaa 922

-12/40-

aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa

958

<210> 6

<211> 591

<212> DNA

<213> Aequorea victoria

<220>

<221> CDS

<222> (1)...(588)

<223> Recombinant Aequorin AEQ1

<400> 6

atg acc agc gaa caa tac tca gtc aag ctt aca cca gac ttc gac aac	48
Met Thr Ser Glu Gln Tyr Ser Val Lys Leu Thr Pro Asp Phe Asp Asn	
1 5 10 15	
cca aaa tgg att gga cga cac aag cac atg ttt aat ttt ctt gat gtc	96
Pro Lys Trp Ile Gly Arg His Lys His Met Phe Asn Phe Leu Asp Val	
20 25 30	
aac cac aat gga agg atc tct ctt gac gag atg gtc tac aag gcg tcc	144
Asn His Asn Gly Arg Ile Ser Leu Asp Glu Met Val Tyr Lys Ala Ser	
35 40 45	
gat att gtt ata aac aat ctt gga gca aca cct gaa caa gcc aaa cgt	192
Asp Ile Val Ile Asn Asn Gly Ala Thr Pro Glu Gln Ala Lys Arg	
50 55 60	
cac aaa gat gct gta gaa gcc ttc ttc gga gga gct gga atg aaa tat	240
His Lys Asp Ala Val Glu Ala Phe Phe Gly Gly Ala Gly Met Lys Tyr	
65 70 75 80	
ggg gta gaa act gaa tgg cct gaa tac atc gaa gga tgg aaa aga ctg	288
Gly Val Glu Thr Glu Trp Pro Glu Tyr Ile Glu Gly Trp Lys Arg Leu	
85 90 95	
gct tcc gag gaa ttg aaa agg tat tca aaa aac caa atc aca ctt att	336
Ala Ser Glu Glu Leu Lys Arg Tyr Ser Lys Asn Gln Ile Thr Leu Ile	
100 105 110	
cgt tta tgg ggt gat gca ttg ttc gat atc att gac aaa gac caa aat	384
Arg Leu Trp Gly Asp Ala Leu Phe Asp Ile Ile Asp Lys Asp Gln Asn	
115 120 125	
gga gct att tca ctg gat gaa tgg aaa gca tac acc aaa tct gat ggc	432
Gly Ala Ile Ser Leu Asp Glu Trp Lys Ala Tyr Thr Lys Ser Asp Gly	
130 135 140	
atc atc caa tcg tca gaa gat tgc gag gaa aca ttc aga gtg tgc gat	480
Ile Ile Gln Ser Ser Glu Asp Cys Glu Glu Thr Phe Arg Val Cys Asp	
145 150 155 160	
att gat gaa agt gga cag ctc gat gtt gat gag atg aca aga caa cat	528
Ile Asp Glu Ser Gly Gln Leu Asp Val Asp Glu Met Thr Arg Gln His	
165 170 175	
tta gga ttt tgg tac acc atg gat cct gct tgc gaa aag ctc tac ggt	576
Leu Gly Phe Trp Tyr Thr Met Asp Pro Ala Cys Glu Lys Leu Tyr Gly	
180 185 190	

-13/40-

gga gct gtc ccc taa 591
 Gly Ala Val Pro
 195

<210> 7
 <211> 591
 <212> DNA
 <213> Aequoria victoria

<220>
 <221> CDS
 <222> (1)...(588)
 <223> Recombinant Aequorin AEQ2

<400> 7
 atg acc agc gaa caa tac tca gtc aag ctt aca tca gac ttc gac aac 48
 Met Thr Ser Glu Gln Tyr Ser Val Lys Leu Thr Ser Asp Phe Asp Asn
 1 5 10 15

cca aga tgg att gga cga cac aag cat atg ttc aat ttc ctt gat gtc 96
 Pro Arg Trp Ile Gly Arg His Lys His Met Phe Asn Phe Leu Asp Val
 20 25 30

aac cac aat gga aaa atc tct ctt gac gag atg gtc tac aag gca tct 144
 Asn His Asn Gly Lys Ile Ser Leu Asp Glu Met Val Tyr Lys Ala Ser
 35 40 45

gat att gtc atc aat aac ctt gga gca aca cct gag caa gcc aaa cga 192
 Asp Ile Val Ile Asn Asn Leu Gly Ala Thr Pro Glu Gln Ala Lys Arg
 50 55 60

cac aaa gat gct gta gaa gcc ttc ttc gga gga gct gga atg aaa tat 240
 His Lys Asp Ala Val Glu Ala Phe Phe Gly Gly Ala Gly Met Lys Tyr
 65 70 75 80

ggc gtg gaa act gat tgg cct gca tat att gaa gga tgg aaa aaa ttg 288
 Gly Val Glu Thr Asp Trp Pro Ala Tyr Ile Glu Gly Trp Lys Lys Leu
 85 90 95

gct act gat gaa ttg gag aaa tac gcc aaa aac gaa cca acg ctc atc 336
 Ala Thr Asp Glu Leu Glu Lys Tyr Ala Lys Asn Glu Pro Thr Leu Ile
 100 105 110

cgt ata tgg ggt gat gct ttg ttc gat atc gtt gac aaa gat caa aat 384
 Arg Ile Trp Gly Asp Ala Leu Phe Asp Ile Val Asp Lys Asp Gln Asn
 115 120 125

gga gcc att aca ctg gat gaa tgg aaa gca tac acc aaa gct gct ggt 432
 Gly Ala Ile Thr Leu Asp Glu Trp Lys Ala Tyr Thr Lys Ala Ala Gly
 130 135 140

atc atc caa tca tca gaa gat tgc gag gaa aca ttc aga gtg tgc gat 480
 Ile Ile Gln Ser Ser Glu Asp Cys Glu Glu Thr Phe Arg Val Cys Asp
 145 150 155 160

att gat gaa agt gga caa ctc gat gtt gat gag atg aca aga caa cat 528
 Ile Asp Glu Ser Gly Gln Leu Asp Val Asp Glu Met Thr Arg Gln His
 165 170 175

-14/40-

```

tta gga ttt tgg tac acc atg gat cct gct tgc gaa aag ctc tac ggt      576
Leu Gly Phe Trp Tyr Thr Met Asp Pro Ala Cys Glu Lys Leu Tyr Gly
      180                      185                      190

gga gct gtc ccc taa      591
Gly Ala Val Pro
      195

<210> 8
<211> 591
<212> DNA
<213> Aequoria victoria

<220>
<221> CDS
<222> (1) ... (588)
<223> Recombinant Aequorin AEQ3

<400> 8
atg acc agc gaa caa tac tca gtc aag ctt aca tca gac ttc gac aac      48
Met Thr Ser Glu Gln Tyr Ser Val Lys Leu Thr Ser Asp Phe Asp Asn
      1              5              10              15

cca aga tgg att gga cga cac aag cat atg ttc aat ttc ctt gat gtc      96
Pro Arg Trp Ile Gly Arg His Lys His Met Phe Asn Phe Leu Asp Val
      20              25              30

aac cac aat gga aaa atc tct ctt gac gag atg gtc tac aag gca tct      144
Asn His Asn Gly Lys Ile Ser Leu Asp Glu Met Val Tyr Lys Ala Ser
      35              40              45

gat att gtc atc aat aac ctt gga gca aca cct gag caa gcc aaa cga      192
Asp Ile Val Ile Asn Asn Leu Gly Ala Thr Pro Glu Gln Ala Lys Arg
      50              55              60

cac aaa gat gct gta gga gac ttc ttc gga gga gct gga atg aaa tat      240
His Lys Asp Ala Val Gly Asp Phe Phe Gly Gly Ala Gly Met Lys Tyr
      65              70              75              80

ggg gtg gaa act gat tgg cct gca tac att gaa gga tgg aaa aaa ttg      288
Gly Val Glu Thr Asp Trp Pro Ala Tyr Ile Glu Gly Trp Lys Lys Leu
      85              90              95

gct act gat gaa ttg gag aaa tac gcc aaa aac gaa cca acg ctc atc      336
Ala Thr Asp Glu Leu Glu Lys Tyr Ala Lys Asn Glu Pro Thr Leu Ile
      100             105             110

cgt ata tgg ggt gat gct ttg ttc gat atc gtt gac aaa gat caa aat      384
Arg Ile Trp Gly Asp Ala Leu Phe Asp Ile Val Asp Lys Asp Gln Asn
      115             120             125

gga gcc att aca ctg gat gaa tgg aaa gca tac acc aaa gct gct ggt      432
Gly Ala Ile Thr Leu Asp Glu Trp Lys Ala Tyr Thr Lys Ala Ala Gly
      130             135             140

atc atc caa tca tca gaa gat tgc gag gaa aca ttc aga gtg tgc gat      480
Ile Ile Gln Ser Ser Glu Asp Cys Glu Glu Thr Phe Arg Val Cys Asp
      145             150             155             160

att gat gaa aat gga caa ctc gat gtt gat gag atg aca aga caa cat      528

```

-15/40-

Ile Asp Glu Asn Gly Gln Leu Asp Val Asp Glu Met Thr Arg Gln His	
165 170 175	
tta gga ttt tgg tac acc atg gat cct gct tgc gaa aag ctc tac ggt	576
Leu Gly Phe Trp Tyr Thr Met Asp Pro Ala Cys Glu Lys Leu Tyr Gly	
180 185 190	
gga gct gtc ccc taa	591
Gly Ala Val Pro	
195	
<210> 9	
<211> 567	
<212> DNA	
<213> Aequoria victoria	
<220>	
<221> CDS	
<222> (1)...(567)	
<223> Aequorin photoprotein	
<400> 9	
gtc aag ctt aca cca gac ttc gac aac cca aaa tgg att gga cga cac	48
Val Lys Leu Thr Pro Asp Phe Asp Asn Pro Lys Trp Ile Gly Arg His	
1 5 10 15	
aag cac atg ttt aat ttt ctt gat gtc aac cac aat gga agg atc tct	96
Lys His Met Phe Asn Phe Leu Asp Val Asn His Asn Gly Arg Ile Ser	
20 25 30	
ctt gac gag atg gtc tac aag gcg tcc gat att gtt ata aac aat ctt	144
Leu Asp Glu Met Val Tyr Lys Ala Ser Asp Ile Val Ile Asn Asn Leu	
35 40 45	
gga gca aca cct gaa caa gcc aaa cgt cac aaa gat gct gta gaa gcc	192
Gly Ala Thr Pro Glu Gln Ala Lys Arg His Lys Asp Ala Val Glu Ala	
50 55 60	
ttc ttc gga gga gct gca atg aaa tat ggt gta gaa act gaa tgg cct	240
Phe Phe Gly Gly Ala Ala Met Lys Tyr Gly Val Glu Thr Glu Trp Pro	
65 70 75 80	
gaa tac atc gaa gga tgg aaa aga ctg gct tcc gag gaa ttg aaa agg	288
Glu Tyr Ile Glu Gly Trp Lys Arg Leu Ala Ser Glu Glu Leu Lys Arg	
85 90 95	
tat tca aaa aac caa atc aca ctt att cgt tta tgg ggt gat gca ttg	336
Tyr Ser Lys Asn Gln Ile Thr Leu Ile Arg Leu Trp Gly Asp Ala Leu	
100 105 110	
ttc gat atc att gac aaa gac caa aat gga gct att tca ctg gat gaa	384
Phe Asp Ile Ile Asp Lys Asp Gln Asn Gly Ala Ile Ser Leu Asp Glu	
115 120 125	
tgg aaa gca tac acc aaa tct gct ggc atc atc caa tcg tca gaa gat	432
Trp Lys Ala Tyr Thr Lys Ser Ala Gly Ile Ile Gln Ser Ser Glu Asp	
130 135 140	
tgc gag gaa aca ttc aga gtg tgc gat att gat gaa agt gga cag ctc	480
Cys Glu Glu Thr Phe Arg Val Cys Asp Ile Asp Glu Ser Gly Gln Leu	
145 150 155 160	

-16/40-

gat gtt gat gag atg aca aga caa cat tta gga ttt tgg tac acc atg 528
Asp Val Asp Glu Met Thr Arg Gln His Leu Gly Phe Trp Tyr Thr Met
165 170 175

gat cct gct tgc gaa aag ctc tac ggt gga gct gtc ccc 567
Asp Pro Ala Cys Glu Lys Leu Tyr Gly Gly Ala Val Pro
180 185

<210> 10
<211> 588
<212> DNA
<213> *Aequoria victoria*

<220>
<221> CDS
<222> (1)...(588)
<223> Aequorin mutant w/increased bioluminescence
activity

<400> 10
atg acc agc gaa caa tac tca gtc aag ctt aca cca gac ttc gac aac 48
Met Thr Ser Glu Gln Tyr Ser Val Lys Leu Thr Pro Asp Phe Asp Asn
1 5 10 15

cca aaa tgg att gga cga cac aag cac atg ttt aat ttt ctt gat gtc 96
Pro Lys Trp Ile Gly Arg His Lys His Met Phe Asn Phe Leu Asp Val
20 25 30

aac cac aat gga agg atc tct ctt gac gag atg gtc tac aag gcg tcc 144
Asn His Asn Gly Arg Ile Ser Leu Asp Glu Met Val Tyr Lys Ala Ser
35 40 45

gat att gtt ata aac aat ctt gga gca aca cct gaa caa gcc aaa cgt 192
Asp Ile Val Ile Asn Asn Leu Gly Ala Thr Pro Glu Gln Ala Lys Arg
50 55 60

cac aaa gat gct gta gaa gcc ttc ttc gga gga gct gca atg aaa tat 240
His Lys Asp Ala Val Glu Ala Phe Phe Gly Gly Ala Ala Met Lys Tyr
65 70 75 80

ggt gta gaa act gaa tgg cct gaa tac atc gaa gga tgg aaa aga ctg 288
Gly Val Glu Thr Glu Trp Pro Glu Tyr Ile Glu Gly Trp Lys Arg Leu
85 90 95

gct tcc gag gaa ttg aaa agg tat tca aaa aac caa atc aca ctt att 336
Ala Ser Glu Glu Leu Lys Arg Tyr Ser Lys Asn Gln Ile Thr Leu Ile
100 105 110

cgt tta tgg ggt gat gca ttg ttc gat atc att tcc aaa gac caa aat 384
Arg Leu Trp Gly Asp Ala Leu Phe Asp Ile Ile Ser Lys Asp Gln Asn
115 120 125

gga gct att tca ctg gat gaa tgg aaa gca tac acc aaa tct gct ggc 432
Gly Ala Ile Ser Leu Asp Glu Trp Lys Ala Tyr Thr Lys Ser Ala Gly
130 135 140

atc atc caa tcg tca gaa gat tgc gag gaa aca ttc aga gtg tgc gat 480
Ile Ile Gln Ser Ser Glu Asp Cys Glu Glu Thr Phe Arg Val Cys Asp
145 150 155 160

-17/40-

```

att gat gaa agt gga cag ctc gat gtt gat gag atg aca aga caa cat      528
Ile Asp Glu Ser   Gly Gln Leu Asp Val Asp Glu Met Thr Arg Gln His
                165                170                175

tta gga ttt tgg tac acc atg gat cct gct tgc gaa aag ctc tac ggt      576
Leu Gly Phe   Trp Tyr Thr Met Asp Pro Ala Cys Glu Lys Leu Tyr Gly
                180                185                190

gga gct gtc ccc                                          588
Gly Ala Val Pro
                195

<210> 11
<211> 588
<212> DNA
<213> Aequorea victoria

<220>
<221> CDS
<222> (1)...(588)
<223> Recombinant site-directed Aequorin mutant

<400> 11
atg acc agc gaa caa tac tca gtc aag ctt aca cca gac ttc gac aac      48
Met Thr Ser Glu Gln Tyr Ser Val Lys Leu Thr Pro Asp Phe Asp Asn
  1                5                10                15

cca aaa tgg att gga cga cac aag cac atg ttt aat ttt ctt gat gtc      96
Pro Lys Trp   Ile Gly Arg His Lys His Met Phe Asn Phe Leu Asp Val
                20                25                30

aac cac aat gga agg atc tct ctt gac gag atg gtc tac aag gcg tcc      144
Asn His Asn Gly Arg Ile Ser Leu Asp Glu Met Val Tyr Lys Ala Ser
                35                40                45

gat att gtt ata aac aat ctt gga gca aca cct gaa caa gcc aaa cgt      192
Asp Ile Val Ile Asn Asn Leu Gly Ala Thr Pro Glu Gln Ala Lys Arg
                50                55                60

cac aaa gat gct gta gaa gcc ttc ttc gga gga gct gca atg aaa tat      240
His Lys Asp Ala Val Glu Ala Phe Phe Gly Gly Ala Ala Met Lys Tyr
                65                70                75                80

ggg gta gaa act gaa tgg cct gaa tac atc gaa gga tgg aaa aga ctg      288
Gly Val Glu Thr Glu Trp Pro Glu Tyr Ile Glu Gly Trp Lys Arg Leu
                85                90                95

gct tcc gag gaa ttg aaa agg tat tca aaa aac caa atc aca ctt att      336
Ala Ser Glu Glu Leu Lys Arg Tyr Ser Lys Asn Gln Ile Thr Leu Ile
                100                105                110

cgt tta tgg ggt gat gca ttg ttc gat atc att tcc aaa gac caa aat      384
Arg Leu Trp Gly Asp Ala Leu Phe Asp Ile Ile Ser Lys Asp Gln Asn
                115                120                125

gga gct att tca ctg gat tca tgg aaa gca tac acc aaa tct gct ggc      432
Gly Ala Ile Ser Leu Asp Ser Trp Lys Ala Tyr Thr Lys Ser Ala Gly
                130                135                140

```


-18/40-

atc atc caa tcg tca gaa gat tgc gag gaa aca ttc aga gtg tgc gat Ile Ile Gln Ser Ser Glu Asp Cys Glu Glu Thr Phe Arg Val Cys Asp 145 150 155 160	480
att gat gaa agt gga cag ctc gat gtt gat gag atg aca aga caa cat Ile Asp Glu Ser Gly Gln Leu Asp Val Asp Glu Met Thr Arg Gln His 165 170 175	528
tta gga ttt tgg tac acc atg gat cct gct tgc gaa aag ctc tac ggt Leu Gly Phe Trp Tyr Thr Met Asp Pro Ala Cys Glu Lys Leu Tyr Gly 180 185 190	576
gga gct gtc ccc Gly Ala Val Pro 195	588
<210> 12	
<211> 588	
<212> DNA	
<213> Aequorea victoria	
<220>	
<221> CDS	
<222> (1)...(588)	
<223> Aequorin mutant with increased bioluminescence activity	
<400> 12	
atg acc agc gaa caa tac tca gtc aag ctt aca cca gac ttc gac aac Met Thr Ser Glu Gln Tyr Ser Val Lys Leu Thr Pro Asp Phe Asp Asn 1 5 10 15	48
cca aaa tgg att gga cga cac aag cac atg ttt aat ttt ctt gat gtc Pro Lys Trp Ile Gly Arg His Lys His Met Phe Asn Phe Leu Asp Val 20 25 30	96
aac cac aat gga agg atc tct ctt gac gag atg gtc tac aag gcg tcc Asn His Asn Gly Arg Ile Ser Leu Asp Glu Met Val Tyr Lys Ala Ser 35 40 45	144
gat att gtt ata aac aat ctt gga gca aca cct gaa caa gcc aaa cgt Asp Ile Val Ile Asn Asn Leu Gly Ala Thr Pro Glu Gln Ala Lys Arg 50 55 60	192
cac aaa gat gct gta gaa gcc ttc ttc gga gga gct gca atg aaa tat His Lys Asp Ala Val Glu Ala Phe Phe Gly Gly Ala Ala Met Lys Tyr 65 70 75 80	240
ggt gta gaa act gaa tgg cct gaa tac atc gaa gga tgg aaa aga ctg Gly Val Glu Thr Glu Trp Pro Glu Tyr Ile Glu Gly Trp Lys Arg Leu 85 90 95	288
gct tcc gag gaa ttg aaa agg tat tca aaa aac caa atc aca ctt att Ala Ser Glu Glu Leu Lys Arg Tyr Ser Lys Asn Gln Ile Thr Leu Ile 100 105 110	336
cgt tta tgg ggt gat gca ttg ttc gat atc att tcc aaa gac caa aat Arg Leu Trp Gly Asp Ala Leu Phe Asp Ile Ile Ser Lys Asp Gln Asn 115 120 125	384

-19/40-

gca gct att tca ctg gat gaa tgg aaa gca tac acc aaa tct gct ggc Ala Ala Ile Ser Leu Asp Glu Trp Lys Ala Tyr Thr Lys Ser Ala Gly 130 135 140	432
atc atc caa tcg tca gaa gat tgc gag gaa aca ttc aga gtg tgc gat Ile Ile Gln Ser Ser Glu Asp Cys Glu Glu Thr Phe Arg Val Cys Asp 145 150 155 160	480
att gat gaa agt gga cag ctc gat gtt gat gag atg aca aga caa cat Ile Asp Glu Ser Gly Gln Leu Asp Val Asp Glu Met Thr Arg Gln His 165 170 175	528
tta gga ttt tgg tac acc atg gat cct gct tgc gaa aag ctc tac ggt Leu Gly Phe Trp Tyr Thr Met Asp Pro Ala Cys Glu Lys Leu Tyr Gly 180 185 190	576
gga gct gtc ccc Gly Ala Val Pro 195	588
 <210> 13 <211> 567 <212> DNA <213> Aequorea victoria <220> <221> CDS <222> (1)...(567) <223> Recombinant apoaeguorin (AQUALITEp)	
 <400> 13	
gtc aag ctt aca cca gac ttc gac aac cca aaa tgg att gga cga cac Val Lys Leu Thr Pro Asp Phe Asp Asn Pro Lys Trp Ile Gly Arg His 1 5 10 15	48
aag cac atg ttt aat ttt ctt gat gtc aac cac aat gga agg atc tct Lys His Met Phe Asn Phe Leu Asp Val Asn His Asn Gly Arg Ile Ser 20 25 30	96
ctt gac gag atg gtc tac aag gcg tcc gat att gtt ata aac aat ctt Leu Asp Glu Met Val Tyr Lys Ala Ser Asp Ile Val Ile Asn Asn Leu 35 40 45	144
gga gca aca cct gaa caa gcc aaa cgt cac aaa gat gct gta gaa gcc Gly Ala Thr Pro Glu Gln Ala Lys Arg His Lys Asp Ala Val Glu Ala 50 55 60	192
ttc ttc gga gga gct gga atg aaa tat ggt gta gaa act gaa tgg cct Phe Phe Gly Gly Ala Gly Met Lys Tyr Gly Val Glu Thr Glu Trp Pro 65 70 75 80	240
gaa tac atc gaa gga tgg aaa aaa ctg gct tcc gag gaa ttg aaa agg Glu Tyr Ile Glu Gly Trp Lys Lys Leu Ala Ser Glu Glu Leu Lys Arg 85 90 95	288
tat tca aaa aac caa atc aca ctt att cgt tta tgg ggt gat gca ttg Tyr Ser Lys Asn Gln Ile Thr Leu Ile Arg Leu Trp Gly Asp Ala Leu 100 105 110	336
ttc gat atc att gac aaa gac caa aat gga gct att ctg tca gat gaa	384

-20/40-

Phe	Asp	Ile	Ile	Asp	Lys	Asp	Gln	Asn	Gly	Ala	Ile	Leu	Ser	Asp	Glu		
		115					120					125					
tgg	aaa	gca	tac	acc	aaa	tct	gat	ggc	atc	atc	caa	tcg	tca	gaa	gat	432	
Trp	Lys	Ala	Tyr	Thr	Lys	Ser	Asp	Gly	Ile	Ile	Gln	Ser	Ser	Glu	Asp		
	130					135					140						
tgc	gag	gaa	aca	ttc	aga	gtg	tgc	gat	att	gat	gaa	agt	gga	cag	ctc	480	
Cys	Glu	Glu	Thr	Phe	Arg	Val	Cys	Asp	Ile	Asp	Glu	Ser	Gly	Gln	Leu		
145					150					155					160		
gat	gtt	gat	gag	atg	aca	aga	caa	cat	tta	gga	ttt	tgg	tac	acc	atg	528	
Asp	Val	Asp	Glu	Met	Thr	Arg	Gln	His	Leu	Gly	Phe	Trp	Tyr	Thr	Met		
				165					170					175			
gat	cct	gct	tgc	gaa	aag	ctc	tac	ggg	gga	gct	gtc	ccc				567	
Asp	Pro	Ala	Cys	Glu	Lys	Leu	Tyr	Gly	Gly	Ala	Val	Pro					
			180					185									

<210> 14
 <211> 236
 <212> PRT
 <213> Vibrio fisheri

<400> 14

Met	Pro	Ile	Asn	Cys	Lys	Val	Lys	Ser	Ile	Glu	Pro	Leu	Ala	Cys	Asn
1				5					10					15	
Thr	Phe	Arg	Ile	Leu	Leu	His	Pro	Glu	Gln	Pro	Val	Ala	Phe	Lys	Ala
			20					25					30		
Gly	Gln	Tyr	Leu	Thr	Val	Val	Met	Gly	Glu	Lys	Asp	Lys	Arg	Pro	Phe
		35					40					45			
Ser	Ile	Ala	Ser	Ser	Pro	Cys	Arg	His	Glu	Gly	Glu	Ile	Glu	Leu	His
	50					55					60				
Ile	Gly	Ala	Ala	Glu	His	Asn	Ala	Tyr	Ala	Gly	Glu	Val	Val	Glu	Ser
65					70					75				80	
Met	Lys	Ser	Ala	Leu	Glu	Thr	Gly	Gly	Asp	Ile	Leu	Ile	Asp	Ala	Pro
				85					90					95	
His	Gly	Glu	Ala	Trp	Ile	Arg	Glu	Asp	Ser	Asp	Arg	Ser	Met	Leu	Leu
			100					105					110		
Ile	Ala	Gly	Gly	Thr	Gly	Phe	Ser	Tyr	Val	Arg	Ser	Ile	Leu	Asp	His
		115					120					125			
Cys	Ile	Ser	Gln	Gln	Ile	Gln	Lys	Pro	Ile	Tyr	Leu	Tyr	Trp	Gly	Gly
	130					135					140				
Arg	Asp	Glu	Cys	Gln	Leu	Tyr	Ala	Lys	Ala	Glu	Leu	Glu	Ser	Ile	Ala
145					150					155					160
Gln	Ala	His	Ser	His	Ile	Thr	Phe	Val	Pro	Val	Val	Glu	Lys	Ser	Glu
				165					170					175	
Gly	Trp	Thr	Gly	Lys	Thr	Gly	Asn	Val	Leu	Glu	Ala	Val	Lys	Ala	Asp
			180				185						190		
Phe	Asn	Ser	Leu	Ala	Asp	Met	Asp	Ile	Tyr	Ile	Ala	Gly	Arg	Phe	Glu
		195					200					205			
Met	Ala	Gly	Ala	Ala	Arg	Glu	Gln	Phe	Thr	Thr	Glu	Lys	Gln	Ala	Lys
	210					215					220				
Lys	Glu	Gln	Leu	Phe	Gly	Asp	Ala	Phe	Ala	Phe	Ile				
225					230						235				

<210> 15
 <211> 1079
 <212> DNA

-21/40-

<213> Renilla mulleri

<220>

<221> CDS

<222> (259)...(975)

<223> Renilla mulleri GFP

<400> 15

ggttatacac aagtgtatcg cgtatctgca gacgcattcta gtgggattat tcgagcggta	60
gtattttacgt cagacctgtc taatcgaaac cacaacaaac tcttaaaata agccacattt	120
acataatatac taagagacgc ctcattttaag agtagtaaaa atataatata tgatagagta	180
tacaactctc gccttagaca gacagtgtgc aacagagtaa ctcttggttaa tgcaatcgaa	240
agcgtcaaga gagataag atg agt aaa caa ata ttg aag aac act tgt tta	291
Met Ser Lys Gln Ile Leu Lys Asn Thr Cys Leu	
1 5 10	
caa gaa gta atg tcg tat aaa gta aat ctg gaa gga att gta aac aac	339
Gln Glu Val Met Ser Tyr Lys Val Asn Leu Glu Gly Ile Val Asn Asn	
15 20 25	
cat gtt ttt aca atg gag ggt tgc ggc aaa ggg aat att tta ttc ggc	387
His Val Phe Thr Met Glu Gly Cys Gly Lys Gly Asn Ile Leu Phe Gly	
30 35 40	
aat caa ctg gtt cag att cgt gtc acg aaa ggg gcc cca ctg cct ttt	435
Asn Gln Leu Val Gln Ile Arg Val Thr Lys Gly Ala Pro Leu Pro Phe	
45 50 55	
gca ttt gat att gtg tca cca gct ttt caa tat ggc aac cgt act ttc	483
Ala Phe Asp Ile Val Ser Pro Ala Phe Gln Tyr Gly Asn Arg Thr Phe	
60 65 70 75	
acg aaa tat ccg aat gat ata tca gat tat ttt ata caa tca ttt cca	531
Thr Lys Tyr Pro Asn Asp Ile Ser Asp Tyr Phe Ile Gln Ser Phe Pro	
80 85 90	
gca gga ttt atg tat gaa cga aca tta cgt tac gaa gat ggc gga ctt	579
Ala Gly Phe Met Tyr Glu Arg Thr Leu Arg Tyr Glu Asp Gly Gly Leu	
95 100 105	
gtt gaa att cgt tca gat ata aat tta ata gaa gac aag ttc gtc tac	627
Val Glu Ile Arg Ser Asp Ile Asn Leu Ile Glu Asp Lys Phe Val Tyr	
110 115 120	
aga gtg gaa tac aaa ggt agt aac ttc cca gat gat ggt ccc gtc atg	675
Arg Val Glu Tyr Lys Gly Ser Asn Phe Pro Asp Asp Gly Pro Val Met	
125 130 135	
cag aag act atc tta gga ata gag cct tca ttt gaa gcc atg tac atg	723
Gln Lys Thr Ile Leu Gly Ile Glu Pro Ser Phe Glu Ala Met Tyr Met	
140 145 150 155	
aat aat ggc gtc ttg gtc ggc gaa gta att ctt gtc tat aaa cta aac	771
Asn Asn Gly Val Leu Val Gly Glu Val Ile Leu Val Tyr Lys Leu Asn	
160 165 170	
tct ggg aaa tat tat tca tgt cac atg aaa aca tta atg aag tcg aaa	819
Ser Gly Lys Tyr Tyr Ser Cys His Met Lys Thr Leu Met Lys Ser Lys	
175 180 185	
ggg gta gta aag gag ttt cct tcg tat cat ttt att caa cat cgt ttg	867

-22/40-

Gly Val Val Lys Glu Phe Pro Ser Tyr His Phe Ile Gln His Arg Leu
 190 195 200

gaa aag act tac gta gaa gac ggg ggg ttc gtt gaa cag cat gag act 915
 Glu Lys Thr Tyr Val Glu Asp Gly Gly Phe Val Glu Gln His Glu Thr
 205 210 215

gct att gct caa atg aca tct ata gga aaa cca cta gga tcc tta cac 963
 Ala Ile Ala Gln Met Thr Ser Ile Gly Lys Pro Leu Gly Ser Leu His
 220 225 230 235

gaa tgg gtt taa acacagttac attactttttt ccaattcgtg tttcatgtca 1015
 Glu Trp Val *

aataataatt ttttaaacaa ttatcaatgt tttgtgatat gtttgtaaaa aaaaaaaaaa 1075
 aaaa 1079

<210> 16
 <211> 238
 <212> PRT
 <213> Renilla mulleri

<400> 16
 Met Ser Lys Gln Ile Leu Lys Asn Thr Cys Leu Gln Glu Val Met Ser
 1 5 10 15
 Tyr Lys Val Asn Leu Glu Gly Ile Val Asn Asn His Val Phe Thr Met
 20 25 30
 Glu Gly Cys Gly Lys Gly Asn Ile Leu Phe Gly Asn Gln Leu Val Gln
 35 40 45
 Ile Arg Val Thr Lys Gly Ala Pro Leu Pro Phe Ala Phe Asp Ile Val
 50 55 60
 Ser Pro Ala Phe Gln Tyr Gly Asn Arg Thr Phe Thr Lys Tyr Pro Asn
 65 70 75 80
 Asp Ile Ser Asp Tyr Phe Ile Gln Ser Phe Pro Ala Gly Phe Met Tyr
 85 90 95
 Glu Arg Thr Leu Arg Tyr Glu Asp Gly Gly Leu Val Glu Ile Arg Ser
 100 105 110
 Asp Ile Asn Leu Ile Glu Asp Lys Phe Val Tyr Arg Val Glu Tyr Lys
 115 120 125
 Gly Ser Asn Phe Pro Asp Asp Gly Pro Val Met Gln Lys Thr Ile Leu
 130 135 140
 Gly Ile Glu Pro Ser Phe Glu Ala Met Tyr Met Asn Asn Gly Val Leu
 145 150 155 160
 Val Gly Glu Val Ile Leu Val Tyr Lys Leu Asn Ser Gly Lys Tyr Tyr
 165 170 175
 Ser Cys His Met Lys Thr Leu Met Lys Ser Lys Gly Val Val Lys Glu
 180 185 190
 Phe Pro Ser Tyr His Phe Ile Gln His Arg Leu Glu Lys Thr Tyr Val
 195 200 205
 Glu Asp Gly Gly Phe Val Glu Gln His Glu Thr Ala Ile Ala Gln Met
 210 215 220
 Thr Ser Ile Gly Lys Pro Leu Gly Ser Leu His Glu Trp Val
 225 230 235

<210> 17
 <211> 1217
 <212> DNA
 <213> Renilla mulleri

<220>

-23/40-

<221> CDS

<222> (31)...(963)

<223> Renilla mulleri luciferase

<400> 17

```

cggcacgagg tttaagaatc aataaaaaaa atg acg tca aaa gtt tac gat cct      54
                               Met Thr Ser Lys Val Tyr Asp Pro
                               1                               5

gaa tta aga aaa cgc atg att act ggt cca caa tgg tgg gca aga tgt      102
Glu Leu Arg Lys Arg Met Ile Thr Gly Pro Gln Trp Trp Ala Arg Cys
   10                               15                               20

aaa caa atg aat gtt ctt gat tca ttt att aat tat tat gat tca gaa      150
Lys Gln Met Asn Val Leu Asp Ser Phe Ile Asn Tyr Tyr Asp Ser Glu
   25                               30                               35                               40

aaa cat gca gaa aat gca gtt ata ttt tta cat ggt aat gca gca tct      198
Lys His Ala Glu Asn Ala Val Ile Phe Leu His Gly Asn Ala Ala Ser
                               45                               50                               55

tct tat tta tgg cgt cat gtt gta cca cat gtt gaa cca gtg gcg cga      246
Ser Tyr Leu Trp Arg His Val Val Pro His Val Glu Pro Val Ala Arg
                               60                               65                               70

tgt att ata cca gat ctt ata ggt atg ggt aaa tca ggc aag tct ggt      294
Cys Ile Ile Pro Asp Leu Ile Gly Met Gly Lys Ser Gly Lys Ser Gly
   75                               80                               85

aat ggt tcc tat aga tta cta gat cat tac aaa tat ctt act gaa tgg      342
Asn Gly Ser Tyr Arg Leu Leu Asp His Tyr Lys Tyr Leu Thr Glu Trp
   90                               95                               100

ttc aaa cat ctt aat tta cca aag aag atc att ttt gtc ggt cat gat      390
Phe Lys His Leu Asn Leu Pro Lys Lys Ile Ile Phe Val Gly His Asp
  105                               110                               115                               120

tgg ggt gct tgt tta gca ttt cat tat tgc tat gaa cat cag gat cgc      438
Trp Gly Ala Cys Leu Ala Phe His Tyr Cys Tyr Glu His Gln Asp Arg
                               125                               130                               135

atc aaa gca gtt gtt cat gct gaa agt gta gta gat gtg att gaa tcg      486
Ile Lys Ala Val Val His Ala Glu Ser Val Val Asp Val Ile Glu Ser
                               140                               145                               150

tgg gac gaa tgg cct gat att gaa gaa gat att gct ttg att aaa tct      534
Trp Asp Glu Trp Pro Asp Ile Glu Glu Asp Ile Ala Leu Ile Lys Ser
                               155                               160                               165

gaa gaa gga gaa aaa atg gtt tta gag aat aac ttc ttc gtg gaa acc      582
Glu Glu Gly Glu Lys Met Val Leu Glu Asn Asn Phe Phe Val Glu Thr
   170                               175                               180

atg ttg cca tca aaa atc atg aga aag ttg gaa cca gag gaa ttt gct      630
Met Leu Pro Ser Lys Ile Met Arg Lys Leu Glu Pro Glu Glu Phe Ala
  185                               190                               195                               200

gct tat ctt gaa cca ttt aaa gag aaa ggt gaa gtt cgt cgt cca aca      678
Ala Tyr Leu Glu Pro Phe Lys Glu Lys Gly Glu Val Arg Arg Pro Thr
                               205                               210                               215

```

-24/40-

tta tca tgg cct cgt gaa atc cct ttg gta aaa ggt ggt aaa ccg gat	726
Leu Ser Trp Pro Arg Glu Ile Pro Leu Val Lys Gly Gly Lys Pro Asp	
220 225 230	
gta gta gaa att gtc agg aat tat aat gct tat ctt cgt gca agt cat	774
Val Val Glu Ile Val Arg Asn Tyr Asn Ala Tyr Leu Arg Ala Ser His	
235 240 245	
gat tta cca aaa atg ttt att gaa tct gat cca gga ttc ttt tcc aat	822
Asp Leu Pro Lys Met Phe Ile Glu Ser Asp Pro Gly Phe Phe Ser Asn	
250 255 260	
gct att gtt gaa ggt gct aag aaa ttc cct aat act gaa ttt gtt aaa	870
Ala Ile Val Glu Gly Ala Lys Lys Phe Pro Asn Thr Glu Phe Val Lys	
265 270 275 280	
gtc aaa ggt ctt cat ttt tca caa gaa gat gca cct gat gaa atg gga	918
Val Lys Gly Leu His Phe Ser Gln Glu Asp Ala Pro Asp Glu Met Gly	
285 290 295	
aat tat ata aaa tcg ttt gtt gag cgt gtt ctt aaa aat gaa caa	963
Asn Tyr Ile Lys Ser Phe Val Glu Arg Val Leu Lys Asn Glu Gln	
300 305 310	
taaactacca gggtttccatg ttgccacttt agctggggttt aataaatttc actatcaatt	1023
tgaacaattt cacattaatt ttaactatta aaaaattatg gacacaggga ttatatcaga	1083
tgattaattt agttgggaac aatgaatacc gaataattatg aattctcttt agctatttat	1143
aataatcaca ttcttatgta ataaaacttt gttttaataa attaattgatt cagaaaaaaa	1203
aaaaaaaaaaaa aaaa	1217

<210> 18

<211> 311

<212> PRT

<213> Renilla mulleri

<400> 18

Met Thr Ser Lys Val Tyr Asp Pro Glu Leu Arg Lys Arg Met Ile Thr	
1 5 10 15	
Gly Pro Gln Trp Trp Ala Arg Cys Lys Gln Met Asn Val Leu Asp Ser	
20 25 30	
Phe Ile Asn Tyr Tyr Asp Ser Glu Lys His Ala Glu Asn Ala Val Ile	
35 40 45	
Phe Leu His Gly Asn Ala Ala Ser Ser Tyr Leu Trp Arg His Val Val	
50 55 60	
Pro His Val Glu Pro Val Ala Arg Cys Ile Ile Pro Asp Leu Ile Gly	
65 70 75 80	
Met Gly Lys Ser Gly Lys Ser Gly Asn Gly Ser Tyr Arg Leu Leu Asp	
85 90 95	
His Tyr Lys Tyr Leu Thr Glu Trp Phe Lys His Leu Asn Leu Pro Lys	
100 105 110	
Lys Ile Ile Phe Val Gly His Asp Trp Gly Ala Cys Leu Ala Phe His	
115 120 125	
Tyr Cys Tyr Glu His Gln Asp Arg Ile Lys Ala Val Val His Ala Glu	
130 135 140	
Ser Val Val Asp Val Ile Glu Ser Trp Asp Glu Trp Pro Asp Ile Glu	
145 150 155 160	
Glu Asp Ile Ala Leu Ile Lys Ser Glu Glu Gly Glu Lys Met Val Leu	
165 170 175	
Glu Asn Asn Phe Phe Val Glu Thr Met Leu Pro Ser Lys Ile Met Arg	
180 185 190	
Lys Leu Glu Pro Glu Glu Phe Ala Ala Tyr Leu Glu Pro Phe Lys Glu	

-26/40-

gac cta tgt gta gac tgc aca act gga tgc ctc aaa ggt ctt gcc aat 486
 Asp Leu Cys Val Asp Cys Thr Thr Gly Cys Leu Lys Gly Leu Ala Asn
 135 140 145 150

gtg caa tgt tct gat tta ctc aag aaa tgg ctg cca caa aga tgt gca 534
 Val Gln Cys Ser Asp Leu Leu Lys Lys Trp Leu Pro Gln Arg Cys Ala
 155 160 165

act ttt gct agc aaa att caa ggc caa gtg gac aaa ata aag ggt gcc 582
 Thr Phe Ala Ser Lys Ile Gln Gly Gln Val Asp Lys Ile Lys Gly Ala
 170 175 180

ggt ggt gat taa tcctaataaga atactgcata actggatgat gatatactag 634
 Gly Gly Asp *
 185

cttattgctc ataaaatggc catttttttgt aacaaatcga gtctatgtaa ttcaaaatc 694
 ctaattaatt gttaatacat atgtaattcc tataaatata atttatgcaa tccaaaaaaa 754
 aaaaaaaaa a 765

<210> 20

<211> 185

<212> PRT

<213> Renilla mulleri

<400> 20

Met Gly Val Lys Val Leu Phe Ala Leu Ile Cys Ile Ala Val Ala Glu
 1 5 10 15
 Ala Lys Pro Thr Glu Asn Asn Glu Asp Phe Asn Ile Val Ala Val Ala
 20 25 30
 Ser Asn Phe Ala Thr Thr Asp Leu Asp Ala Asp Arg Gly Lys Leu Pro
 35 40 45
 Gly Lys Lys Leu Pro Leu Glu Val Leu Lys Glu Met Glu Ala Asn Ala
 50 55 60
 Arg Lys Ala Gly Cys Thr Arg Gly Cys Leu Ile Cys Leu Ser His Ile
 65 70 75 80
 Lys Cys Thr Pro Lys Met Lys Lys Phe Ile Pro Gly Arg Cys His Thr
 85 90 95
 Tyr Glu Gly Asp Lys Glu Ser Ala Gln Gly Gly Ile Gly Glu Ala Ile
 100 105 110
 Val Asp Ile Pro Glu Ile Pro Gly Phe Lys Asp Leu Glu Pro Met Glu
 115 120 125
 Gln Phe Ile Ala Gln Val Asp Leu Cys Val Asp Cys Thr Thr Gly Cys
 130 135 140
 Leu Lys Gly Leu Ala Asn Val Gln Cys Ser Asp Leu Leu Lys Lys Trp
 145 150 155 160
 Leu Pro Gln Arg Cys Ala Thr Phe Ala Ser Lys Ile Gln Gly Gln Val
 165 170 175
 Asp Lys Ile Lys Gly Ala Gly Gly Asp
 180 185

<210> 21

<211> 1146

<212> DNA

<213> Gaussia

<220>

<221> CDS

<222> (1)...(1146)

<223> Nucleotide sequence encoding a CBD-Gaussia

-27/40-

luciferase fusion protein

```

<400> 21
atg tca gtt gaa ttt tac aac tct aac aaa tca gca caa aca aac tca      48
Met Ser Val Glu Phe Tyr Asn Ser Asn Lys Ser Ala Gln Thr Asn Ser
1          5          10          15

att aca cca ata atc aaa att act aac aca tct gac agt gat tta aat      96
Ile Thr Pro Ile Ile Lys Ile Thr Asn Thr Ser Asp Ser Asp Leu Asn
          20          25          30

tta aat gac gta aaa gtt aga tat tat tac aca agt gat ggt aca caa      144
Leu Asn Asp Val Lys Val Arg Tyr Tyr Tyr Thr Ser Asp Gly Thr Gln
          35          40          45

gga caa act ttc tgg tgt gac cat gct ggt gca tta tta gga aat agc      192
Gly Gln Thr Phe Trp Cys Asp His Ala Gly Ala Leu Leu Gly Asn Ser
          50          55          60

tat gtt gat aac act agc aaa gtg aca gca aac ttc gtt aaa gaa aca      240
Tyr Val Asp Asn Thr Ser Lys Val Thr Ala Asn Phe Val Lys Glu Thr
          65          70          75          80

gca agc cca aca tca acc tat gat aca tat gtt gaa ttt gga ttt gca      288
Ala Ser Pro Thr Ser Thr Tyr Asp Thr Tyr Val Glu Phe Gly Phe Ala
          85          90          95

agc gga gca gct act ctt aaa aaa gga caa ttt ata act att caa gga      336
Ser Gly Ala Ala Thr Leu Lys Lys Gly Gln Phe Ile Thr Ile Gln Gly
          100          105          110

aga ata aca aaa tca gac tgg tca aac tac act caa aca aat gac tat      384
Arg Ile Thr Lys Ser Asp Trp Ser Asn Tyr Thr Gln Thr Asn Asp Tyr
          115          120          125

tca ttt gat gca agt agt tca aca cca gtt gta aat cca aaa gtt aca      432
Ser Phe Asp Ala Ser Ser Thr Pro Val Val Asn Pro Lys Val Thr
          130          135          140

gga tat ata ggt gga gct aaa gtt ctt ggt aca gca cca ggt tcc gcg      480
Gly Tyr Ile Gly Gly Ala Lys Val Leu Gly Thr Ala Pro Gly Ser Ala
          145          150          155          160

ggt ctg gtg cca cgc ggt agt act gca att ggt atg aaa gaa acc gct      528
Gly Leu Val Pro Arg Gly Ser Thr Ala Ile Gly Met Lys Glu Thr Ala
          165          170          175

gct gct aaa ttc gaa cgc cag cac atg gac agc cca gat ctg ggt acc      576
Ala Ala Lys Phe Glu Arg Gln His Met Asp Ser Pro Asp Leu Gly Thr
          180          185          190

gat gac gac gac aag atg gga gtg aaa gtt ctt ttt gcc ctt att tgt      624
Asp Asp Asp Asp Lys Met Gly Val Lys Val Leu Phe Ala Leu Ile Cys
          195          200          205

att gct gtg gcc gag gcc aaa cca act gaa aac aat gaa gat ttc aac      672
Ile Ala Val Ala Glu Ala Lys Pro Thr Glu Asn Asn Glu Asp Phe Asn
          210          215          220

att gta gct gta gct agc aac ttt gct aca acg gat ctc gat gct gac      720
Ile Val Ala Val Ala Ser Asn Phe Ala Thr Thr Asp Leu Asp Ala Asp

```

-28/40-

225	230	235	240	
cgt ggt aaa ttg	ccc gga aaa aaa tta	cca ctt gag gta	ctc aaa gaa	768
Arg Gly Lys Leu	Pro Gly Lys Lys Leu	Pro Leu Glu Val	Leu Lys Glu	
	245	250	255	
atg gaa gcc aat	gct agg aaa gct	ggc tgc act agg	gga tgt ctg ata	816
Met Glu Ala Asn	Ala Arg Lys Ala	Gly Cys Thr Arg	Gly Cys Leu Ile	
	260	265	270	
tgc ctg tca cac	atc aag tgt aca	ccc aaa atg aag	ttt atc cca	864
Cys Leu Ser His	Ile Lys Cys Thr	Pro Lys Met Lys	Phe Ile Pro	
	275	280	285	
gga aga tgc cac	acc tat gaa gga	gac aaa gaa agt	gca cag gga gga	912
Gly Arg Cys His	Thr Tyr Gln Gly	Asp Lys Glu Ser	Ala Gln Gly Gly	
	290	295	300	
ata gga gag gct	att gtt gac att	cct gaa att cct	ggg ttt aag gat	960
Ile Gly Glu Ala	Ile Val Asp Ile	Pro Glu Ile Pro	Gly Phe Lys Asp	
	310	315	320	
ttg gaa ccc atg	gaa caa ttc att	gca caa gtt gac	cta tgt gta gac	1008
Leu Glu Pro Met	Glu Gln Phe Ile	Ala Gln Val Asp	Leu Cys Val Asp	
	325	330	335	
tgc aca act gga	tgc ctc aaa ggt	ctt gcc aat gtg	caa tgt tct gat	1056
Cys Thr Thr Gly	Cys Leu Lys Gly	Leu Ala Asn Val	Gln Cys Ser Asp	
	340	345	350	
tta ctc aag aaa	tgg ctg cca caa	aga tgt gca act	ttt gct agc aaa	1104
Leu Leu Lys Lys	Trp Leu Pro Gln	Arg Cys Ala Thr	Phe Ala Ser Lys	
	355	360	365	
att caa ggc caa	gtg gac aaa ata	aag ggt gcc ggt	ggt gat	1146
Ile Gln Gly Gln	Val Asp Lys Ile	Lys Gly Ala Gly	Gly Asp	
	370	375	380	

<210> 22
 <211> 382
 <212> PRT
 <213> Gaussia

<400> 22
 Met Ser Val Glu Phe Tyr Asn Ser Asn Lys Ser Ala Gln Thr Asn Ser
 1 5 10 15
 Ile Thr Pro Ile Ile Lys Ile Thr Asn Thr Ser Asp Ser Asp Leu Asn
 20 25 30
 Leu Asn Asp Val Lys Val Arg Tyr Tyr Tyr Thr Ser Asp Gly Thr Gln
 35 40 45
 Gly Gln Thr Phe Trp Cys Asp His Ala Gly Ala Leu Leu Gly Asn Ser
 50 55 60
 Tyr Val Asp Asn Thr Ser Lys Val Thr Ala Asn Phe Val Lys Glu Thr
 65 70 75 80
 Ala Ser Pro Thr Ser Thr Tyr Asp Thr Tyr Val Glu Phe Gly Phe Ala
 85 90 95
 Ser Gly Ala Ala Thr Leu Lys Lys Gly Gln Phe Ile Thr Ile Gln Gly
 100 105 110
 Arg Ile Thr Lys Ser Asp Trp Ser Asn Tyr Thr Gln Thr Asn Asp Tyr
 115 120 125

-29/40-

Ser Phe Asp Ala Ser Ser Ser Thr Pro Val Val Asn Pro Lys Val Thr
 130 135 140
 Gly Tyr Ile Gly Gly Ala Lys Val Leu Gly Thr Ala Pro Gly Ser Ala
 145 150 155 160
 Gly Leu Val Pro Arg Gly Ser Thr Ala Ile Gly Met Lys Glu Thr Ala
 165 170 175
 Ala Ala Lys Phe Glu Arg Gln His Met Asp Ser Pro Asp Leu Gly Thr
 180 185 190
 Asp Asp Asp Asp Lys Met Gly Val Lys Val Leu Phe Ala Leu Ile Cys
 195 200 205
 Ile Ala Val Ala Glu Ala Lys Pro Thr Glu Asn Asn Glu Asp Phe Asn
 210 215 220
 Ile Val Ala Val Ala Ser Asn Phe Ala Thr Thr Asp Leu Asp Ala Asp
 225 230 235 240
 Arg Gly Lys Leu Pro Gly Lys Lys Leu Pro Leu Glu Val Leu Lys Glu
 245 250 255
 Met Glu Ala Asn Ala Arg Lys Ala Gly Cys Thr Arg Gly Cys Leu Ile
 260 265 270
 Cys Leu Ser His Ile Lys Cys Thr Pro Lys Met Lys Lys Phe Ile Pro
 275 280 285
 Gly Arg Cys His Thr Tyr Glu Gly Asp Lys Glu Ser Ala Gln Gly Gly
 290 295 300
 Ile Gly Glu Ala Ile Val Asp Ile Pro Glu Ile Pro Gly Phe Lys Asp
 305 310 315 320
 Leu Glu Pro Met Glu Gln Phe Ile Ala Gln Val Asp Leu Cys Val Asp
 325 330 335
 Cys Thr Thr Gly Cys Leu Lys Gly Leu Ala Asn Val Gln Cys Ser Asp
 340 345 350
 Leu Leu Lys Lys Trp Leu Pro Gln Arg Cys Ala Thr Phe Ala Ser Lys
 355 360 365
 Ile Gln Gly Gln Val Asp Lys Ile Lys Gly Ala Gly Gly Asp
 370 375 380

<210> 23

<211> 864

<212> DNA

<213> Renilla renifomis

<220>

<221> CDS

<222> (61)...(762)

<223> GFP Clone-1

<400> 23

ggcaccgaggg tttcctgaca caataaaaac ctttcaaatt gtttctctgt agcagtaagt 60
 atg gat ctc gca aaa ctt ggt ttg aag gaa gtg atg cct act aaa atc 108
 Met Asp Leu Ala Lys Leu Gly Leu Lys Glu Val Met Pro Thr Lys Ile
 1 5 10 15
 aac tta gaa gga ctg gtt ggc gac cac gct ttc tca atg gaa gga gtt 156
 Asn Leu Glu Gly Leu Val Gly Asp His Ala Phe Ser Met Glu Gly Val
 20 25 30
 ggc gaa ggc aac ata ttg gaa gga act caa gag gtg aag ata tcg gta 204
 Gly Glu Gly Asn Ile Leu Glu Gly Thr Gln Glu Val Lys Ile Ser Val
 35 40 45
 aca aaa ggc gca cca ctc cca ttc gca ttt gat atc gta tct gtg gct 252
 Thr Lys Gly Ala Pro Leu Pro Phe Ala Phe Asp Ile Val Ser Val Ala
 50 55 60

-30/40-

ttt tca tat ggg aac aga gct tat acc ggt tac cca gaa gaa att tcc	300
Phe Ser Tyr Gly Asn Arg Ala Tyr Thr Gly Tyr Pro Glu Glu Ile Ser	
65 70 75 80	
gac tac ttc ctc cag tcg ttt cca gaa ggc ttt act tac gag aga aac	348
Asp Tyr Phe Leu Gln Ser Phe Pro Glu Gly Phe Thr Tyr Glu Arg Asn	
85 90 95	
att cgt tat caa gat gga gga act gca att gtt aaa tct gat ata agc	396
Ile Arg Tyr Gln Asp Gly Gly Thr Ala Ile Val Lys Ser Asp Ile Ser	
100 105 110	
ttg gaa gat ggt aaa ttc ata gtg aat gta gac ttc aaa gcg aag gat	444
Leu Glu Asp Gly Lys Phe Ile Val Asn Val Asp Phe Lys Ala Lys Asp	
115 120 125	
cta cgt cgc atg gga cca gtc atg cag caa gac atc gtg ggt atg cag	492
Leu Arg Arg Met Gly Pro Val Met Gln Gln Asp Ile Val Gly Met Gln	
130 135 140	
cca tcg tat gag tca atg tac acc aat gtc act tca gtt ata ggg gaa	540
Pro Ser Tyr Glu Ser Met Tyr Thr Asn Val Thr Ser Val Ile Gly Glu	
145 150 155 160	
tgt ata ata gca ttc aaa ctt caa act ggc aag cat ttc act tac cac	588
Cys Ile Ile Ala Phe Lys Leu Gln Thr Gly Lys His Phe Thr Tyr His	
165 170 175	
atg agg aca gtt tac aaa tca aag aag cca gtg gaa act atg cca ttg	636
Met Arg Thr Val Tyr Lys Ser Lys Lys Pro Val Glu Thr Met Pro Leu	
180 185 190	
tat cat ttc atc cag cat cgc ctc gtt aag acc aat gtg gac aca gcc	684
Tyr His Phe Ile Gln His Arg Leu Val Lys Thr Asn Val Asp Thr Ala	
195 200 205	
agt ggt tac gtt gtg caa cac gag aca gca att gca gcg cat tct aca	732
Ser Gly Tyr Val Val Gln His Glu Thr Ala Ile Ala Ala His Ser Thr	
210 215 220	
atc aaa aaa att gaa ggc tct tta cca tag atacctgtac acaattattc	782
Ile Lys Lys Ile Glu Gly Ser Leu Pro *	
225 230	
tatgcacgta gcattttttt ggaaatataa gtgggtattgt tcaataaaaat attaaatata	842
aaaaaaaaa aaaaaaaaaa aa	864
<210> 24	
<211> 860	
<212> DNA	
<213> Renilla reniformis	
<220>	
<221> CDS	
<222> (57)...(758)	
<223> GFP Clone-2	
<400> 24	
ggcacgaggc tgacacaata aaaaaccttt caaattgttt ctctgtagca ggaagt atg	59
Met	

-31/40-

1

gat ctc gca aaa ctt ggt ttg aag gaa gtg atg cct act aaa atc aac	107
Asp Leu Ala Lys Leu Gly Leu Lys Glu Val Met Pro Thr Lys Ile Asn	
5 10 15	
tta gaa gga ctg gtt ggc gac cac gct ttc tca atg gaa gga gtt ggc	155
Leu Glu Gly Leu Val Gly Asp His Ala Phe Ser Met Glu Gly Val Gly	
20 25 30	
gaa ggc aac ata ttg gaa gga act caa gag gtg aag ata tcg gta aca	203
Glu Gly Asn Ile Leu Glu Gly Thr Gln Glu Val Lys Ile Ser Val Thr	
35 40 45	
aaa ggc gca cca ctc cca ttc gca ttt gat atc gta tct gtt gct ttc	251
Lys Gly Ala Pro Leu Pro Phe Ala Phe Asp Ile Val Ser Val Ala Phe	
50 55 60 65	
tca tat ggg aac aga gct tat act ggt tac cca gaa gaa att tcc gac	299
Ser Tyr Gly Asn Arg Ala Tyr Thr Gly Tyr Pro Glu Glu Ile Ser Asp	
70 75 80	
tac ttc ctc cag tcg ttt cca gaa ggc ttt act tac gag aga aac att	347
Tyr Phe Leu Gln Ser Phe Pro Glu Gly Phe Thr Tyr Glu Arg Asn Ile	
85 90 95	
cgt tat caa gat gga gga act gca att gtt aaa tct gat ata agc ttg	395
Arg Tyr Gln Asp Gly Gly Thr Ala Ile Val Lys Ser Asp Ile Ser Leu	
100 105 110	
gaa gat ggt aaa ttc ata gtg aat gta gac ttc aaa gcg aag gat cta	443
Glu Asp Gly Lys Phe Ile Val Asn Val Asp Phe Lys Ala Lys Asp Leu	
115 120 125	
cgt cgc atg gga cca gtc atg cag caa gac atc gtg ggt atg cag cca	491
Arg Arg Met Gly Pro Val Met Gln Gln Asp Ile Val Gly Met Gln Pro	
130 135 140 145	
tcg tat gag tca atg tac acc aat gtc act tca gtt ata ggg gaa tgt	539
Ser Tyr Glu Ser Met Tyr Thr Asn Val Thr Ser Val Ile Gly Glu Cys	
150 155 160	
ata ata gca ttc aaa ctt caa act ggc aaa cat ttc act tac cac atg	587
Ile Ile Ala Phe Lys Leu Gln Thr Gly Lys His Phe Thr Tyr His Met	
165 170 175	
agg aca gtt tac aaa tca aag aag cca gtg gaa act atg cca ttg tat	635
Arg Thr Val Tyr Lys Ser Lys Lys Pro Val Glu Thr Met Pro Leu Tyr	
180 185 190	
cat ttc atc cag cat cgc ctc gtt aag acc aat gtg gac aca gcc agt	683
His Phe Ile Gln His Arg Leu Val Lys Thr Asn Val Asp Thr Ala Ser	
195 200 205	
ggt tac gtt gtg caa cac gag aca gca att gca gcg cat tct aca atc	731
Gly Tyr Val Val Gln His Glu Thr Ala Ile Ala Ala His Ser Thr Ile	
210 215 220 225	
aaa aaa att gaa ggc tct tta cca tag atatctatac acaattattc	778
Lys Lys Ile Glu Gly Ser Leu Pro *	
230	

-32/40-

tatgcacgta gcattttttt ggaaatataa gtggatttgt tcaataaaat attaaatata 838
 aaaaaaaaaa aaaaaaaaaa aa 860

<210> 25
 <211> 873
 <212> DNA
 <213> Renilla reniformis

<220>
 <221> CDS
 <222> (61)...(762)
 <223> GFP Clone-3

<400> 25
 ggcacgagggg tttcctgaca caataaaaac ctttcaaatt gtttctctgt agcagtaagt 60
 atg gat ctc gca aaa ctt ggt ttg aag gaa gtg atg cct act aaa atc 108
 Met Asp Leu Ala Lys Leu Gly Leu Lys Glu Val Met Pro Thr Lys Ile
 1 5 10 15

aac tta gaa gga ctg gtt ggc gac cac gct ttc tca atg gaa gga gtt 156
 Asn Leu Glu Gly Leu Val Gly Asp His Ala Phe Ser Met Glu Gly Val
 20 25 30

ggc gaa ggc aac ata ttg gaa gga act caa gag gtg aag ata tcg gta 204
 Gly Glu Gly Asn Ile Leu Glu Gly Thr Gln Glu Val Lys Ile Ser Val
 35 40 45

aca aaa ggc gca cca ctc cca ttc gca ttt gat atc gta tct gtg gct 252
 Thr Lys Gly Ala Pro Leu Pro Phe Ala Phe Asp Ile Val Ser Val Ala
 50 55 60

ttt tca tat ggg aac aga gct tat acc ggt tac cca gaa gaa att tcc 300
 Phe Ser Tyr Gly Asn Arg Ala Tyr Thr Gly Tyr Pro Glu Glu Ile Ser
 65 70 75 80

gac tac ttc ctc cag tcg ttt cca gaa ggc ttt act tac gag aga aac 348
 Asp Tyr Phe Leu Gln Ser Phe Pro Glu Gly Phe Thr Tyr Glu Arg Asn
 85 90 95

att cgt tat caa gat gga gga act gca att gtt aaa tct gat ata agc 396
 Ile Arg Tyr Gln Asp Gly Gly Thr Ala Ile Val Lys Ser Asp Ile Ser
 100 105 110

ttg gaa gat ggt aaa ttc ata gtg aat gta gac ttc aaa gcg aag gat 444
 Leu Glu Asp Gly Lys Phe Ile Val Asn Val Asp Phe Lys Ala Lys Asp
 115 120 125

cta cgt cgc atg gga cca gtc atg cag caa gac atc gtg ggt atg cag 492
 Leu Arg Arg Met Gly Pro Val Met Gln Gln Asp Ile Val Gly Met Gln
 130 135 140

cca tcg tat gag tca atg tac acc aat gtc act tca gtt ata ggg gaa 540
 Pro Ser Tyr Glu Ser Met Tyr Thr Asn Val Thr Ser Val Ile Gly Glu
 145 150 155 160

tgt ata ata gca ttc aaa ctt caa act ggc aag cat ttc act tac cac 588
 Cys Ile Ile Ala Phe Lys Leu Gln Thr Gly Lys His Phe Thr Tyr His
 165 170 175

-33/40-

```

atg agg aca gtt tac aaa tca aag aag cca gtg gaa act atg cca ttg      636
Met Arg Thr Val Tyr Lys Ser Lys Lys Pro Val Glu Thr Met Pro Leu
180 185 190

tat cat ttc atc cag cat cgc ctc gtt aag acc aat gtg gac aca gcc      684
Tyr His Phe Ile Gln His Arg Leu Val Lys Thr Asn Val Asp Thr Ala
195 200 205

agt ggt tac gtt gtg caa cac gag aca gca att gca gcg cat tct aca      732
Ser Gly Tyr Val Val Gln His Glu Thr Ala Ile Ala Ala His Ser Thr
210 215 220

atc aaa aaa att gaa ggc tct tta cca tag atacctgtac acaattattc      782
Ile Lys Lys Ile Glu Gly Ser Leu Pro *
225 230

tatgcacgta gcattttttt ggaaatataa gtgggtattgt tcaataaaaat attaaatata      842
tgctttttgca aaaaaaaaaa aaaaaaaaaa a
873

<210> 26
<211> 864
<212> DNA
<213> Renilla reniformis

<220>
<221> CDS
<222> (61)...(759)
<223> Human codon optimized Renilla reniformis GFP

<400> 26
ggcagcagggg tttctgaca caataaaaac ctttcaaatt gtttctctgt agcagtaagt      60
atg gac ctg gcc aag ctg ggc ctg aag gag gtg atg ccc acc aag atc      108
Met Asp Leu Ala Lys Leu Gly Leu Lys Glu Val Met Pro Thr Lys Ile
1 5 10 15

aac ctg gag ggc ctg gtg ggc gac cac gcc ttc tcg atg gag ggc gtg      156
Asn Leu Glu Gly Leu Val Gly Asp His Ala Phe Ser Met Glu Gly Val
20 25 30

ggc gag ggc aac atc ttg gag ggc acc cag gag gtg aag atc agc gtg      204
Gly Glu Gly Asn Ile Leu Glu Gly Thr Gln Glu Val Lys Ile Ser Val
35 40 45

acc aag ggc gcc ccc ctg ccc ttc gcc ttc gac atc gtg agc gtg gcc      252
Thr Lys Gly Ala Pro Leu Pro Phe Ala Phe Asp Ile Val Ser Val Ala
50 55 60

ttc agc tac ggc aac cgg gcc tac acc ggc tac ccc gag gag atc agc      300
Phe Ser Tyr Gly Asn Arg Ala Tyr Thr Gly Tyr Pro Glu Glu Ile Ser
65 70 75 80

gac tac ttc ctg cag agc ttc ccc gag ggc ttc acc tac gag cgg aac      348
Asp Tyr Phe Leu Gln Ser Phe Pro Glu Gly Phe Thr Tyr Glu Arg Asn
85 90 95

atc cgg tac cag gac ggc ggc acc gcc atc gtg aag agc gac atc agc      396
Ile Arg Tyr Gln Asp Gly Gly Thr Ala Ile Val Lys Ser Asp Ile Ser
100 105 110

ctg gag gac ggc aag ttc atc gtg aac gtg gac ttc aag gcc aag gac      444

```


-34/40-

Leu Glu Asp Gly Lys Phe Ile Val Asn Val Asp Phe Lys Ala Lys Asp
 115 120 125
 ctg cgg cgg atg ggc ccc gtg atg cag cag gac atc gtg ggc atg cag 492
 Leu Arg Arg Met Gly Pro Val Met Gln Gln Asp Ile Val Gly Met Gln
 130 135 140
 ccc agc tac gag agc atg tac acc aac gtg acc agc gtg atc ggc gag 540
 Pro Ser Tyr Glu Ser Met Tyr Thr Asn Val Thr Ser Val Ile Gly Glu
 145 150 155 160
 tgc atc atc gcc ttc aag ctg cag acc ggc aag cac ttc acc tac cac 588
 Cys Ile Ile Ala Phe Lys Leu Gln Thr Gly Lys His Phe Thr Tyr His
 165 170 175
 atg cgg acc gtg tac aag agc aag aag ccc gtg gag acc atg ccc ctg 636
 Met Arg Thr Val Tyr Lys Ser Lys Lys Pro Val Glu Thr Met Pro Leu
 180 185 190
 tac cac ttc atc cag cac cgg ctg gtg aag acc aac gtg gac acc gcc 684
 Tyr His Phe Ile Gln His Arg Leu Val Lys Thr Asn Val Asp Thr Ala
 195 200 205
 agc ggc tac gtg gtg cag cac gag aca gcc atc gcc gcc cac agc acc 732
 Ser Gly Tyr Val Val Gln His Glu Thr Ala Ile Ala Ala His Ser Thr
 210 215 220
 atc aag aag atc gag ggc agc ctg ccc tagatacctg tacacaatta 779
 Ile Lys Lys Ile Glu Gly Ser Leu Pro
 225 230
 ttctatgcac gtagcatttt tttggaaata taagtggat tgttcaataa aatattaaat 839
 ataaaaaaaa aaaaaaaaaa aaaaa 864
 <210> 27
 <211> 233
 <212> PRT
 <213> Renilla reniformis
 <400> 27
 Met Asp Leu Ala Lys Leu Gly Leu Lys Glu Val Met Pro Thr Lys Ile
 1 5 10 15
 Asn Leu Glu Gly Leu Val Gly Asp His Ala Phe Ser Met Glu Gly Val
 20 25 30
 Gly Glu Gly Asn Ile Leu Glu Gly Thr Gln Glu Val Lys Ile Ser Val
 35 40 45
 Thr Lys Gly Ala Pro Leu Pro Phe Ala Phe Asp Ile Val Ser Val Ala
 50 55 60
 Phe Ser Tyr Gly Asn Arg Ala Tyr Thr Gly Tyr Pro Glu Glu Ile Ser
 65 70 75 80
 Asp Tyr Phe Leu Gln Ser Phe Pro Glu Gly Phe Thr Tyr Glu Arg Asn
 85 90 95
 Ile Arg Tyr Gln Asp Gly Gly Thr Ala Ile Val Lys Ser Asp Ile Ser
 100 105 110
 Leu Glu Asp Gly Lys Phe Ile Val Asn Val Asp Phe Lys Ala Lys Asp
 115 120 125
 Leu Arg Arg Met Gly Pro Val Met Gln Gln Asp Ile Val Gly Met Gln
 130 135 140
 Pro Ser Tyr Glu Ser Met Tyr Thr Asn Val Thr Ser Val Ile Gly Glu
 145 150 155 160
 Cys Ile Ile Ala Phe Lys Leu Gln Thr Gly Lys His Phe Thr Tyr His

-35/40-

[illegible]

```
<210> 28
<211> 861
<212> DNA
<213> Pleuromamma
```

```
<220>
<221> CDS
<222> (148)...(741)
<223> Pleuormamma luciferase
```

```

<400> 28
cggcacgaga ttttgtctgt ggtgattggg attgtctgtc tctcagggtca agcagaaaagt      60
tcgtgtaaag gtgattttctg tagtgatggt tcttctctggg atgtgatcaa gtacaacact      120
gagagtcgac aatgctgtga cacaaaaa atg ctt aga aat tgc gct agg aag caa      174
              Met Leu Arg Asn Cys Ala Arg Lys Gln
              1               5

```

gag caa gtt tgc gcc gat gtg acc gag atg aaa tgc caa gca gtt gct 222
Glu Gln Val Cys Ala Asp Val Thr Glu Met Lys Cys Gln Ala Val Ala
10 15 20 25

tgg gcc gac tgt gga ccc aga ttt gat tcc act ggc agg aat aga tgc 270
Trp Ala Asp Cys Gly Pro Arg Phe Asp Ser Thr Gly Arg Asn Arg Cys
 30 35 40

caa gtt caa tac aag gac tac gcg tac aag tcc tgc gtg gaa gtt gat 318
Gln Val Gln Tyr Lys Asp Tyr Ala Tyr Lys Ser Cys Val Glu Val Asp
45 50 55

tac act gta ccg cac agg aag caa gtt cca gag tgc aaa caa gtc act 366
 Tyr Thr Val Pro His Arg Lys Gln Val Pro Glu Cys Lys Gln Val Thr
 60 65 70

aaa gat aac tgc gtt act gat tgg gaa gtt gac gcc aat ggc aac aag 414
Lys Asp Asn Cys Val Thr Asp Trp Glu Val Asp Ala Asn Gly Asn Lys
75 80 85

gtt tgg ggt ggt acc gag aaa tgc act cct gtc act tgg gaa gaa tgt 462
Val Trp Gly Gly Thr Glu Lys Cys Thr Pro Val Thr Trp Glu Glu Cys
90 95 100 105

aat atc gtg gag aaa gat gta gat ttt cca act gtc aag acg gaa tgc 510
Asn Ile Val Glu Lys Asp Val Asp Phe Pro Thr Val Lys Thr Glu Cys
110 115 120

ggc atc ctg tct cac ctt aag tat gca gac ttc ata gag gga cct tcc 558
Gly Ile Leu Ser His Leu Lys Tyr Ala Asp Phe Ile Glu Gly Pro Ser
125 130 135

cac tct ttg tct atg aga acc aat tgt cag gtc aag agt tca ttg gac 606

-36/40-

His Ser Leu Ser Met Arg Thr Asn Cys Gln Val Lys Ser Ser Leu Asp
 140 145 150
 tgc cgg cct gtt aag acc agg aag tgt gca acg gtc gag tac cac gaa 654
 Cys Arg Pro Val Lys Thr Arg Lys Cys Ala Thr Val Glu Tyr His Glu
 155 160 165
 tgc agc atg aag ccc caa gaa gac tgc agc cca gtc act gtt cat att 702
 Cys Ser Met Lys Pro Gln Glu Asp Cys Ser Pro Val Thr Val His Ile
 170 175 180 185
 cct gac cag gag aaa gtt cac cag aag aag tgc ctc aca taaatgttat 751
 Pro Asp Gln Glu Lys Val His Gln Lys Lys Cys Leu Thr
 190 195
 caatttttagc tcttactaat ttaaacataa taaatatcac atcgaagccc tttattttat 811
 agaagtgttaa tgcttgaata aatctagtga ataaaaaaaa aaaaaaaaaa 861
 <210> 29
 <211> 198
 <212> PRT
 <213> Pleuromamma
 <400> 29
 Met Leu Arg Asn Cys Ala Arg Lys Gln Glu Gln Val Cys Ala Asp Val
 1 5 10 15
 Thr Glu Met Lys Cys Gln Ala Val Ala Trp Ala Asp Cys Gly Pro Arg
 20 25 30
 Phe Asp Ser Thr Gly Arg Asn Arg Cys Gln Val Gln Tyr Lys Asp Tyr
 35 40 45
 Ala Tyr Lys Ser Cys Val Glu Val Asp Tyr Thr Val Pro His Arg Lys
 50 55 60
 Gln Val Pro Glu Cys Lys Gln Val Thr Lys Asp Asn Cys Val Thr Asp
 65 70 75 80
 Trp Glu Val Asp Ala Asn Gly Asn Lys Val Trp Gly Gly Thr Glu Lys
 85 90 95
 Cys Thr Pro Val Thr Trp Glu Glu Cys Asn Ile Val Glu Lys Asp Val
 100 105 110
 Asp Phe Pro Thr Val Lys Thr Glu Cys Gly Ile Leu Ser His Leu Lys
 115 120 125
 Tyr Ala Asp Phe Ile Glu Gly Pro Ser His Ser Leu Ser Met Arg Thr
 130 135 140
 Asn Cys Gln Val Lys Ser Ser Leu Asp Cys Arg Pro Val Lys Thr Arg
 145 150 155 160
 Lys Cys Ala Thr Val Glu Tyr His Glu Cys Ser Met Lys Pro Gln Glu
 165 170 175
 Asp Cys Ser Pro Val Thr Val His Ile Pro Asp Gln Glu Lys Val His
 180 185 190
 Gln Lys Lys Cys Leu Thr
 195
 <210> 30
 <211> 1104
 <212> DNA
 <213> Ptilosarcus gurneyi
 <220>
 <221> CDS
 <222> (34)...(747)
 <223> Ptilosarcus Green Fluorescent Protein

-37/40-

<400> 30

tcggc	cacgag	ctggc	ctcca	cacttt	tagac	aaa	atg	aac	cgc	aac	gta	tta	aag		54	
							Met	Asn	Arg	Asn	Val	Leu	Lys			
							1				5					
aac	act	gga	ctg	aaa	gag	att	atg	tcg	gca	aaa	gct	agc	gtt	gaa	gga	102
Asn	Thr	Gly	Leu	Lys	Glu	Ile	Met	Ser	Ala	Lys	Ala	Ser	Val	Glu	Gly	
		10					15					20				
atc	gtg	aac	aat	cac	gtt	ttt	tcc	atg	gaa	gga	ttt	gga	aaa	ggc	aat	150
Ile	Val	Asn	Asn	His	Val	Phe	Ser	Met	Glu	Gly	Phe	Gly	Lys	Gly	Asn	
		25					30					35				
gta	tta	ttt	gga	aac	caa	ttg	atg	caa	atc	cgg	gtt	aca	aag	gga	ggc	198
Val	Leu	Phe	Gly	Asn	Gln	Leu	Met	Gln	Ile	Arg	Val	Thr	Lys	Gly	Gly	
		40					45				50					
ccg	ttg	cca	ttc	gct	ttc	gat	att	gtt	tcc	ata	gct	ttc	caa	tac	ggg	246
Pro	Leu	Pro	Phe	Ala	Phe	Asp	Ile	Val	Ser	Ile	Ala	Phe	Gln	Tyr	Gly	
				60						65					70	
aat	cgc	act	ttc	acg	aaa	tac	cca	gac	gac	att	gcg	gac	tac	ttt	gtt	294
Asn	Arg	Thr	Phe	Thr	Lys	Tyr	Pro	Asp	Asp	Ile	Ala	Asp	Tyr	Phe	Val	
				75					80							
caa	tca	ttc	ccg	gct	gga	ttt	ttc	tac	gaa	aga	aat	cta	cgc	ttt	gaa	342
Gln	Ser	Phe	Pro	Ala	Gly	Phe	Phe	Tyr	Glu	Arg	Asn	Leu	Arg	Phe	Glu	
			90					95					100			
gat	ggc	gcc	att	gtt	gac	att	cgt	tca	gat	ata	agt	tta	gaa	gat	gat	390
Asp	Gly	Ala	Ile	Val	Asp	Ile	Arg	Ser	Asp	Ile	Ser	Leu	Glu	Asp	Asp	
		105					110					115				
aag	ttc	cac	tac	aaa	gtg	gag	tat	aga	ggc	aac	ggc	ttc	cct	agt	aac	438
Lys	Phe	His	Tyr	Lys	Val	Glu	Tyr	Arg	Gly	Asn	Gly	Phe	Pro	Ser	Asn	
					125					130					135	
gga	ccc	gtg	atg	caa	aaa	gcc	atc	ctc	ggc	atg	gag	cca	tcg	ttt	gag	486
Gly	Pro	Val	Met	Gln	Lys	Ala	Ile	Leu	Gly	Met	Glu	Pro	Ser	Phe	Glu	
				140						145					150	
gtg	gtc	tac	atg	aac	agc	ggc	gtt	ctg	gtg	ggc	gaa	gta	gat	ctc	gtt	534
Val	Val	Tyr	Met	Asn	Ser	Gly	Val	Leu	Val	Gly	Glu	Val	Asp	Leu	Val	
			155					160					165			
tac	aaa	ctc	gag	tca	ggg	aac	tat	tac	tcg	tgc	cac	atg	aaa	acg	ttt	582
Tyr	Lys	Leu	Glu	Ser	Gly	Asn	Tyr	Tyr	Ser	Cys	His	Met	Lys	Thr	Phe	
		170					175					180				
tac	aga	tcc	aaa	ggc	gga	gtg	aaa	gaa	ttc	ccg	gaa	tat	cac	ttt	atc	630
Tyr	Arg	Ser	Lys	Gly	Gly	Val	Lys	Glu	Phe	Pro	Glu	Tyr	His	Phe	Ile	
		185					190				195					

-38/40-

ggc tcc ctt cat gaa tgg gtg tagaaaatga ccaatatact ggggaaaccg 777
 Gly Ser Leu His Glu Trp Val
 235

ataaccgttt ggaagcttgt gtatacaaat tatttggggt cattttgtaa tgtgtatgtg 837
 tgttgtatga tcaatagacg tcgtcattca tagcttgaat ccttcagcaa aagaaacctc 897
 gaagcatatt gaaacctcga agcatattga aacctcgacg gagagcgtaa agagaccgca 957
 caaattaacg cgtttcaacc agcagtttga atctttaaac cgatcaaaac tattaatata 1017
 aatatatata ccctgtataa cttatatata tctatatagt ttgatattga ttaaattctgt 1077
 tcttgatcaa aaaaaaaaaa aaaaaaa 1104

<210> 31
 <211> 1279
 <212> DNA
 <213> *Ptilosarcus gurneyi*

<220>
 <221> CDS
 <222> (7)...(720)
 <223> *Ptilosarcus* Green Flourescent Protein (GFP)

<400> 31
 gacaaa atg aac cgc aac gta tta aag aac act gga ctg aaa gag att 48
 Met Asn Arg Asn Val Leu Lys Asn Thr Gly Leu Lys Glu Ile
 1 5 10

atg tcg gca aaa gct agc gtt gaa gga atc gtg aac aat cac gtt ttt 96
 Met Ser Ala Lys Ala Ser Val Glu Gly Ile Val Asn Asn His Val Phe
 15 20 25 30

tcc atg gaa gga ttt gga aaa ggc aat gta tta ttt gga aac caa ttg 144
 Ser Met Glu Gly Phe Gly Lys Gly Asn Val Leu Phe Gly Asn Gln Leu
 35 40 45

atg caa atc cgg gtt aca aag gga ggt ccg ttg cca ttc gct ttc gac 192
 Met Gln Ile Arg Val Thr Lys Gly Gly Pro Leu Pro Phe Ala Phe Asp
 50 55 60

att gtt tcc ata gct ttc caa tac ggg aat cgc act ttc acg aaa tac 240
 Ile Val Ser Ile Ala Phe Gln Tyr Gly Asn Arg Thr Phe Thr Lys Tyr
 65 70 75

cca gac gac att gcg gac tac ttt gtt caa tca ttt ccg gct gga ttt 288
 Pro Asp Asp Ile Ala Asp Tyr Phe Val Gln Ser Phe Pro Ala Gly Phe
 80 85 90

ttc tac gaa aga aat cta cgc ttt gaa gat ggc gcc att gtt gac att 336
 Phe Tyr Glu Arg Asn Leu Arg Phe Glu Asp Gly Ala Ile Val Asp Ile
 95 100 105 110

cgt tca gat ata agt tta gaa gat gat aag ttc cac tac aaa gtg gag 384
 Arg Ser Asp Ile Ser Leu Glu Asp Asp Lys Phe His Tyr Lys Val Glu
 115 120 125

tat aga ggc aac ggt ttc cct agt aac gga ccc gtg atg caa aaa gcc 432
 Tyr Arg Gly Asn Gly Phe Pro Ser Asn Gly Pro Val Met Gln Lys Ala
 130 135 140

atc ctc ggc atg gag cca tcg ttt gag gtg gtc tac atg aac agc ggc 480
 Ile Leu Gly Met Glu Pro Ser Phe Glu Val Val Tyr Met Asn Ser Gly

-39/40-

145	150	155	
gtt ctg gtg ggc gaa gta gat ctc gtt tac aaa ctc gag tca ggg aac			528
Val Leu Val Gly Glu Val Asp Leu Val Tyr Lys Leu Glu Ser Gly Asn			
160	165	170	
tat tac tcg tgc cac atg aaa acg ttt tac aga tcc aaa ggt gga gtg			576
Tyr Tyr Ser Cys His Met Lys Thr Phe Tyr Arg Ser Lys Gly Gly Val			
175	180	185	190
aaa gaa ttc ccg gaa tat cac ttt atc cat cat cgt ctg gag aaa acc			624
Lys Glu Phe Pro Glu Tyr His Phe Ile His Arg Leu Glu Lys Thr			
195	200	205	
tac gtg gaa gaa gga agc ttc gtg gaa caa cac gag acg gcc att gca			672
Tyr Val Glu Glu Gly Ser Phe Val Glu Gln His Glu Thr Ala Ile Ala			
210	215	220	
caa ctg acc aca att gga aaa cct ctg ggc tcc ctt cat gaa tgg gtg			720
Gln Leu Thr Thr Ile Gly Lys Pro Leu Gly Ser Leu His Glu Trp Val			
225	230	235	
tagaaaaatga ccaatatact ggggaaaatc accaatatac tggggaaaat gaccaattta			780
ctggggaaaaa tgaccaatat actgtagaaa atcaccaata tactggggaa aatgaccaat			840
ttactggggga aatgaccaat ttactgtaga aaatcaccaa tatactgtgg aaaatgacca			900
aaatactgta gaaatgttca cactgggttg ataaccgttt cgataaccgt ttggaagctt			960
gtgtatacaa gttatttggg gtcattttgt aatgtgtatg tgtgttgtat gatctataga			1020
cgctgctatt catagcttga atccttcagc aaaagaaacc tcgaagcata ttgaaacctc			1080
gacggagagc ataaagagac cgcacgtaca caaattataa taccagcagt tggaatcttt			1140
aaaccgatca aaactattaa tatatatata caccctgtat aacatatata tatatatata			1200
tctacatagt ttgatattga ttaaactctgt tcttgatcac taaaaaaaaa aaaaaaaaaa			1260
aaaaaaaaaa aaaaaaaaaa			1279

<210> 32

<211> 238

<212> PRT

<213> *Ptilosarcus gurneyi*

<400> 32

Met Asn Arg Asn Val Leu Lys Asn Thr Gly Leu Lys Glu Ile Met Ser	
1 5 10 15	
Ala Lys Ala Ser Val Glu Gly Ile Val Asn Asn His Val Phe Ser Met	
20 25 30	
Glu Gly Phe Gly Lys Gly Asn Val Leu Phe Gly Asn Gln Leu Met Gln	
35 40 45	
Ile Arg Val Thr Lys Gly Gly Pro Leu Pro Phe Ala Phe Asp Ile Val	
50 55 60	
Ser Ile Ala Phe Gln Tyr Gly Asn Arg Thr Phe Thr Lys Tyr Pro Asp	
65 70 75 80	
Asp Ile Ala Asp Tyr Phe Val Gln Ser Phe Pro Ala Gly Phe Phe Tyr	
85 90 95	
Glu Arg Asn Leu Arg Phe Glu Asp Gly Ala Ile Val Asp Ile Arg Ser	
100 105 110	
Asp Ile Ser Leu Glu Asp Asp Lys Phe His Tyr Lys Val Glu Tyr Arg	
115 120 125	
Gly Asn Gly Phe Pro Ser Asn Gly Pro Val Met Gln Lys Ala Ile Leu	
130 135 140	
Gly Met Glu Pro Ser Phe Glu Val Val Tyr Met Asn Ser Gly Val Leu	
145 150 155 160	
Val Gly Glu Val Asp Leu Val Tyr Lys Leu Glu Ser Gly Asn Tyr Tyr	
165 170 175	

-40/40-

Ser Cys His Met Lys Thr Phe Tyr Arg Ser Lys Gly Gly Val Lys Glu
 180 185 190
 Phe Pro Glu Tyr His Phe Ile His His Arg Leu Glu Lys Thr Tyr Val
 195 200 205
 Glu Glu Gly Ser Phe Val Glu Gln His Glu Thr Ala Ile Ala Gln Leu
 210 215 220
 Thr Thr Ile Gly Lys Pro Leu Gly Ser Leu His Glu Trp Val
 225 230 235

<210> 33

<211> 233

<212> PRT

<213> Renilla Reniformis mutein

<400> 33

Met Asp Leu Ala Lys Leu Gly Leu Lys Glu Val Met Pro Thr Lys Ile
 1 5 10 15
 Asn Leu Glu Gly Leu Val Gly Asp His Ala Phe Ser Met Glu Gly Val
 20 25 30
 Gly Glu Gly Asn Ile Leu Glu Gly Thr Gln Glu Val Lys Ile Ser Val
 35 40 45
 Thr Lys Gly Ala Pro Leu Pro Phe Ala Phe Asp Ile Val Ser Val Ala
 50 55 60
 Phe Ser Tyr Gly Asn Arg Ala Tyr Thr Gly Tyr Pro Glu Glu Ile Ser
 65 70 75 80
 Asp Tyr Phe Leu Gln Ser Phe Pro Glu Gly Phe Thr Tyr Glu Arg Asn
 85 90 95
 Ile Arg Tyr Gln Asp Gly Gly Thr Ala Ile Val Asp Ser Asp Ile Ser
 100 105 110
 Leu Glu Asp Gly Lys Phe Ile Val Asn Val Asp Phe Lys Ala Asp Asp
 115 120 125
 Leu Arg Asp Met Gly Pro Val Met Gln Gln Asp Ile Val Gly Met Gln
 130 135 140
 Pro Ser Tyr Glu Ser Met Tyr Thr Asn Val Thr Ser Val Ile Gly Glu
 145 150 155 160
 Cys Ile Ile Ala Phe Lys Leu Gln Thr Gly Lys Asp Phe Thr Tyr His
 165 170 175
 Met Arg Thr Val Tyr Lys Ser Lys Lys Pro Val Glu Thr Met Pro Leu
 180 185 190
 Tyr His Phe Ile Gln His Asp Leu Val Lys Thr Asn Val Asp Thr Ala
 195 200 205
 Ser Gly Tyr Val Val Gln His Glu Thr Ala Ile Ala Ala His Ser Thr
 210 215 220
 Ile Asp Lys Ile Glu Gly Ser Leu Pro
 225 230

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68824 A3

(51) International Patent Classification⁷: C12N 9/02, C07K 14/435, C12N 15/12, 15/10, 15/11, 15/66, A61K 49/00, A01H 5/00, A01K 67/027, A61K 38/17, F41C 3/00, F21S 10/00, A23L 1/00, C12G 1/00, C07K 16/18, C12N 5/10

Pittsburgh, PA 15237 (US). **SZCZEPANIAK, William** [US/US]; 372 South Highland Avenue #203, Pittsburgh, PA 15206 (US).

(21) International Application Number: PCT/US01/08277

(74) Agent: **TOWNER, Alan, G.**; Pietragallo, Bosick & Gordon, One Oxford Centre, 38th Floor, 301 Grant Street, Pittsburgh, PA 15219 (US).

(22) International Filing Date: 15 March 2001 (15.03.2001)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(26) Publication Language: English

(30) Priority Data:
60/189,691 15 March 2000 (15.03.2000) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/189,691 (CON)
Filed on 15 March 2000 (15.03.2000)

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **PROLUME, LTD.** [US/US]; 1085 William Pitt Way, Pittsburgh, PA 15236 (US).

Published:

— with international search report

(71) Applicant and

(72) Inventor: **BRYAN, Bruce, J.** [US/US]; 716 N. Arden Drive, Beverly Hills, CA 90210 (US).

(88) Date of publication of the international search report:
2 May 2002

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **SZENT-GY-ORGYI, Christopher** [US/US]; 719 Duncan Avenue.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: RENILLA RENIFORMIS FLUORESCENT PROTEINS, NUCLEIC ACIDS ENCODING THE FLUORESCENT PROTEINS AND THE USE THEREOF IN DIAGNOSTICS, HIGH THROUGHPUT SCREENING AND NOVELTY ITEMS

(57) Abstract: Isolated and purified nucleic acids encoding green fluorescent proteins from *Renilla reniformis* and the green fluorescent protein encoded thereby are also provided. Mutants of the nucleic acid molecules and the modified encoded proteins are also provided. Compositions and combinations comprising the green fluorescent proteins and/or the luciferase are further provided.

WO 01/68824 A3

INTERNATIONAL SEARCH REPORT

Int. l. Application No

PCT/US 01/08277

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N9/02 C07K14/435 C12N15/12 C12N15/10 C12N15/11
 C12N15/66 A61K49/00 A01H5/00 A01K67/027 A61K38/17
 F41C3/00 F21S10/00 A23L1/00 C12G1/00 C07K16/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EMBL, WPI Data, PAJ, EPO-Internal, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 49019 A (SZENT GYORGYI CHRISTOPHER ;BRYAN BRUCE J (US); PROLUME LTD (US)) 30 September 1999 (1999-09-30) cited in the application the whole document ---	1-77
X	WO 98 14605 A (UNIV LOMA LINDA) 9 April 1998 (1998-04-09) the whole document ---	38,39, 41,42, 44,47-56
X	WO 92 15673 A (UNIV GEORGIA) 17 September 1992 (1992-09-17) page 4, line 20-35 figures 1,2 ---	41
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

5 October 2001

Date of mailing of the international search report

15/10/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
 Fax: (+31-70) 340-3016

Authorized officer

Mateo Rosell, A.M.

INTERNATIONAL SEARCH REPORT

Int. :ional Application No

PCT/US 01/08277

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N5/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
<p>X WARD W W ET AL: "AN ENERGY TRANSFER PROTEIN IN COELENTERATE BIO LUMINESCENCE CHARACTERIZATION OF THE RENILLA-RENIFORMIS GREEN FLUORESCENT PROTEIN" JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 254, no. 3, 1979, pages 781-788, XP002935261 EN ISSN: 0021-9258 the whole document</p>	1-14
<p>X US 5 876 995 A (BRYAN BRUCE) 2 March 1999 (1999-03-02) cited in the application the whole document</p>	23-35, 75-77

-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

5 October 2001

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Mateo Rosell, A.M.

INTERNATIONAL SEARCH REPORT

Int lional Application No

PCT/US 01/08277

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 26277 A (BRYAN BRUCE J ;PROLUME LTD (US); GAALEMA STEPHEN (US); MURPHY RAND) 18 June 1998 (1998-06-18) cited in the application the whole document	70-74
A	FRATAMICO P M ET AL: "Construction and characterization of Escherichia coli 0157:H7 strains expressing firefly luciferase and green fluorescent protein and their use in survival studies" JOURNAL OF FOOD PROTECTION, DES MOINES, IO, US, vol. 60, no. 10, 1997, pages 1167-1173, XP002118033 ISSN: 0362-028X the whole document	38
A	XU Y ET AL: "A bioluminescence resonance energy transfer (BRET) system: application to interacting circadian clock proteins" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, US, vol. 96, no. 1, 5 January 1999 (1999-01-05), pages 151-156, XP002118035 ISSN: 0027-8424 the whole document	64-69
A	HEIM R ET AL: "ENGINEERING GREEN FLUORESCENT PROTEIN FOR IMPROVED BRIGHTNESS, LONGER WAVELENGTHS AND FLUROESCENCE RESONANCE ENERGY TRANSFER" CURRENT BIOLOGY, CURRENT SCIENCE,, GB, vol. 6, no. 2, 1 February 1996 (1996-02-01), pages 178-182, XP000676582 ISSN: 0960-9822 the whole document	1,15,16, 20
A	EHRIG T ET AL: "GREEN-FLUORESCENT PROTEIN MUTANTS WITH ALTERED FLUORENCE EXCITATIONSPECTRA" FEBS LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 367, 1995, pages 163-166, XP000579119 ISSN: 0014-5793 the whole document	15,16,20
E	WO 01 32688 A (UNIV RUTGERS ;WARD WILLIAM W (US)) 10 May 2001 (2001-05-10) the whole document	1-22,36, 37,64-69

INTERNATIONAL SEARCH REPORT

Information on patent family members

Int: ional Application No

PCT/US 01/08277

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9949019	A	30-09-1999	AU 3210399 A	18-10-1999
			EP 1064360 A2	03-01-2001
			WO 9949019 A2	30-09-1999
			US 6232107 B1	15-05-2001
WO 9814605	A	09-04-1998	US 5976796 A	02-11-1999
			AU 730040 B2	22-02-2001
			AU 4500497 A	24-04-1998
			EP 0934425 A1	11-08-1999
			JP 2001501100 T	30-01-2001
			WO 9814605 A1	09-04-1998
WO 9215673	A	17-09-1992	WO 9215673 A1	17-09-1992
			DE 69131780 D1	16-12-1999
			DE 69131780 T2	16-11-2000
			DK 575319 T3	10-07-2000
			EP 0575319 A1	29-12-1993
			ES 2142801 T3	01-05-2000
			GR 3032608 T3	31-05-2000
			JP 3126980 B2	22-01-2001
			JP 6504903 T	09-06-1994
US 5876995	A	02-03-1999	US 6247995 B1	19-06-2001
			AU 2255597 A	28-08-1997
			CA 2245594 A1	14-08-1997
			EP 0879383 A2	25-11-1998
			JP 11504822 T	11-05-1999
			WO 9729319 A2	14-08-1997
			US 6152358 A	28-11-2000
			US 6113886 A	05-09-2000
WO 9826277	A	18-06-1998	AU 6012898 A	03-07-1998
			EP 1015872 A2	05-07-2000
			JP 2001511887 T	14-08-2001
			WO 9826277 A2	18-06-1998
WO 0132688	A	10-05-2001	AU 1446801 A	14-05-2001
			WO 0132688 A1	10-05-2001

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.